

Chiral magnetic effect search in p+Au, d+Au and Au+Au collisions at RHIC


Jie Zhao (for the STAR collaboration)
Sep. 15 2017

Purdue University, West Lafayette

- **Chiral Magnetic Effect (CME)**
- **CME in small systems**
- **RHIC-STAR experiment**
- **Results in p/d+A and A+A collisions**
- **Identification of backgrounds and possible CME**
- **Summary**

Chiral Magnetic Effect (CME)

D. Kharzeev, etc. NPA 803, 227(2008)



$$j_V = \frac{N_c e}{2\pi^2} \mu_A B, \quad \Rightarrow \quad \text{electric charge separation along the B field}$$

Configuration with non-zero topological charge (Q_w) converts left (right)-handed fermions to right (left)-handed fermions, generating electric current along B direction and leading to electric charge separation

STAR Charged-Particle Azimuthal Correlations


S. A. Voloshin, Phys.Rev. C 70 (2004) 057901

Particle distribution effectively can be described by:

$$\frac{dN_{\pm}}{d\phi} \propto (1 + 2v_1 \cos(\Delta\phi) + 2v_2 \cos(2\Delta\phi) + \dots + 2a_{\pm} \sin(\Delta\phi)), \quad (1)$$

$a > 0$ preferential emission along the angular momentum, $a_+ = -a_-$.

The sign of Q_w **can vary** event to event and domain to domain \rightarrow one has to measure correlations, $\langle a_{\alpha} a_{\beta} \rangle$, P-even quantity (!) -- possibility of contribution from effects not related to P-violation



slide from S. A. Voloshin

Predictions:

$$a_+ = -a_-$$


$$a_+ a_+ = a_- a_- = -a_+ a_-$$

$a \sim 10^{-2}$ for midcentral coll-ns

$$\begin{aligned} \langle \cos(\phi_{\alpha} + \phi_{\beta} - 2\Psi_{RP}) \rangle &= \\ &= \langle \cos(\phi_{\alpha} - \Psi_{RP}) \cos(\phi_{\beta} - \Psi_{RP}) \rangle - \langle \sin(\phi_{\alpha} - \Psi_{RP}) \sin(\phi_{\beta} - \Psi_{RP}) \rangle \\ &\approx (v_{1,\alpha} v_{1,\beta} - a_{\alpha} a_{\beta}) \end{aligned}$$

Harmonic planes in small systems

CMS collaboration, PRL 118(2017)122301; R. Belmont and J.L. Nagle, arXiv:1610.07964v1




Ψ_2 : second order event plane; Ψ_1 : first order event plane



- Ψ_2 related to flow, related to -> flow background
- Ψ_1 related to the magnetic direction (B), useful for -> CME signal
- Ψ_1 and Ψ_2 correlated in A+A, signal and background entangled
- Ψ_1 and Ψ_2 not correlated in p+A, d+A, signal and background disentangled

Charge dependent signal at RHIC


STAR collaboration, PRL 103(2009)251601; PRC 81(2010)54908; PRC 88 (2013) 64911



- Correlator indicates charge dependent signal
- Consistent between different years (2004 and 2007)
- Consistent with the 1st-order EP (from spectator neutron v_1)



- “The observed signal as functions of multiplicity and η gap, are of similar magnitude in p+Pb and Pb+Pb collisions at the same multiplicities”
- “The results pose a challenge for the interpretation of charge-dependent azimuthal correlations in heavy-ion collisions in terms of the CME”
- There might be energy dependent difference, more studies at RHIC are needed



RHIC

PHENIX

STAR

LINAC




NSRL

EBIS



BOOSTER

AGS

The STAR detector




- **Time Projection Chamber** ($0 < \phi \leq 2\pi$, $|\eta| < 1$)
 - Tracking – momentum
 - Ionization energy loss – dE/dx (particle identification)
- **Time Of Flight detector** ($0 < \phi \leq 2\pi$, $|\eta| < 0.9$)
 - Timing resolution $< 100\text{ps}$ - significant improvement of PID



$N(\alpha/\beta)$ represents the charged(+/-) particle multiplicity used for the correlator

- Sizeable charge dependent signal in small system p+Au and d+Au collisions with respect to second order event plane Ψ_2
- $v_2\{2\}$ with η gap of 1.0


STAR Rapidity-gap dependence in small systems




- Correlator as a function of the η gap between the two charged particles in p+Au, d+Au and peripheral Au+Au collisions
- Peripheral Au+Au data are similar to those of p+Au and d+Au

Multiplicity dependence

$v_2\{2\}$ with eta gap of 1.0



$N^* < \cos(\phi_\alpha + \phi_\beta - 2\phi_c) > / v_2^2\{2\}$ OS-SS





- Background expectation: N dilution, proportional to flow $v_2\{2\}$
- Right plot: if intrinsic particle pair-wise correlation is independent of N , background scenario would yield a constant as a function of N
- With topological charge sign fluctuations and magnetic field direction fluctuations, **CME might yield different multiplicity dependence**

Identify the backgrounds

Resonance decay background

Fuqiang Wang, Jie Zhao, Phys.Rev.C 95,051901(R) (2017)

STAR, Phys.Rev.Lett.92,092301 (2004)




Resonance decay:

$$\Delta\gamma = \cos(\alpha + \beta - 2\Psi_{RP})$$



$$\propto \langle \cos(\alpha + \beta - 2\phi_\rho) \rangle \cos 2(\phi_\rho - \Psi_{RP})$$

$$\approx \langle \cos(\alpha + \beta - 2\phi_\rho) \rangle v_{2,\rho}$$

Resonance bkg.:
resonance decay
coupled with v_2 ,
will give a CME
-like $\Delta\gamma$ signal




AMPT simulation




$b: 6.8-8.2$ fm
 $\Delta\gamma: (81.2 \pm 1.2)E-6$
 $m > 2 \text{ GeV}/c^2: (-5.8 \pm 8.1)E-6$

- AMPT has **no CME, only background**
- AMPT show resonance structure in $\Delta\gamma$ as function of mass
- At large mass with smaller abundance difference between the unlike-sign and like-sign pairs, $\Delta\gamma$ consistent with zero

Identify resonance bkg. by **invariant mass**




STAR data





- Identify resonance background by the **invariant mass**
- **Data show resonance structure in $\Delta\gamma$ as function of mass**
- $\Delta\gamma$ decrease as r decrease with m increase, larger $r \rightarrow$ larger res. contribution \rightarrow larger $\Delta\gamma$

$\Delta\gamma$ traces r

Go to high mass region




Centrality	All (A)	M>1.5 (B)	B/A
50-80%	(7.45±0.21)E-4	(1.3±5.7)E-5	(1.8±7.6)%
20-50%	(1.82±0.03)E-4	(7.7±9.0)E-6	(4.3±4.9)%
0-20%	(3.70±0.67)E-5	(1.4±1.8)E-5	(-3.8±49)%



- Decreasing resonance contribution to unlike-sign pair excess with increasing mass
- At $m > 1.5 \text{ GeV}/c^2$, $\Delta\gamma$ consistent with zero

Identify resonance bkg. and possible CME



➤ Resonance bkg.:

$$\Delta\gamma = \cos(\alpha + \beta - 2\Psi_{RP})$$

$$\infty \langle \cos(\alpha + \beta - 2\phi_{res.}) \rangle \cos 2(\phi_{res.} - \Psi_{RP})$$

$$\approx \langle \cos(\alpha + \beta - 2\phi_{res.}) \rangle v_{2res.}$$

➤ Resonance bkg. + CME

$$\Delta\gamma(m) = r(m) * \cos(\alpha + \beta - 2\phi_{res.}) * v_{2res.} + \text{CME}$$

$$= r(m) * f(m) * v_{2res.} + \text{CME}$$



peaky



smooth

different dep. on mass, viable way to distinguish

Default $\Delta\gamma$	$(1.82 \pm 0.03)E-4$	
	constant CME	exponential CME
	$(5.9 \pm 9.0)E-6$	$(3.0 \pm 2.0)E-5$
Ratio to default:	$3.2 \pm 4.9\%$	$16 \pm 11\%$

- Data fitted with a constant CME, and a CME exponential in mass
- In the current approach, the statistical uncertainty is dominant

Summary

- With respect to Ψ_2 : p+Au and d+Au charge dependent correlations are background. Peripheral Au+Au data are similar to that of p+Au and d+Au
- The scaled correlators from peripheral to mid-central Au+Au collisions are approximately constant over multiplicity. **These data do not currently allow conclusive statements to be made regarding the presence of the CME**
- Identify resonance bkg. by the **invariant mass**
- At $m > 1.5 \text{ GeV}/c^2$, $\Delta\gamma$ is consistent with zero within uncertainty
- Observation of resonance structure in $\Delta\gamma$ at $m < 1.5 \text{ GeV}/c^2$.
Two component fit is used to isolate the possible CME from bkg.