

# Dependence of $J/\psi$ production on Charged-Particle Multiplicity in p+p Collisions at $\sqrt{s}=200$ GeV with the STAR Experiment

8<sup>th</sup> International Workshop on Multiple Parton Interactions at the LHC, 2016





Zhenyu Ye<sup>1,2</sup> (for the STAR collaboration)

1. University of Illinois at Chicago
2. Central China Normal University


# Motivation

- Quarkonium production mechanisms in p+p collisions are not fully understood
  - Hard processes for heavy quark pair production - pQCD
  - Non-perturbative soft processes for hadronization – CEM, CSM, COM
- More differential and comprehensive studies at different energies may shed new lights on quarkonium production and QCD
  - production cross-section
  - polarization
  - yield vs event activity ( $N_{ch}$ )
  - ...
- Quarkonium yield vs event activity can provide insights into multiple parton interactions



Courtesy: Wei Yang Wang

# Open Charm Yield vs $N_{ch}$ at LHC




Faster-than-linear rise of open charm production vs  $N_{ch}$  in p+p @ 7 TeV

- PYTHIA8:** including MPI:  


$$N_{hard} \propto N_{ch} \propto N_{MPI}$$
- EPOS3:** use Gribov-Regge multiple parton scattering for initial conditions:  

$$N_{hard} \propto N_{ch} \propto N_{MPI}$$
- EPOS3+Hydro:** energy density in 7 TeV p+p collisions is high enough to apply hydrodynamic evolution to the core of the collisions.  
 **$N_{hard}$  rises faster than  $N_{ch}$**  in certain  $p_T$  range
- Percolation model:** exchange color sources in collisions. High energy density suppresses soft process more than hard process  
 **$N_{hard}$  rises faster than  $N_{ch}$**

# Relativistic Heavy Ion Collider



# STAR Experiment at RHIC



Magnet

TPC

TOF




BEMC

VPD

BBC



TPC/TOF/BEMC: full azimuthal acceptance in  $|\eta| < 1$

# Electron Identification




$$n\sigma_{dE/dx}^e = \frac{\ln(dE/dx_{\text{measured}}) - \ln(dE/dx_{\text{exp}})}{R[\ln(dE/dx_{\text{measured}})]}$$

# J/ψ Reconstruction



- Trigger: VPD for MB, BEMC for HT0(2) with  $E_T > \sim 2.5$  (4.3) GeV/c
- HT0(2): triggered electron  $p_T > 2.5$  (4.3) GeV/c
- Electron ID: TPC+EMC or TPC+TOF
- $2.9 < m_{ee} < 3.2$  GeV/c<sup>2</sup>,  $-1 < y_{ee} < 1$

# J/ψ Cross-section in p+p@200 GeV



STAR 2012 data preliminary

$$B_{ee} \frac{d\sigma_{J/\psi}}{dy} = 47.3 \pm 2.9 \pm 6.1 \pm 3.8 \text{ nb}$$

PHENIX Publication

$$B_{ee} \frac{d\sigma_{J/\psi}}{dy} = 42.5 \pm 1.4 \pm 4.8 \pm 3.1 \text{ nb}$$

- Consistent with PHENIX result; Better precision at  $p_T > 2 \text{ GeV}/c$
- NRQCD describes data fairly well; Small tension at  $p_T < 1.5 \text{ GeV}/c$  with CGC+NRQCD


PHENIX: PRD **82** (2010) 012001

NLO NRQCD: Ma et al., PRL **106** (2011) 042002

CGC+NRQCD: Ma, Venugopalan, PRL **113** (2014) 192301



# Measuring $N_{ch}$ at STAR



- Reconstruct charged particles with  $p_T > 0.2$  GeV/c and  $|\eta| < 1$  with  $\geq 20$  TPC hits.
- “TPC-only” tracks receive large contribution from pile-up collisions, which can be suppressed by requiring tracks match to fast “TOF” hits.
- Dependences of pile-up effect, TPC and TOF efficiencies on instantaneous luminosity ( $zdcX$ ) need to be taken into account.

TPC raw: detector-level TPC-only  $N_{ch}$


TOF raw: detector-level TPC+TOF  $N_{ch}$

TPC cor: TPC-only  $N_{ch}$  corrected by TPC efficiency

TOF cor: TPC+TOF  $N_{ch}$  corrected by lumi-independent TPC+TOF efficiency



TOF eff weight: TPC+TOF  $N_{ch}$  corrected by lumi-dependent TPC+TOF efficiency

# Measuring $N_{ch}$ at STAR




- Charged particles with  $p_T > 0.2 \text{ GeV}/c$  and  $|\eta| < 1$  are reconstructed by the TPC and matched with TOF hits in NSD-enhanced ( $BBC 2.2 < |\eta| < 5.2$ ) events.
- Raw  $N_{ch}$  distributions are unfolded to  $p_T > 0 \text{ GeV}/c$  and  $|\eta| < 1$  (corrected for detector efficiency and secondary particles from weak decays)

# $N_{ch}$ in p+p Collisions at 200 GeV




- Unfolded  $N_{ch}$  distribution can be described by negative binomial distribution.
- $dN_{ch}/d\eta = 3.13 \pm 0.27$  consistent with previous STAR result  $2.98 \pm 0.34$

# J/ψ Yield vs $N_{ch}$ in p+p @ 200 GeV




- Unfolded  $N_{ch}$  distribution for J/ψ events can not be described by PYTHIA8.  $\langle N_{ch} \rangle$  in data higher than that in PYTHIA8 for J/ψ production.

# J/ψ Yield vs $N_{ch}$ in p+p @ 200 GeV



- Unfolded  $N_{ch}$  distribution for J/ψ events can not be described by PYTHIA8.  $\langle N_{ch} \rangle$  in data higher than that in PYTHIA8 for J/ψ production.
- Relative J/ψ yield vs  $N_{ch}$  increases faster than linear. Such an increase is qualitatively described by PYTHIA8 and Percolation model but the increase is underestimated.

# J/ψ Yield vs Event Activity ( $N_{ch}$ )



Faster-than-linear rise of open charm and J/ψ production vs  $N_{ch}$  in p+p @ 7 TeV

- **Percolation model:** exchange color sources in collisions. High energy density suppresses soft processes more than hard processes

$N_{hard}$  rises faster than  $N_{ch}$  at LHC


- **EPOS3+Hydro:** energy density in 7 TeV p+p collisions is high enough to apply hydrodynamic evolution to the core of the collisions

$N_{hard}$  rises faster than  $N_{ch}$  at LHC

- **PYTHIA8 and EPOS3:** with MPI underestimate the increase

$$N_{hard} \propto N_{ch} \propto N_{MPI}$$

# J/ψ Yield vs Event Activity ( $N_{ch}$ )



- Percolation model:** exchange color sources in collisions. High energy density suppresses soft processes more than hard processes  
 $N_{hard}$  rises faster than  $N_{ch}$  at LHC  
 Small collisional energy dependence  
 $N_{hard}$  rises faster than  $N_{ch}$  at RHIC
- EPOS3+Hydro:** energy density in 7 TeV p+p collisions is high enough to apply hydrodynamic evolution to the core of the collisions  
 $N_{hard}$  rises faster than  $N_{ch}$  at LHC  
 Strong collision energy dependence  
 $\langle dN_{ch}/d\eta \rangle \sim 3$  at 200 GeV  
 $\sim 6$  at 7 TeV  
 $N_{hard}$  rises linearly as  $N_{ch}$  at RHIC

Stronger-than-linear rise following the same trend at 200 GeV and 7 TeV, suggests not a hot medium effect assumed in EPOS3+Hydro for p+p collisions

# Summary and Outlook

$J/\psi$  yield vs event activity ( $N_{\text{ch}}$ ) in p+p collisions at 200 GeV has been studied at STAR and compared to LHC results at 7 TeV

- Strong and similar correlation between  $J/\psi$  production and  $N_{\text{ch}}$  at 0.2 and 7 TeV
- PYTHIA8 qualitatively describes RHIC and LHC data but underestimates at high  $N_{\text{ch}}$
- EPOS3 with MPI predicts linear increase and underestimates the data
- EPOS3+Hydro and Percolation model predict faster-than-linear increase, but the former expects strong dependence on  $\sqrt{s}$ , inconsistent with RHIC data.


More precise results from significantly increased data size and upgraded STAR detectors are expected.

- Improved low  $p_{\text{T}}$   $J/\psi$  precision by Muon Telescope Detector
- Open heavy flavor and Upsilon production, p+p at 500 GeV



# Backup

# $N_{ch}$ Unfolding



# $N_{ch}$ Unfolding

