

Deutsches Elektronen-Synchrotron
(DESY), Hamburg




Monte Carlo tuning and development in the CMS collaboration

[Paolo Gunnellini](#)

on behalf of the CMS Collaboration




MPI@LHC 2016
San Cristobal de las Chiapas
November 2016, Mexico




- Brief introduction
- Before and during Run II situation
- Considered measurements for CMS tunes
- Tuning value of α_S
- Validation plots at 13 TeV
- Herwig7 and Sherpa tuning
- Conclusions

The underlying event at the LHC



From Frank Siegert


The underlying event at the LHC



A hard pp -collision at the LHC can be interpreted as a hard scattering between partons, accompanied by the underlying event (UE) consisting of:

- Initial and final state radiation
- Beam Remnants
- Multiple Parton Interactions (MPI)
- Hadronization

Many processes are included in the nomenclature "UE" at different scales



Double Parton Scattering (DPS)
Diffractive processes
Semi-hard multiparton interactions

How do we deal with that?



Monte Carlo event generators (PYTHIA, HERWIG, SHERPA..)



Parameters need to be adjusted (tuned) to describe data

- MPI

e.g. $p_T^0 = p_T^{ref} \cdot (E/E_{ref})^\epsilon$
Proton matter distribution profile
Colour reconnection

- Primordial k_T

e.g. Width of the gaussian used for modelling the parton primordial k_T inside the proton

- Parton shower

e.g. Strong coupling value
Regularization cut-off
Upper scale

- Hadronization



e.g. Length of fragmentation strings
Strange baryon suppression

How does one tune all these?

- Choice of parameter ranges and sensitive observables
- Predictions for different parameter choices and interpolation of the MC response
- Data-MC difference and minimisation over parameter space

Before Run II data: trying to predict

Charged particle mult. in the MAX reg. @ 0.9 (left) and 7 (right) TeV



New tunes!



- PYTHIA 8 (CUETP8)
 - HERWIG++ (CUETHpp)
- with various PDFs

Better constrain of the energy extrapolation CR changes with the choice of the PDF

Rising part and plateaux region are well predicted by the new tunes

(EPJC 76 (2016) 155)

After Run II data: the outcome



$\sqrt{s} = 13$ TeV



TOP:
 $dN/d\eta$

ATLAS-CONF-2015-028, PLB 751
(2015) 143

BOTTOM:
 N_{ch} vs p_T^{lead}

ATLAS-PHYS-2015-019,
CMS-FSQ-15-007

None of the tunes
reproduce the data
perfectly!



The energy
dependence of the
MPI fitted to lower
energies is not
optimal

$$p_T^0 = p_T^{ref} \cdot (E/E_{ref})^c$$

Similar strategy used for obtaining CUETP8M1 (BUT..without energy dependence)


GEN-14-001, EPJC 76 (2016) 155

Two types of measurements considered for the fit:

- UE observables (charged particle multiplicity and p_T sum) at 13 TeV in MIN and MAX region as a function of leading track p_T
- Charged particle multiplicity as a function of η in MB collisions
- **CUETP8M2T4: New UE/MB tune at 13 TeV**

TOP-16-021, CMS-GEN-XXX: paper in preparation

Measurement of UE observables in the transverse region by CMS



Transverse regions: $60^\circ < |\Delta\phi| < 120^\circ$:

- **TransMAX:** maximum activity side, often containing a 3rd jet → **MPI/BR + ISR/FSR**
- **TransMIN:** minimum activity side → **MPI/BR**

$$\text{TransAVE} = (\text{TransMAX} + \text{TransMIN})/2$$

$$\text{TransDIF} = \text{TransMAX} - \text{TransMIN} \rightarrow \text{ISR/FSR}$$

Observables:

- average charged-particle multiplicity density (particle density) : $\langle N_{\text{ch}} \rangle / [\Delta\eta \Delta(\Delta\phi)]$
- average transverse-momentum scalar sum density (energy density) : $\langle \Sigma p_T \rangle / [\Delta\eta \Delta(\Delta\phi)]$

MAX and MIN regions included in the tune
CMS-PAS-FSQ-15-007

Considered measurements

Measurement of charged-particle multiplicities by ATLAS and CMS

→ ATLAS measurement: $p_T > 500$ MeV (100 MeV)

$$\frac{1}{N_{\text{ev}}} \cdot \frac{dN_{\text{ch}}}{d\eta}, \quad \frac{1}{N_{\text{ev}}} \cdot \frac{1}{2\pi p_T} \cdot \frac{d^2 N_{\text{ch}}}{d\eta d p_T}, \quad \frac{1}{N_{\text{ev}}} \cdot \frac{dN_{\text{ev}}}{dn_{\text{ch}}} \quad \text{and} \quad \langle p_T \rangle \text{ v.s. } n_{\text{ch}}$$

Not included in the tune

arXiv:1602.01633

→ CMS measurement: any particle $p_T (> 0$ MeV)

Included in the tune

PLB 751 (2015) 143


Measurement of charged-particle multiplicities by CMS

(A) At least 1 charged particle $\left\{ \begin{array}{l} p_T > 0.5 \text{ GeV} \\ |\eta| < 2.4 \end{array} \right.$


- ✦ **Activity**: at least 1 particle with $E > 5 \text{ GeV}$
- ✦ **Veto**: no particle with $E > 5 \text{ GeV}$

- **Inclusive**: (A)


- **Inelastic enhanced**: (A) + **Activity** in **at least one** Forward Region



- **NSD enhanced**: (A) + **Activity** in **both** Forward Regions



- **SD enhanced**: (A) + **Activity** in one Forward Region **and Veto** in the other side



(From Juan's slides)



Not included in the tune
CMS-PAS-FSQ-15-008

The starting point of the Underlying Event tune

Top events are important background for searches (e.g. ttH)

Low jet multiplicity is sensitive to ME and matching to PS

High jet multiplicity is sensitive to PS (i.e. UE tune)



Any considered prediction overestimates the jet multiplicity, when jets come from the parton shower!

Effect seen also at 8 TeV

TOP-12-041, TOP-16-011, TOP-16-021



The starting point of the Underlying Event tune

Need for improvement of the jet multiplicity in top events \rightarrow tune of α_S^{ISR} and h_{damp}
 $\rightarrow h_{damp}$ is an internal parameter inside the POWHEG ME simulation, which regulates the amount of additional hard radiation

Results

$$\alpha_S^{ISR} = 0.1108^{+0.0144}_{-0.0142}$$



$$h_{damp} = 1.581^{+0.658}_{-0.585}$$



TOP-12-041, TOP-16-021

The new Underlying Event Tune

The fit includes five histograms for the UE and MB measurements!



PYTHIA8.219

PDF set: NNPDF30_lo_as_0130

ISR $\alpha_S = 0.1108$ (previous slide)

MultipartonInteractions:ecmPow=0.25208
(from CUETP8M1)

Baseline: Monash tune



Parameter	Tuning Range
MultipartonInteractions:pT0Ref	1.0-3.0
MultipartonInteractions:expPow	0.4-10.0
ColourReconnection:range	0.0-9.0

TOP: chg part. mult. in trans MIN region
BOTTOM: $dN/d\eta$

CMS-GEN-XXX in prep.

Performance of the new tune

Charged particle mult. in the MIN region and $dN/d\eta$ @13 TeV



The new tune has a better description of the plateau region

Rising part of the spectrum seems to prefer a double gaussian matter distribution profile





CMS-FSQ-15-007

PLB 751 (2015) 143

CMS-TOP-16-021

Performance of the new tune

Charged particle mult. in the MIN region and $dN/d\eta$ @13 TeV



Single-diffractive enhanced observables and inelastic cross sections not well described

NEED FOR TUNING DIFFRACTIVE PART OF THE SIMULATION!



CMS-FSQ-15-007

PLB 751 (2015) 143

CMS-TOP-16-021

Immediate to-do list from the CMS side





- Rising part of UE observables solved by choosing (**and tuning**) a double gaussian matter distribution profile
- Try to perform a tune with measured inelastic and diffractive cross sections as inputs

```
SigmaTotal:setOwn = on      ! default = off  
SigmaTotal:sigmaTot = 109.10 ! Total X-Section 13 TeV  
SigmaTotal:sigmaEl = 29.80  ! Elastic X-Section 13 TeV  
SigmaTotal:sigmaXB = 2.538  ! SD X-Section fDD = 0.04  
SigmaTotal:sigmaAX = 2.538  ! SD X-Section fDD = 0.04  
SigmaTotal:sigmaXX = 3.172  ! DD X-Section fDD = 0.04  
SigmaTotal:sigmaAXB = 0.793 ! CD X-Section fCD = 0.01
```

GOALS:

- Good cross section predictions of tunes at 13 TeV
- Cross section obtaining from pile-up reweighting procedure lower than ~ 78 mb (effect also seen by ATLAS?)

CMS is on track for Herwig7 as well!



Available tunes

- CUETHppS1 (CTEQ6L1)
- Default (MMHT14)

CMS Herwig++ tune is not performing well!
Need for tune update

Default tune is good for UE and ND observables

CMS-FSQ-15-007



PLB 751 (2015) 143

CMS-FSQ-15-008

CMS is on track on using Herwig7!

PRD 89 (2014) 092010





EPJC 76 (2016) 451



Two pending issues from Run I:

- H7 LO dijet ME with default tune ($\sigma_{eff} = 15$ mb) describes DPS-sensitive observables at 7 TeV! Very promising comparison!
- Inclusive jet cross section at 7, 13 TeV still not well described by H7 LO..but much better than H++

CMS is on track on Sherpa as well!



Available tunes

- Default Shrimp Tune
- Default Amisic Tune

Good behaviour only of strictly MB observables ($dN/d\eta$ for charged part. with $p_T > 0\text{ GeV}$)


When applying p_T cut on the charged particles, the agreement gets worse..

CMS-FSQ-15-007

PLB 751 (2015) 143

CMS-FSQ-15-008

Sherpa tuning studies (II)



$2 \rightarrow 2$ process from $\hat{p}_T > 3.84 \text{ GeV}$
PDF: CT10
Tune: default Amisic

SOURCE: MCPLOTS4CMS

Bad description of UE observables
at 13 TeV..is it only a tune issue?

**Tuning CMS effort is ongoing
but nothing conclusive on UE
observables!**

CMS-FSQ-15-007

- CMS has a great interest on Monte Carlo models and follows closely tuning issues during LHC Run II
- **A new PYTHIA 8 tune is ready after first RunII data!**
 - It is able to describe UE and MB observables at the same time and uses a lower value of ISR α_S , tuned to jet multiplicities in top events
 - Cross checks with other observables suggest that the new tune behaves very well in general at 13 TeV
 - At 7 TeV, it is also performing well
- **Herwig7 default tune describes well 13 TeV data**
 - Old Herwig++ tunes can not be ported to Herwig7 but need retuning
- **Still some things to understand for Sherpa available tunes**

- CMS has a great interest on Monte Carlo models and follows closely tuning issues during LHC Run II
- **A new PYTHIA 8 tune is ready after first RunII data!**
 - It is able to describe UE and MB observables at the same time and uses a lower value of ISR α_S , tuned to jet multiplicities in top events
 - Cross checks with other observables suggest that the new tune behaves very well in general at 13 TeV
 - At 7 TeV, it is also performing well
- **Herwig7 default tune describes well 13 TeV data**
 - Old Herwig++ tunes can not be ported to Herwig7 but need retuning
- **Still some things to understand for Sherpa available tunes**

THANKS FOR YOUR ATTENTION

BACKUP SLIDES

ATLAS released a CONF NOTE with a new tune for improving the description of soft QCD observables

\sqrt{s}	Measurement type	Rivet name
13 TeV	MB	ATLAS_2016_I1419652 [3]
13 TeV	INEL XS	MC_XS [5]
7 TeV	MB	ATLAS_2010_S8918562 [11]
7 TeV	INEL XS	ATLAS_2011_I89486 [4]
7 TeV	RAPGAP	ATLAS_2012_I1084540 [15]
7 TeV	ETFLOW	ATLAS_2012_I1183818 [14]
900 GeV	MB	ATLAS_2010_S8918562 [11]
2.36 TeV	MB	ATLAS_2010_S8918562 [11]
8 TeV	MB	ATLAS_2016_I1426695 [16]

Parameter	Sampling range	
MultipartonInteractions:pT0Ref	1.00	– 3.60
MultipartonInteractions:ecmPow	0.10	– 0.35
MultipartonInteractions:coreRadius	0.40	– 1.00
MultipartonInteractions:coreFraction	0.50	– 1.00
BeamRemnants:reconnectRange	0.50	– 10.0
Diffraction:PomFluxEpsilon	0.02	– 0.12
Diffraction:PomFluxAlphaPrime	0.10	– 0.40

Table 1: Tuning parameters and sampling range

- Need to improve the description at the new energy
- Focus on total inelastic cross section, $dN/d\eta$ and particle multiplicities at different energy
- Choice of double gaussian matter distribution profile
- First attempt to include diffractive parameters in the procedure

ATL-PHYS-PUB-2016-017