

ISMD 2017 – Tlaxcala City (Mexico)

Heavy flavors in quark gluon plasma: group chasing of the transport coefficients



P.B. Gossiaux
SUBATECH, UMR 6457
Université de Nantes, IMT Atlantique, IN2P3/CNRS


Special thanks to Shanshan CAO and Marlene NAHRGANG

Why heavy flavors in A-A ?

1. Produced early, number conserved through time evolution (even at LHC) \Rightarrow signature (hard probes) of early (hot) phase
2. Weakly affected by late time evolution (heavy, colour transparency)
3. *Strongly affected by the QGP phase ... probe the QGP strength in a way that is arguably under better control from the theory view point*
4. Allows *some* pQCD calculations for the production and annihilation,...
5. Clear decay channel of quarkonia in leptons


See f.i talk by
R. Rapp

Seems to be an ideal alternate probe of dense matter as compared to light sector...



Why open heavy flavors in A-A ?

- Those are for sure sensitive to the early stages
- Much simpler than quarkonia and also sensitive to the medium properties (t_{equil} ($\propto M_Q/T^2$) \Rightarrow clear hierarchy for s, c and b).
- Mandatory to understand Q-Qbar evolution in QGP & quarkonia product.



The Trilogy: \equiv **densitometer**

Thermalisation &
collectivity

\equiv **barometer**

Quenching (leading
hadron)

HQ
Hidden
c & b

\equiv **thermometer**

Quarkonia suppression and
Dimuons product; **heavy Q**
thermal production

HQ are imbedded in expanding matter \Rightarrow they
participate to collective motion and gain elliptic
flow (v_2 :azimutal asymmetry) at finite b


Challenges:

- Description of HQ E-loss / equilibration from fundamental theory
- Joint v_2 - R_{AA} explanation could help to better constrain free parameters...
- Dual game “probe the medium evolution” vs “understands the force”

Paradigm of HF evolution in URHIC


For low and intermediate p_T

No force on HQ before thermalization of QGP




Paradigm of HF evolution in URHIC

For low and intermediate p_T




Somehow arbitrary T_c between
QGP and hadronic phase

Paradigm of HF evolution in URHIC



Basic Consequences of HQ interaction with QGP for the R_{AA}



Flow bump: due to

- *(radial) flow of the medium and coupling at small p_T*
- *recombination with light quarks*


shadowing: due to *initial state nuclear effects*

energy loss: due to

- elastic and inelastic scatterings
- *opacity of the medium*
- coherence important at larger p_T

Italic: extrinsic to the HF coupling with QGP AKA « energy loss model »

Basic Consequences of HQ interaction with QGP for the v_2



Small p_T : height of v_2 at low p_T

sensitive to:

- Bulk anisotropy, mostly at the late times
- The drag force acting locally on HF

high p_T non-0 v_2 is due to anisotropic Eloss (same ingredients as for the RAA + geometrical anisotropy of initial distribution of matter)

intermediate p_T : onset and offset of many competing effects.


2 Important remarks:

- Any energy loss model, even the roughest one, will generate these typical structures in the R_{AA} and the v_2 . Getting a correct **quantitative** agreement is much more involved.
- Quantitative predictions also depends on numerous extra ingredients

Setting the scene: E-Loss and thermalization

(init) $P_T \approx m_Q$

- Bulk part of Q production
- E gain becomes probable
- HQ scatter and can thermalize with the medium
- very \neq from light quarks
- *Dominated by collisional processes and diffusion*
- Non perturbative effect (small momentum transfert, coalescence with light quark)
- 1 dominant transport parameter




(init) $P_T \gg m_Q$

- Rare processes
- Mostly Energy loss
- HQ go on nearly straight lines and probe the opacity of matter. Little thermalization
- \sim light quarks (s.e.p.)
- *Coherent radiative + collisional processes*
- Good test of pQCD... Theory at work (a priori)
- Several transport coefficients implied (dE/dx , B_T , ...)

Various approaches to modeling

Bottom-up schemes (microscopic -> mesoscopic):

- Assume (effective) degrees of freedom and (effective) interactions
- Take insights and constraints from the fundamental QCD theory, but often includes some free parameter
- Rely on more or less sophisticated realizations of the transport theory



Various approaches to modeling: Langevin Dynamics

- FP equation for the HQ distribution

$$\frac{\partial}{\partial t} f_Q(\mathbf{p}, t) = \nabla_i [\textcolor{blue}{A^i} f_Q(\mathbf{p}, t) + \nabla_j (\textcolor{red}{B_{ij}} f_Q(\mathbf{p}, t))]$$

with (in local isotropic medium):

$$A^i(\mathbf{p}, T) = \underbrace{\eta(p, T) p^i}_{\text{Friction (drag) coefficient; relaxation rate}}$$

and

Longitudinal and transverse projectors

$$\hat{B} = \textcolor{red}{B_1} \hat{P}_L + \textcolor{red}{B_0} \hat{P}_T$$

Longitudinal and transverse diffusion coefficients

- ⇒ Langevin Dynamics for each individual HQ: $\frac{d\mathbf{p}}{dt} = -\eta(p)\mathbf{p} + \hat{\xi}$

with $\langle \xi_i(t) \xi_j(t') \rangle = 2B_{ij}\delta(t - t')$

- In most implementations: Einstein relation imposed in order to obtain asymptotic Boltzmann – Jüttner distributions :


$$\textcolor{red}{B_1} (= \textcolor{red}{B_0}) = \textcolor{blue}{\eta} E_Q T \quad \text{(in post point Ito; no systematic prescription)}$$

pQCD inspired models (f.i. Nantes)


Colisional component

- One-gluon exchange model: reduced IR regulator λm_D^2 in the hard propagator, fixed on HTL Energy loss
- Running coupling $\alpha_{\text{eff}}(t)$ and self consistent Debye mass

$$m_{D\text{self}}^2(T) = (1+n_f/6) 4\pi \alpha_{\text{eff}}(m_{D\text{self}}^2) T^2$$



Radiative component



- Extension of Gunion-Bertsch approximation beyond mid-rapidity and to finite mass m_Q) distribution of induced gluon radiation per collision ($\Delta E_{\text{rad}} \propto E L$):

$$P_g(x, \mathbf{k}_\perp, \mathbf{q}_\perp, m_Q) = \frac{3\alpha_s}{\pi^2} \frac{1-x}{x} \left(\frac{\mathbf{k}_\perp}{\mathbf{k}_\perp^2 + xm_Q^2} - \frac{\mathbf{k}_\perp - \mathbf{q}_\perp}{(\mathbf{k}_\perp - \mathbf{q}_\perp)^2 + xm_Q^2} \right)^2$$

- LPM effect for moderate gluon energy

Implemented in MC@HQ + EPOS2(3) through Boltzmann

But also BAMPS, LBL-CCNU, Duke,...

Potential models (TAMU)

- Thermodynamic T-matrix approach, $T = V + VGT$, given by a two-body driving kernel V , estimated from the IQCD internal/free energy for a static $Q\bar{Q}$ pair; increase of coupling with QGP at small momentum

D. Cabrera, R. Rapp PRD 76 (2007); H. van Hees, M. Mannarelli, V. Greco, R. Rapp PRL 100 (2008)


- Comprehensive sQGP approach for the EoS, light quark & gluon spectral functions, quarkonium correlators and HQ diffusion.

F. Riek, R. Rapp PRC 82 (2010); S. Liu, R. Rapp arxiv:1612.09138


- Resonance correlations in the T-matrix naturally lead to recombination (resonance recombination model) near T_c from the same underlying interactions!

M. He, R. Fries, R. Rapp PRC 82 (2010), PRC 86 (2012)

- Implemented through Langevin dynamics in hydro evolution or in URQMD



No good q-particle at low p



Large coupling at small p_Q 13

Quasi particle models (f.i DQPM)

- Nonperturbative effects near T_c are captured by $\alpha_s(T)$, leading to thermal masses/widths, determined from fits to IQCD EoS.

A. Peshier et al. PLB 337 (1994), PRD 70 (2004); M. Bluhm et al. EPJC 49 (2007); W. Cassing et al. NPA 795 (2007)




- Relaxation rates larger than in pQCD for all T relevant for QGP, slightly smaller than the ones from TAMU

H. Berrehrah et al, PHYSICAL REVIEW C 90, 064906 (2014)


- Implemented for HF dynamics in e.g. PHSD (full off-shell, off-equilibrium transport).

T. Song et al. PRC 92 (2015), PRC 93 (2016)


But also CATANIA



Various models... with various coupling strength of heavy quarks to the QGP !



Models vs DATA at LHC: Sapore Gravis Report



Despite various prescriptions for Energy loss, a lot a model can cope with the data₁₆

Models vs DATA at LHC: Sapore Gravis Report

Sapore Gravis I3HP Network (Led by G. Martinez) => comprehensive report

Eur. Phys. J. C (2016) 76:107
DOI 10.1140/epjc/s10052-015-3819-5

THE EUROPEAN
PHYSICAL JOURNAL C



CrossMark

Review

Heavy-flavour and quarkonium production in the LHC era: from proton–proton to heavy-ion collisions

A. Andronic¹, F. Arleo^{2,3}, R. Arnaldi⁴, A. Beraudo⁴, E. Bruna⁴, D. Caffarri⁵, Z. Conesa del Valle⁶, J. G. Contreras⁷, T. Dahms⁸, A. Dainese⁹, M. Djordjevic¹⁰, E. G. Ferreiro¹¹, H. Fujii¹², P.-B. Gossiaux¹³, R. Granier de Cassagnac², C. Hadjidakis⁶, M. He¹⁴, H. van Hees¹⁵, W. A. Horowitz¹⁶, R. Kolevatov^{13,17}, B. Z. Kopeliovich¹⁸, J.-P. Lansberg⁶, M. P. Lombardo¹⁹, C. Lourenço⁵, G. Martinez-Garcia¹³, L. Massacrier^{6,13,20,a}, C. Mironov², A. Mischke^{21,22}, M. Nahrgang²³, M. Nguyen², J. Nystrand²⁴, S. Peigné¹³, S. Porteboeuf-Houssais²⁵, I. K. Potashnikova¹⁸, A. Rakotozafindrabe²⁶, R. Rapp²⁷, P. Robbe²⁰, M. Rosati²⁸, P. Rosnet²⁵, H. Satz²⁹, R. Schicker³⁰, I. Schienbein³¹, I. Schmidt¹⁸, E. Scomparin⁴, R. Sharma³², J. Stachel³⁰, D. Stocco¹³, M. Strickland³³, R. Tieulent³⁴, B. A. Trzeciak⁷, J. Uphoff³⁵, I. Vitev³⁶, R. Vogt^{37,38}, K. Watanabe^{39,40}, H. Woeber⁵, P. Zhuang⁴¹


- Compilation of all existing (published) experimental results vs a large number of models
- Explicitation of these models on the same footing (as much as possible)
- (partial) compilation of « Fokker Plank coefficients » (drag & diffusionS)

Models vs DATA at LHC: Sapore Gravis Report



The models considered
In SG. New ones in the
meanwhile (pHSD, CCNU,
Catania)

Model	Heavy-quark production	Medium modelling	Quark–medium interactions	Heavy-quark hadronisation	Tuning of medium coupling (or density) parameter(s)
Djordjevic et al. [511–515]	FONLL no PDF shadowing	Glauber model nuclear overlap no fl. dyn. evolution	Rad. + coll. energy loss finite magnetic mass	Fragmentation	Medium temperature fixed separately at RHIC and LHC
WHDG [459,519]	FONLL no PDF shadowing	Glauber model nuclear overlap no fl. dyn. evolution	Rad. + coll. energy loss	Fragmentation	RHIC (then scaled with $dN_{ch}/d\eta$)
Vitev et al. [422,460]	Non-zero-mass VFNS no PDF shadowing	Glauber model nuclear overlap ideal fl. dyn. 1 + 1d Bjorken expansion	Radiative energy loss in-medium meson dissociation	Fragmentation	RHIC (then scaled with $dN_{ch}/d\eta$)
AdS/CFT (HG) [624, 625]	FONLL no PDF shadowing	Glauber model nuclear overlap no fl. dyn. evolution	AdS/CFT drag	Fragmentation	RHIC (then scaled with $dN_{ch}/d\eta$)
POWLANG [507–509,585,586]	POWHEG (NLO) EPS09 (NLO) PDF shadowing	2 + 1d expansion with viscous fl. dyn. evolution	Transport with Langevin eq. collisional energy loss	Fragmentation recombination	Assume pQCD (or I-QCD U potential)
MC@_sHQ+EPOS2 [528–530]	FONLL EPS09 (LO) PDF shadowing	3 + 1d expansion (EPOS model)	Transport with Boltzmann eq. rad. + coll. energy loss	Fragmentation recombination	QGP transport coefficient fixed at LHC, slightly adapted for RHIC
BAMPS [537–540]	MC@NLO no PDF shadowing	3 + 1d expansion parton cascade	Transport with Boltzmann eq. rad. + coll. energy loss	Fragmentation	RHIC (then scaled with $dN_{ch}/d\eta$)
TAMU [491,565, 606]	FONLL EPS09 (NLO) PDF shadowing	2+1d expansion ideal fl. dyn.	Transport with Langevin eq. collisional energy loss diffusion in hadronic phase	Fragmentation recombination	Assume I-QCD U potential
UrQMD [608–610]	PYTHIA no PDF shadowing	3+1d expansion ideal fl. dyn.	Transport with Langevin eq. collisional energy loss	fragmentation recombination	Assume I-QCD U potential
Duke [587,628]	PYTHIA EPS09 (LO) PDF shadowing	2+1d expansion viscous fl. dyn.	Transport with Langevin eq. rad. + coll. energy loss	Fragmentation recombination	QGP transport coefficient fixed at RHIC and LHC (same value)

Extended p_T measurement by CMS...



Models vs DATA at RHIC



Which information / properties to extract from all of this ?


Some people are confident they can...

Inspiration from the JET Collaboration

PHYSICAL REVIEW C **90**, 014909 (2014)

Extracting the jet transport coefficient from jet quenching in high-energy heavy-ion collisions

Karen M. Burke,¹ Alessandro Buzzatti,² Ningbo Chang,³ Charles Gale,⁴ Miklos Gyulassy,⁵ Ulrich Heinz,⁶ Sangyong Jeon,⁴ Abhijit Majumder,¹ Berndt Müller,⁷ Guang-You Qin,^{1,3} Björn Schenke,⁷ Chun Shen,⁶ Xin-Nian Wang,^{2,3,*} Jiechen Xu,⁵ Clint Young,⁸ and Hanzhong Zhang³
(JET Collaboration)



$$\hat{q} = \left\langle \frac{\Delta p_T^2(L)}{L} \right\rangle$$


$\sim 4 \times B_0$ FP Coefficient in
the case of HQ

See talk by A. Majumder

Which relevant quantity to measure with HQ ?

Reminder: The JET collaboration calibrated their transport coefficient on the quenching : $\Delta E = F(E, L, \hat{q}, \dots)$

For Heavy Flavor at small and intermediate p_T , the link between ΔE and \hat{q} is more intricate => « extract » $\langle dP \rangle / dt$?



Several issues:

- Non-trivial momentum dependence
- No prediction from lQCD


Which relevant quantity to measure with HQ ?

At $p_T \approx 0$: All FP coefficients are equivalent and measure the (long wave) coupling strength to the medium :

$$B_1 = B_0 = B = \eta m_Q T$$



Slope of $\langle dP \rangle / dp$ at $p \rightarrow 0$



Most often recatch into the spatial diffusion coefficient D_s $\left(= \frac{1}{6} \lim_{t \rightarrow \infty} \frac{\langle (\mathbf{x}(t) - \mathbf{x}(0))^2 \rangle}{t} \right)$

$$D_s = \frac{B}{m_Q^2 \eta^2} \Big|_{p \approx 0} = \frac{T}{m_Q \eta} = \frac{T^2}{B}$$

Good News: Lattice QCD calculation for D_s

Small $D_s \Leftrightarrow$ large
relaxation \Leftrightarrow large
fluctuations

IQCD Calculation of D_s

- Lattice QCD at finite T is performed in Euclidean space notoriously difficult to calculate dynamical quantities.
- Up to 2014, D_s was evaluated directly through the (**narrow**) diffusion peak of the spectral function evaluated from current – current correlator (**hard**)
- From 2014: Use of the field – field correlator in order to obtain a better shaped spectral function:

$$G_E(\tau) \equiv -\frac{1}{3} \sum_{i=1}^3 \frac{\left\langle \text{Re Tr} \left[U(\beta; \tau) gE_i(\tau, \mathbf{0}) U(\tau; 0) gE_i(0, \mathbf{0}) \right] \right\rangle}{\left\langle \text{Re Tr} [U(\beta; 0)] \right\rangle}$$


- Then obtain the variance κ of stochastic forces (a transport coefficient; $\kappa = 2 \times B$) from the slope of spectral function ρ_E at $\omega = 0$:

$$\kappa \equiv \lim_{\omega \rightarrow 0} \frac{2T \rho_E(\omega)}{\omega} \quad \text{with } \rho_E \text{ extracted from } G_E(\tau) = \int_0^\infty \frac{d\omega}{\pi} \rho_E(\omega) \frac{\cosh[\omega(\frac{\beta}{2} - \tau)]}{\sinh[\frac{\omega\beta}{2}]}$$

- Main result : $\kappa/T^3 = 1.8 \dots 3.4$ then convert to D_s

lQCD Calculation of D_s


- This leads to $2\pi T D_s = 3.7 \dots 6.9$



- Still drastic approximations/limitations: quenched QCD, heavy quark vs. charm quark, (no) continuum extrapolation,...
- Relaxation time $\eta^{-1} = \frac{m_Q D_s}{T} = \frac{0.59 \dots 1.1 m_Q}{T^2} \approx 2.5 \dots 5 \text{ fm/c}$

Model + Exp. Constrains on D_s

- F.i., recent STAR study:



STAR Collab. PRL 118 (2017)

compare with	$2\pi TD_s$	χ^2/NDF	p -value
SUBATECH [11]	2–4	15.2 / 8	0.06
TAMU c quark diff. [13]	5–12	10.0 / 8	0.26
TAMU no c quark diff. [13]	-	29.5 / 8	2×10^{-4}
Duke [14]	7	35.7 / 8	2×10^{-5}
LBT [15]	3–6	11.1 / 8	0.19
PHSD [10]	5–12	8.7 / 7	0.28
3D viscous hydro [42]	-	3.6 / 6	0.73

TABLE I. $D^0 v_2$ in 0–80% centrality Au+Au collisions compared with model calculations, quantified by χ^2/NDF and the p -value (the probability of observing a χ^2 that exceeds the current measured χ^2 by chance). $2\pi TD_s$ values quoted are in the range of T_c to $2T_c$. χ^2/NDF is calculated in the p_T range wherever the model calculation is available.

- Not really conclusive; χ^2/NDF is not a smooth function of D_s : large residuals
- Beware: Models are essentially validated at finite $p_{(T)}$; extrapolation at 0 momentum might contain further uncertainties
- Urgent need for collective actions to better control the « residuals »

Recent Collective actions beyond Sapore Gravis

- **Heavy Quark – Working Group** (convener: X-N Wang); in the spirit of the Jet Collaboration, the goal is, in a first stage, to :
 - Collect and compare the transport coefficients from various models,
 - Measure and understand their consequences by first studying a simpler brick problem
 - Estimate some systematics + uncertainties
- **EMMI Rapid Reaction Task Force** (organizers: A. Andronic, R. Averbeck, PB Gossiaux, S. Maschiocchi, R. Rapp):
 - Global strategy to extract the diffusion coefficient from the intercomparison between models and data
 - Collect and analyse all ingredients from various models
 - Identify constraints from lQCD
 - Initiate discussions to assess the limitations of some existing models.
- **Lorentz Workshop** on “Tomography of the quark-gluon plasma with heavy quarks”; oct 2016 (organizers: J. Aichelin, R. Granier de Cassagnac, M.P. Lombardo, A. Mischke, N. Xu): The goal of this workshop is to identify common understandings and develop strategies for the upcoming 5 years to achieve a profound knowledge of the dynamical properties of the quark-gluon plasma

Heavy-Quark Working Group

- LBL-CCNU (**XN Wang, S. Cao**)
- Duke (S. Bass , S. Cao, M. Nahrgang, Y Xu)
- Catania (V. Greco, S. Das, S. Plumari, F. Scardina)
- TAMU (R. Rapp + M. He)
- Frankfurt pHSD (E. Bratkovskaya, T. Song, H. Berrehrah)
- Nantes (J. Aichelin, PB Gossiaux, M. Nahrgang)



PHYSICAL REVIEW C ~~90, 014909 (2014)~~
(2017)

Extracting the ~~jet~~ transport coefficient from ~~jet quenching~~ in high-energy heavy-ion collisions

charm Drag coefficient D-meson observables
Fluctuation tensor

Heavy-Quark Working Group



After 3 meetings, footprints of the physics start to emerge... but no firm conclusion yet



For step 1: Compare HQ spectra from different models in static medium with common initial condition


Models	note	basic	tune 1
LBL-CCNU	fix α_s	$\alpha_s = 0.3$	$\alpha_s = 0.26$
Duke	fix α_s	$\alpha_s = 0.3$	$\alpha_s = 0.23$
Catania QPM	$\alpha_s(T)$	$K = 1$	$K = 2$
Catania pQCD	$\alpha_s(T)$	$K = 1$	$K = 3.4$
TAMU	U -potential	no tuning	no tuning
Frankfurt PHSD	$\alpha_s(T)$	no tuning	no tuning
Nantes col. + rad.	$\alpha_s(q^2)$	$K = 1$	$K = 0.8$
Nantes col. only	$\alpha_s(q^2)$	$K = 1$	$K = 1.5$

★: Radiative included

Basic: original model


Tune 1: favorite tuning of each group in order to describe D meson data with their own ingredients (background, hadronization,...) ; K = rate multiplier

qhat – momentum broadening per unit time (due to elastic scattering) ONLY




Tune 1 would be the « extraction »... Large « error bands » except at small p_T !!!
Nature is maybe not so kind...

A (drag force) – p_L change per unit time (total)





Tune 1 would be the « extraction »... Large « error bands » as well

Time evolution of average quantities (thermalization process)



Looking at the evolution in a brick, still large fluctuations observed

Time evolution of average quantities (thermalization process)







Models without radiation =>
larger transverse broadening at large momentum

Looking at the evolution in a brick, still large fluctuations observed



E and k_T distribution at 4 fm

k_T



Looking at the outcome of the evolution in a brick, still large fluctuations observed, especially in long direction -> seek for more exclusive variables !!!

R_{AA} of c -quark in static medium



Initial spectrum: simple power-law parametrization:
G. Moore and D. Teaney, Phys. Rev. C71 (2005) 064904. Eq. 5.12


$$\frac{dN}{dy d\eta d^2 p_T} \propto \delta(y - \eta) \frac{1}{(p_T^2 + \Lambda^2)^\alpha}$$

$$\alpha = 3.5, \Lambda = 1.849 \text{ GeV}$$

« fluctuations » of the $R_{AA}(c)$ in a brick... much larger than the $R_{AA}(D)$ in URHIC 35



R_{AA} of c -quark in static medium



... but rather good hierarchical ordering as comp. to A (not so clear with \hat{q}/T^3)

Heavy-Quark Working Group: preliminary conclusions

Previous studies suggest that the other « ingredients » play a crucial role in each prediction and prevent accurate extraction of Fokker Planck coefficient from the URHIC data...

.... assuming it makes sense otherwise !

=> Difficult to extract physics out of the variety of {tune 1}; **conclusion reinforced by this study**

=> **Method and further work: Use the brick configuration and some « Gedanken Experiment » to provide information on the smallest achievable ERROR-BAND WIDTH (and not on the absolute values of the FP coefficients)**

Stay tuned !

EMMI RRTF

Thanks to the generosity of EMMI !!!

1st meeting: 18-22 July 2016; GSI (Germany)

2nd meeting: 12-14 Dec 2016; GSI (Germany)

Researchers involved:

- Organizers
- HQ-WG members
- Other key players in HF – QGP tomography (A. BERAUDO, M. DJORDJEVIC, C. GREINER , G. INGHIRAMI, H. VAN HEES, I. VITEV, CUEJET...)
- IQCD experts (O. KACZMAREK, P. PETRECZKY)
- QCD and EFT experts (G. MOORE , J. PAWLOWSKI)
- Selected experimentalists (J. BIELCIK, P. BRAUN MUNZINGER, E. BRUNA, Z CONESSA, A. DAINESSE, YJ LEE, F. PRINO, J. STACHEL)

EMMI RRTF

Goal to attack the problem with a broad view right from the beginning...



Topics:

- Initial spectra and shadowing
- Bulk evolution and consequence on HF observables
- Transport implementation
- Hadronization
- Microscopic models for HF energy loss and constrains from QCD at low and high momentum
- Future observables

Selected topics presented here

EMMI RRTF : Initial HF spectra and shadowing

Collection of models vs data by Yen-Jie Lee:





Some « outliers »... Is it acceptable or does it has measurable consequence either on the « extraction » of the transport coefficient (for « tunable » models) or on the agreement with experiment.

« We all do FONLL / GM - VFNS » ... yes, but with slightly different parameters !

EMMI RRTF : Initial HF spectra and shadowing

Right now: data better than uncertainty band in theory:

3 best fits extracted from members of the ALICE collaboration, with a fair wish to explore various hardness (BASELINE)




Consequence of the baseline on the RAA of D mesons in the Nantes Model
 (thermalisation at small p_T => the initial profile has a large impact in the ratio)

Consequence for the collective chase: ideally, all models should adopt the baseline; minimal action: check that the χ^2 with their own production model is at least as small as the one found in the baseline (rejection if $\chi^2/\text{NDF} > 2$).

EMMI RRTF : Initial HF spectra and shadowing

- RAA for c quarks with MNR+EPS09NLO
- Proposal: multiply the input c-quark p_T spectrum from FONLL by this RAA and use the band to define a band on final R_{AA} and v_2



EMMI RRTF : Initial HF spectra and shadowing

Consequence for the collective chase:

- ideally, all models should adopt the same prescription for shadowing, and the uncertainties on the shadowing should then be recast in a global systematic error on the D_s , *common to all models*
- If some models have an intrinsic theory to evaluate shadowing and are unwilling to modify this for the sake of consistency (f.i. EPOS3), the minimal « quality control » should be to implement the common prescription for shadowing and display the consequences on the observables and on their extraction of the transport parameters in order to document the origin of possible differences.
- Perspective (apart from hoping on better control on shadowing):
 - Go for B or to v_2 , less sensitive to shadowing
 - Uncertainties on the shadowing may partly factor out if the ratio of R_{AA} at different centralities is considered



Shadowing seems to act in a nearly multiplicative way

EMMI RRTF : Consequences from the **bulk choice**


Question: What is the role of the different medium evolution models, and how do different predictions for the bulk cooling and expansion temperature in the current models manifest themselves in HF observables ?

Method: adopt a common $\alpha_s=0.4$ -pQCD $\times 5$ cross section for thermal light partons acting on c-quarks (or associated FP coefficients for models based on FP) in all frameworks.

This allows to probe the effect of the bulk with a mechanism that has a D_s roughly similar to the one extracted from IQCD
(caution: of course the weighting in the $T(t)$ might differ a bit)



Consequences from the bulk choice on c-quarks




EMMI RRTF : Consequences from the **bulk choice**

Under work: comparison of genuine bulk quantities (spectra and v2 of hadrons at freeze out) in order to establish some systematics

Conclusions:


- (Even state of the art) Bulk evolution can have significant consequences on the HF observables (v_2 : $\approx \pm 33\%$ for most of them but can be more in some case)
- Off equilibrium effects present in Boltzman and Kadanoff-Baym modeling of the bulk must be understood in order to provide a reliable estimation of the transport coefficients from the data.
- For all frameworks relying on fluid dynamics to describe the bulk evolution, adopting a limited number of « common hydro » would permit to shrink the residuals.
- In //: need for a more precise study of the quantitative bulk characteristics that impact the HF observables



Global Extraction


Theory « Error bar » ?

Where we stand...




Understanding the role of various ingredients

Method : One group collects one given ingredient of all others and proceed to systematic comparison



Global Extraction

Suggested method once the role of each component is understood: adopt common bona fide ingredients (for some « standard component »)... Collective => need some consensus




Preliminary conclusions

- After 10 years of individual work, the open heavy flavour community has moved towards a bit of collectivity.
- Benefits are expected in the upcoming times
- This also implies that some models will be ruled out, while others – whose applicability ranges remain to be settled – will allow a more robust description of QGP probing with heavy flavors...
- ... which is exactly what science should be.

Mid-term Perspectives

- More sophisticated observables (higher flow components, correlations,...):




Up to now, no smoking gun => should just be part of the global fit

- « No model approach » : Bayesian analysis pursued by the Duke group


$$D_s(T, p) = \frac{1}{1+(\gamma^2 p)^2} (D_s 2\pi T)^{\text{lin}}(T; \alpha, \beta) + \frac{(\gamma^2 p)^2}{1+(\gamma^2 p)^2} (D_s 2\pi T)^{\text{PQCD}}(T, p)$$

Data




Mid-term Perspectives


- Each individual model designed to cover the full p_T range should satisfy « basic constrains » at the phase boundaries (low and high p_T ; low and high T)



Be in the bulk part of world IQCD data on D_s at small p




Be compatible with the state of the art multi-gluonic radiative Energy loss at large p_T



Reproduce the gradual transition from heavy quarks to heavy mesons around T_c

Global Extraction

Method : « use constrains »... collaborative; too generic, exhausts human ressources



Long-term Perspectives


- Input from IQCD at finite heavy quark momentum
- More insight from IQCD on the QGP effective DF and their dynamics
- Schwinger – Dyson at finite temperature

Back UP

Some thoughts on methodology

2005-2015 was the era of model development for OHF in URHIC (nearly every idea was publishable)... Now it is time to think...

But maybe physics is not so bad with us ... what if some “mesoscopic” coefficients – like f.i. momentum loss per unit time -- have a major influence (nearly) irrespective of the (microscopic) model ?



EMMI RRTF: Some focus

Transport coefficients

4 Transport Coefficients

- 4.1 General Features
- 4.2 Comparison of Existing Models
- 4.3 Information and Constraints from Lattice QCD
 - 4.3.1 Computations of Diffusion Coefficient
 - 4.3.2 Susceptibilities and Role of Heavy-Quark Mass
- 4.4 Repository of Fokker-Planck Coefficients
- 4.5 High- p_T Energy Loss and \hat{q}
- 4.6 “Applicability Chart” in (p, T)

Up to there, we were all good friends..

Consensus on the fact that 2 clear cut regimes:

- Small p_T , where lQCD should provide strong constrains in the future
- Large p_T , where all models should converge -> *unique* pQCD with *unique* transp. Coeff.

No Consensus on :

- Room for intermediate p_T
- Whether physics boils down to Fokker Planck & Transport coefficients
- Should we abandon HTL inspired models around T_c
- What to do with models not fulfilling these constrains ?
- How to go beyond the present models (not the goal of EMMI RRTF)

Open issues

EMMI RRTF: Some focus

Transport coefficients

4 Transport Coefficients

4.1 General Features
4.2 Comparison of Existing Models
4.3 Information and Constraints from Lattice QCD
4.3.1 Computations of Diffusion Coefficient
4.3.2 Susceptibilities and Role of Heavy-Quark Mass
4.4 Repository of Fokker-Planck Coefficients
4.5 High- p_T Energy Loss and \hat{q}
4.6 “Applicability Chart” in (p, T)

Up to there, we were all good friends..



What is a c « quark » around T_c ?



- A quasi particle, as treated by many practitioners ? Seems to be ruled out by IQCD
- A spectral function
- A mix state between c-state and D-state ?

How to deal consistently with the transport of such object ? Outside the reach of EMMI

Boltzmann vs Langevin Dynamics

Langevin from Boltzman view point:

- For « exclusive process », momentum distributions differ significantly, even after imposing Einstein relation (ER):




- These differences should me seen in observables like g-HQ correlations

Boltzmann vs Langevin Dynamics

Langevin from Boltzman view point:

- For coarse grained observables like the R_{AA} and the v_2 , the agreement between the 2 transport schemes essentially depends on the isotropization strength of the cross section (i.e., the Debye mass of the gluon propagator)




S Das et al, Phys. Rev. C 90,
044901 (2014)

- For $m_D = g T \approx 2 T$ found f.i. in the Quasi Particle Model, extra coupling is found for the R_{AA} using LV, which can be suppressed by reducing the FP coefficients by $\approx 30\%$

Boltzmann vs Langevin Dynamics

Boltzman from Langevin view point:

- There are a lot of situations where Langevin dynamics applies, but not Boltzmann, thanks to the large mass of the particle.
- It is even a result proven for dynamical systems (conditions on the velocity applies as well)
- In a dense strongly coupled system, this is likely to be the case !



Constraints at high p_T

- Any reliable Energy loss model should be compatible with the D_s IQCD world data at small p_T and with state of the art energy loss calculations at large p_T





Figure 21: Relative energy loss, $\Delta E/E$, of c -quarks (upper panels) and b -quarks (lower panels) for the NLO calculation presented in Sec. 5.4, including multi-gluon emission, as a function of the path-length L for various initial energies, E , and fixed QGP temperatures of $T=0.2$ GeV (left panels) and $T=0.3$ GeV (right panels).

The weak to strong axis for HQ

“Naive” pQCD
(WHDG, ASW,...)
 $\hat{q} \approx 1 \text{ GeV}^2/\text{fm}$

“Optimized”
pQCD (ok with
sions)

ASW (pure rad. energy loss;
extended BDMPS)



Armesto et al Dainese, Phys. Rev D (hep-ph/0501225) & Phys.Lett. B637 (2006) 362-366 hep-ph/0511257



Conclude to rough agreement,
subjected to b/c ratio in p-p

So-called “Failure of pQCD approach” aka
“the non photonic single electron
puzzle”

coll Eloss (BT and TG) + radiative Eloss

WHDG

Beauty is
the
problem...
but beauty
is found to
contribute



M Aggarwal et al, STAR, PRL 105 202301



The weak to strong axis for HQ

“Naive” pQCD
(WHDG, ASW,...)
 $\hat{q} \approx 1 \text{ GeV}^2/\text{fm}$

“Optimized”
pQCD

So-called “Failure of pQCD approach” aka
“the non photonic single electron
puzzle”

Collisional model with running α_s and optimized
gluon propagator (Peshier, Gossiaux and Aichelin,
BAMPS)



The weak to strong axis for HQ


“Naive” pQCD
(WHDG, ASW,...)
 $\hat{q} \approx 1 \text{ GeV}^2/\text{fm}$

So-called “Failure of pQCD approach” aka
“the non photonic single electron
puzzle”

“Optimized”
pQCD

Running α_s (Peshier, Gossiaux & Aichelin, Uphoff &
Greiner)

Distortion of heavy meson fragmentation functions due to the
existence of bound mesons in QGP, R. Sharma, I. Vitev & B-W
Zhang 0904.0032v1 [hep-ph]



The weak to strong axis for HQ


“Naive” pQCD
(WHDG, ASW,...)
 $\hat{q} \approx 1 \text{ GeV}^2/\text{fm}$

“Optimized”
pQCD

Distortion
fragments
existence
QGP, R. S
Zhang 09(

So-called “Failure of pQCD approach” aka
“the non photonic single electron
puzzle”

Running α_s (Peshier, Gossiaux & Aichelin, Uphoff &
Greiner)



Bound states diffusion or non-
perturbative, lattice potential
scattering models (see R. Rapp
and H Van Hees 0903.1096 [hep-
ph] for a review)

FIG. 40: (Color online) Comparison of Langevin-based models from [74–76] to the heavy flavor electron R_{AuAu} for 0–10% centrality and v_2 for minimum-bias collisions.

The weak to strong axis for HQ

“Naive” pQCD
(WHDG, ASW,...)
 $\hat{q} \approx 1 \text{ GeV}^2/\text{fm}$


“Optimized”
pQCD

Distortion of heavy meson
fragmentation functions due to the
existence of bound mesons in
QGP, R. Sharma, I. Vitev & B-W
Zhang 0904.0032v1 [hep-ph]

So-called “Failure of pQCD approach” aka
“the non photonic single electron
puzzle”

Running α_s (Peshier, Gossiaux & Aichelin, Uphoff &
Greiner)

Bound states diffusion or non-
perturbative, lattice potential
scattering models (see R. Rapp
03.1096 [hep-



ADS/CFT
(akamatsu et
al)

The weak to strong axis for HQ

