

CHROMOMAGNETIC AND CHROMOELECTRIC DIPOLE MOMENTS OF THE TOP QUARK IN THE 4GTHDM

ALAN IGNACIO HERNÁNDEZ JUÁREZ, AGUSTÍN MOYOTL
ACUAHUITL AND GILBERTO TAVARES VELASCO

FCFM, BUAP



XXXII REUNIÓN ANUAL DIVISIÓN DE PARTÍCULAS Y CAMPOS,
SMF
MAY 29, 2018

INTRODUCTION


- Since its discovery in 1995 the top quark has played a special role in the study of the phenomenology of the standard model (SM).
- The LHC is a top quark factory, then the study of the new contributions to the chromomagnetic dipole moment (CMDM) and chromoelectric dipole moment (CEDM) of the top quark is a topic worth studying as they could be at the reach of experimental measurement in the near future.
- A nonzero CEDM is a clear evidence of CP violation.
- CP violation is necessary to explain the baryon asymmetry of the universe (Sakharov's criteria).

INTRODUCTION

- In the SM, the complex phase of the CKM matrix gives rise to CP violation, but it is still not enough to explain the baryon asymmetry.
- New sources of CP violation beyond the SM are required.
- The CMDM and CEDM can be induced via the following Lagrangian

$$\mathcal{L} = -\frac{g_s T^a}{2} \bar{t} \frac{a_t}{2m_t} \sigma^{\mu\nu} t G_{\mu\nu}^a - \frac{T^a}{2} \bar{t} i \sigma^{\mu\nu} \gamma^5 d_t t G_{\mu\nu}^a,$$

- Where a_t is the CMDM, while d_t is the CEDM.



4GTHDM

- The 4GTHDM is a variation of a THDM type-II which introduces a fourth generation of fermions. ([PhysRevD.84.053009](#))
- A model with a fourth generation of SM-Like quarks was studied in the past (SM4).
- Unfortunately, the SM4 is not consistent with the Higgs production at the LHC.
- In the 4GTHDM the theoretical prediction for Higgs boson production at the LHC remains unchanged. ([PhysRevD.86.115008](#))

4GTHDM

- The Yukawa Lagrangian of the quark sector can be written as follows:

$$\mathcal{L}_Y = -\bar{Q}_L \left(\Phi_1 \mathbf{F} \cdot \left(\mathbf{I} - \mathbf{I}_d^{\alpha_d \beta_d} \right) + \Phi_2 \mathbf{F} \cdot \mathbf{I}_d^{\alpha_d \beta_d} \right) d_R \\ - \bar{Q}_L \left(\tilde{\Phi}_1 \mathbf{G} \cdot \left(\mathbf{I} - \mathbf{I}_u^{\alpha_u \beta_u} \right) + \Phi_2 \mathbf{G} \cdot \mathbf{I}_u^{\alpha_u \beta_u} \right) u_R + \text{H.c.},$$

- We focus in the 4GTHDM-I, where $(\alpha_b, \beta_{b'}, \alpha_t, \beta_{t'}) = (0, 1, 0, 1)$: Φ_h gives masses to the fermions of the fourth family only, whereas Φ_ℓ gives masses to the remaining fermions.

4GTHDM

- Flavor changing neutral currents (FCNCs) in the 4GTHDM arise at the tree level in the scalar sector and in general can be written as

$$\mathcal{L} = \frac{g}{2m_W} f^\phi \bar{q}_i \left(S_{ij}^\phi + P_{ij}^\phi \gamma_5 \right) q_j \phi + \text{H.c.},$$

Where i (j) run over up (down) quarks type for h^0 , H^0 and A^0 , while for H^\pm runs over up (down) quarks.

ϕ	f^ϕ	S_{ij}^ϕ	P_{ij}^ϕ
h^0	$\frac{c_\alpha}{s_\beta} + \frac{s_\alpha}{c_\beta}$	$\frac{m_{q_i}}{f_\beta} \frac{s_\alpha}{c_\beta} \delta_{ij} - f_{ij}^+$	$-f_{ij}^-$
H^0	$\frac{c_\alpha}{c_\beta} - \frac{s_\alpha}{s_\beta}$	$-\frac{m_{q_i}}{f_\beta} \frac{c_\alpha}{c_\beta} \delta_{ij} + f_{ij}^+$	f_{ij}^-
A^0	$2iI_q(t_\beta + \frac{1}{t_\beta})$	f_{ij}^-	$-\frac{m_{q_i}}{f_\beta} t_\beta \delta_{ij} + f_{ij}^+$
H^\pm	$\frac{2}{\sqrt{2}}$	$\frac{1}{2}t_\beta U_{ij}(m_{d_j} - m_{u_i}) + h_{ij}^-$	$\frac{1}{2}t_\beta U_{ij}(m_{d_j} + m_{u_i}) - h_{ij}^+$

- U_{ij} are entries of the new 4x4 CKM matrix, while I_q is the weak isospin.

4GTHDM

- $f_{ij}^{\pm} = \frac{1}{2} (m_{q_i} \Sigma_{ij}^q \pm m_{q_j} \Sigma_{ji}^{q*})$, with $q = u$ (d) for up (down) quarks.


$$h_{ij}^{\pm} = \frac{1}{2} (t_{\beta} + \frac{1}{t_{\beta}}) (m_{u_k} \Sigma_{ki}^{u*} U_{kj} \pm m_{d_k} \Sigma_{kj}^d U_{ik})$$

- $\Sigma_{ij}^{u,d}$ are entries of the new complex 4x4 mixing matrices.

$$\Sigma^{u,d} \simeq \left(\begin{array}{c|cc} \mathbf{0} & & \mathbf{0} \\ \hline \mathbf{0} & |\epsilon_{u,d}|^2 & \epsilon_{u,d}^* \left(1 - \frac{|\epsilon_{u,d}|^2}{2} \right) \\ & \epsilon_{u,d} \left(1 - \frac{|\epsilon_{u,d}|^2}{2} \right) & \left(1 - \frac{|\epsilon_{u,d}|^2}{2} \right) \end{array} \right),$$

CMDM AND CEDM IN THE 4GTHDM

- The CMDM and CEDM of the top quark arise through the generic Feynman diagram



- Where $Q = t, t'$ for h^0, H^0 and A^0 , whereas $Q = b, b'$ for H^\pm .

CMDM AND CEDM IN THE 4GTHDM

- The contributions to CMDM and CEDM are

$$a_t^\phi(m_Q) = \left(\frac{g}{2r_W} \right)^2 \frac{|f_\phi|^2}{8\pi^2} \left(|\tilde{S}_{tQ}^\phi|^2 F(r_Q, r_\phi) + |\tilde{P}_{tQ}^\phi|^2 F(-r_Q, r_\phi) \right),$$

$$d_t^\phi(m_Q) = \frac{g_s}{m_t} \left(\frac{g}{2r_W} \right)^2 \frac{|f_\phi|^2}{8\pi^2} \text{Im}(\tilde{S}_{tQ}^\phi \tilde{P}_{tQ}^{\phi*}) G(r_Q, r_\phi),$$

- Where

$$F(x, y) = \int_0^1 dz \frac{(1-z)^2(z+x)}{(1-z)(x^2-z) + zy^2},$$

$$G(x, y) = x \int_0^1 dz \frac{(1-z)^2}{(1-z)(x^2-z) + zy^2},$$

$$r_a = m_a/m_t, \quad \tilde{S}_{ij}^\phi = S_{ij}^\phi/m_t, \quad \text{and} \quad \tilde{P}_{ij}^\phi = P_{ij}^\phi/m_t$$

CMDM AND CEDM IN THE 4GTHDM

- The equation for CMDM contribution can be reduced to SM contribution

$$a_t^{h_{\text{SM}}^0} = \frac{G_F m_t^2}{4\sqrt{2}\pi^2} \int_0^1 dz \frac{(1+z)(1-z)^2}{(1-z)^2 + z r_{h_{\text{SM}}^0}^2},$$

- We obtain

$$a_t^{h_{\text{SM}}^0} = 3.78 \times 10^{-3}$$

CMDM AND CEDM IN THE 4GTHDM

- In our numerical analysis the parameter space is taken as

Parameter	Value
t_β	1
χ	0.1
$m_{t'}$	400 – 1000 GeV
Δm_4	100 GeV
m_ϕ	400 – 1000 GeV
$ U_{tb} , U_{t'b'} $	0.99
$ U_{t'b} , U_{tb'} $	0.1
$\rho_{t'b}$	0
$ \epsilon_t , \epsilon_b $	0.5, 0.05 (scenario I)
$ \epsilon_t , \epsilon_b $	0.05, 0.05 (scenario II)
$\rho_{tb'}, \delta_t, \delta_b$	$\pi/2, \pi/4, \pi/4$ (scenarios I and II)

CMDM AND CEDM IN THE 4GTHDM

- The new physics contribution is given as follows

$$\delta a_t^{4GTHDM} = a_t^{3rd} + a_t^{4th},$$


- Where

$$a_t^{3rd} = \delta a_t^{h^0}(m_t) + \sum_{\phi=H^0, A^0} a^\phi(m_t) + a^{H^\pm}(m_b),$$


$$a_t^{4th} = \sum_{\phi=h^0, H^0, A^0} a^\phi(m_{t'}) + a^{H^\pm}(m_{b'}).$$

$$\delta a_t^{h^0}(m_t) = a_t^{h^0}(m_t) - a_t^{h^0_{SM}}$$

CMDM AND CEDM IN THE 4GTHDM



CMDM AND CEDM IN THE 4GTHDM



CONCLUSIONS

- A fourth generation of fermions is still consistent with the 125 GeV Higgs measured in 2012 in the framework of the 4GTHDM.
- New sources of CP violation are required in order to explain baryon asymmetry of the universe.
- We obtain contributions of order $10^{-4} - 10^{-3}$ for the CMDM and order $10^{-18} - 10^{-19} e \cdot cm$ for the CEDM.
- The most recent bound to the CEDM is $10^{-18} e \cdot cm$, so our result is very close. ([PhysRevD.88.034033](#))
- The fourth family gives a high contribution to the CMDM and CEDM.

THANK YOU