

Lepton-pair production in di-pion τ lepton decays

Content

We study the $\tau \rightarrow \nu_\tau \pi^- \pi^0 \ell^+ \ell^-$ ($\ell=e, \mu$) decays, which are $O(\alpha^2)$ -suppressed with respect to the dominant di-pion tau decay channel. Both the inner-bremsstrahlung and the structure- (and model-)dependent contributions are considered. In the $\ell=e$ case, structure-dependent effects are $O(1\%)$ in the decay rate, yielding a clean prediction of its branching ratio, 2.3×10^{-5} , measurable with BaBar or Belle(-II) data. For $\ell=\mu$, both contributions have similar magnitude and we get a branching fraction of $(1.6 \pm 0.3) \times 10^{-7}$, reachable by the end of Belle-II operation. These decays allow to study the dynamics of strong interactions with simultaneous weak and electromagnetic probes; their knowledge will contribute to reducing backgrounds in lepton flavor/number violating searches.

Summary

Primary author(s) : Mr. GUTIÉRREZ SANTIAGO, Jorge Luis (Cinvestav)

Co-author(s) : Dr. LOPEZ CASTRO, Gabriel (cinvestav); Dr. ROIG, Pablo (Cinvestav)

Presenter(s) : Mr. GUTIÉRREZ SANTIAGO, Jorge Luis (Cinvestav)