

# Charmed baryon decays at BESIII

Xiao-Rui Lyu

University of Chinese Academy of Sciences, Beijing

(On behalf of the BESIII collaboration)

10th International Workshop on Charm Physics (CHARM 2020)

from 31 May 2021 to 5 June 2021  
Asia/Shanghai timezone



# Outline


- Physics related to the charmed baryon
- Recent results on  $\Lambda_c$  decays at BESIII
- Future prospects
- Summary

# Why $\Lambda_c^+$ is interesting



- An important intermediate particle:
  - **corner stone** of the charmed baryon spectra
  - many b-baryon decays to  $\Lambda_c$
- Its decays reveal information of strong- and weak-interactions in charm region, complementary to D/Ds





$\Xi_c^+$ : relative to the decay of  $\Xi^- 2\pi^+$

Mode	Fraction ( $\Gamma_i / \Gamma$ )
No absolute branching fractions have been measured. The following are branching to $\Xi^- \pi^+$ . Cabibbo-favored ( $S = -2$ ) decays – relative to $\Xi^- \pi^+$	
$\Gamma_1$ $p 2 K_S^0$	$0.087 \pm 0.021$
$\Gamma_2$ $\Lambda \bar{K}^0 \pi^+$	
$\Gamma_3$ $\Sigma(1385)^+ \bar{K}^0$	$1.0 \pm 0.5$
$\Gamma_4$ $\Lambda K^- 2 \pi^+$	$0.323 \pm 0.033$
$\Gamma_5$ $\Lambda \bar{K}^* (892)^0 \pi^+$	$< 0.16$
$\Gamma_6$ $\Sigma(1385)^+ K^- \pi^+$	$< 0.23$
$\Gamma_7$ $\Sigma^+ K^- \pi^+$	$0.94 \pm 0.10$
$\Gamma_8$ $\Sigma^+ \bar{K}^* (892)^0$	$0.81 \pm 0.15$
$\Gamma_9$ $\Sigma^0 K^- 2 \pi^+$	$0.27 \pm 0.12$
$\Gamma_{10}$ $\Xi^0 \pi^+$	$0.55 \pm 0.16$
$\Gamma_{11}$ $\Xi^- 2 \pi^+$	<b>DEFINED AS 1</b>

### Decay Modes $\Omega_c^0$



Mode	Fraction ( $\Gamma_i / \Gamma$ )
No absolute branching fractions have been measured. The following are branching ratios relative to $\Omega_c^0 \rightarrow \Xi^- \pi^+$ . Cabibbo-favored ( $S = -3$ ) decays – relative to $\Omega_c^0 \rightarrow \Xi^- \pi^+$	
$\Gamma_6$ $\Xi^0 \bar{K}^0$	$1.64 \pm 0.29$
$\Gamma_7$ $\Xi^0 K^- \pi^+$	$1.20 \pm 0.18$
$\Gamma_8$ $\Xi^0 \bar{K}^{*0}, \bar{K}^{*0} \rightarrow K^- \pi^+$	$0.68 \pm 0.16$
$\Gamma_9$ $\Xi^- \bar{K}^0 \pi^+$	$2.12 \pm 0.28$
$\Gamma_{10}$ $\Xi^- K^- 2 \pi^+$	$0.63 \pm 0.09$
$\Gamma_{11}$ $\Xi(1530)^0 K^- \pi^+, \Xi^{*0} \rightarrow \Xi^- \pi^+$	$0.21 \pm 0.06$
$\Gamma_{12}$ $\Xi^- \bar{K}^{*0} \pi^+$	$0.34 \pm 0.11$
$\Gamma_{13}$ $\Sigma^+ K^- K^- \pi^+$	$< 0.32$
$\Gamma_{14}$ $\Lambda \bar{K}^0 \bar{K}^0$	$1.72 \pm 0.35$



**2014 : 0.567 fb<sup>-1</sup> at 4.6 GeV**

corresponds to 0.1M  $\Lambda_c$  pairs

- $E_{\text{cms}} - 2M_{\Lambda_c} = 26\text{MeV}$  only!
- $\Lambda_c^+ \Lambda_c^-$  produced in pairs with no additional accompany hadrons.
  - $e^+e^- \rightarrow \gamma^* \rightarrow \Lambda_c^+ \Lambda_c^-$
- Clean backgrounds and well constrained kinematics.
- Typically, two ways to study  $\Lambda_c^+$  decays:
  - **Single Tag(ST):** detect only one of the  $\Lambda_c^+ \Lambda_c^-$ .
    - =>Relative higher backgrounds
    - =>Higher efficiencies
    - =>Full reconstruction
  - **Double Tag(DT):** detect both of  $\Lambda_c^+ \Lambda_c^-$ 
    - =>Smaller backgrounds.
    - =>Missing technique.
    - =>Lower efficiencies.
    - =>Systematic in tag side are mostly cancelled.



*Hadronic decay***2014 : 0.567 fb<sup>-1</sup> at 4.6 GeV** $\Lambda_c^+ \rightarrow pK^- \pi^+ + 11 \text{ CF modes}$ 

PRL 116, 052001 (2016)

 $\Lambda_c^+ \rightarrow pK^+ K^-, p\pi^+ \pi^-$ 

PRL 117, 232002 (2016)

 $\Lambda_c^+ \rightarrow nK_s \pi^+$ 

PRL 118, 12001 (2017)

 $\Lambda_c^+ \rightarrow p\eta, p\pi^0$ 

PRD 95, 111102(R) (2017)

 $\Lambda_c^+ \rightarrow \Sigma^- \pi^+ \pi^+ \pi^0$ 

PLB 772, 388 (2017)

 $\Lambda_c^+ \rightarrow \Xi^{0(*)} K^+$ 

PLB783, 200 (2018)

 $\Lambda_c^+ \rightarrow \Lambda \eta \pi^+$ **PRD99, 032010 (2019)** $\Lambda_c^+ \rightarrow \Sigma^+ \eta, \Sigma^+ \eta'$ **CPC43, 083002 (2019)** $\Lambda_c^+ \rightarrow \text{BP decay asymmetries}$ **PRD100, 072004 (2019)** $\Lambda_c^+ \rightarrow pK_s \eta$ **PLB 817, 136327 (2021)** $\Lambda_c^+$  spin determination**PRD 103, L091101(2021)***Semi-leptonic decay* $\Lambda_c^+ \rightarrow \Lambda e^+ \nu_e$ 

PRL 115, 221805(2015)

 $\Lambda_c^+ \rightarrow \Lambda \mu^+ \nu_\mu$ 

PLB 767, 42 (2017)

*Inclusive decay* $\Lambda_c^+ \rightarrow \Lambda X$ 

PRL121, 062003 (2018)

 $\Lambda_c^+ \rightarrow e^+ X$ **PRL 121 251801(2018)** $\Lambda_c^+ \rightarrow K_s^0 X$ **EPJC 80, 935 (2020)***Production* $\Lambda_c^+ \Lambda_c^-$  cross section




PRL 120,132001(2018)

# $\Lambda_c^+ \rightarrow \Lambda \eta \pi^+$

PRD 99, 032010,(2019)

- current world result has large uncertainty
- potential to study intermediate states, such as  $a_0(980)$  and  $\Lambda(1670)$

Decay	CLEO in 1995	CLEO in 2003	PDG average
$B(\Lambda_c^+ \rightarrow \Lambda \eta \pi^+)/B(\Lambda_c^+ \rightarrow pK^- \pi^+)$	$0.35 \pm 0.05 \pm 0.06$	$0.41 \pm 0.17 \pm 0.10$	$0.36 \pm 0.07$
$B(\Lambda_c^+ \rightarrow \Lambda \eta \pi^+)$	----	----	$(2.3 \pm 0.5)\%$





- branching fraction  $B(\Lambda_c^+ \rightarrow \Lambda \eta \pi^+)$  measured to be  $(1.84 \pm 0.21 \pm 0.15)\%$  more precise than previous results
- $B(\Lambda_c^+ \rightarrow \Sigma^{*+} \eta)$  measured as  $(0.91 \pm 0.08 \pm 0.09)\%$  more precise than the previous result  $(1.24 \pm 0.37)\%$

# $\Lambda_c^+ \rightarrow \Sigma^+ \eta, \Sigma^+ \eta'$

CPC 43, 083002, (2019)

- Decay through internal W-emission and W-exchange.
- Both are non-factorable in theoretic calculation.



$$\frac{\mathcal{B}(\Lambda_c^+ \rightarrow \Sigma^+ \eta)}{\mathcal{B}(\Lambda_c^+ \rightarrow \Sigma^+ \pi^0)} = 0.35 \pm 0.16 \pm 0.03$$

$$\frac{\mathcal{B}(\Lambda_c^+ \rightarrow \Sigma^+ \eta')}{\mathcal{B}(\Lambda_c^+ \rightarrow \Sigma^+ \omega)} = 0.86 \pm 0.34 \pm 0.07$$

$$\frac{\mathcal{B}(\Lambda_c^+ \rightarrow \Sigma^+ \eta')}{\mathcal{B}(\Lambda_c^+ \rightarrow \Sigma^+ \eta)} = 3.5 \pm 2.1 \pm 0.4$$

Our measurement contradict with most theoretical calculations.







Decay mode	Körner [5]	Sharma [3]	Zenczykowski [4]	Ivanov [6]	CLEO [12]	This work
$\Lambda_c^+ \rightarrow \Sigma^+ \eta$	0.16	0.57	0.94	0.11	$0.70 \pm 0.23$	$0.41 \pm 0.20$ ( $< 0.68$ )
$\Lambda_c^+ \rightarrow \Sigma^+ \eta'$	1.28	0.10	0.12	0.12	-	$1.34 \pm 0.57$ ( $< 1.9$ )



- Previous CLEO result of  $B(\Lambda_c^+ \rightarrow p K_S^0 \eta) = (0.8 \pm 0.2)\%$ , while theoretical calculations based on SU(3) symmetry give  $(0.35 \sim 0.45)\%$
- A potential channel to study a puzzling  $N(1535)$  which has nontrivial decay rate of  $\eta N$  and  $K \Lambda$
- 2D fit to  $M_{BC}$  and  $\Delta E$  distributions are implemented

single tag method

significance  $5.3 \sigma$



$42.0 \pm 8.5$  signals

$B(\Lambda_c^+ \rightarrow p K_S^0 \eta) = (0.414 \pm 0.084 \pm 0.028)\%$

compatible with CLEO result within  $1.5 \sigma$





single tag method

PRD100, 072004 (2019)

4(6)-fold angular analysis of the cascade decays of  $\Lambda_c^+ \rightarrow pK_S, \Lambda\pi^+, \Sigma^+\pi^0$  and  $\Sigma^0\pi^+$  based on 567/pb data

$$\mathcal{W} \propto 1 + \alpha_0 \cos^2 \theta_0 + \mathcal{P}_T \alpha_{BP}^+ \sin \theta_1 \sin \phi_1$$

$$\mathcal{P}_T(\cos \theta_0) \equiv \sqrt{1 - \alpha_0^2} \cos \theta_0 \sin \theta_0 \sin \Delta_0$$






$$\frac{dN}{d \cos \theta_2} \propto 1 + \alpha_{\Lambda\pi^+(\Sigma^+\pi^0)}^+ \alpha_{\Lambda(\Sigma^+)} \cos \theta_2.$$

$\Lambda_c^+ \rightarrow$	$pK_S^0$	$\Lambda\pi^+$	$\Sigma^+\pi^0$	$\Sigma^0\pi^+$	
$\alpha_{BP} = \frac{2\text{Re}(s^*p)}{ s ^2 +  p ^2}$	Predicted	-1.0 [16], 0.51 [11]	-0.70 [16], -0.67 [11]	0.71 [16], 0.92 [11]	0.70 [16], 0.92 [11]
		-0.49 [10], -0.90 [10]	-0.95 [10], -0.99 [10]	0.79 [10], -0.49 [10]	0.78 [10], -0.49 [10]
		-0.49 [17], -0.97 [18]	-0.96 [17], -0.95 [18]	0.83 [17], 0.43 [18]	0.83 [17], 0.43 [18]
		-0.66 [19], -0.90 [30]	-0.99 [19], -0.86 [30]	0.39 [19], -0.76 [30]	0.39 [19], -0.76 [30]
		-0.99 [20], -0.91 [31]	-0.99 [20], -0.94 [31]	-0.31 [20], -0.47 [31]	-0.31 [20], -0.47 [31]
PDG [2]		$-0.91 \pm 0.15$	$-0.45 \pm 0.32$		
This work	$0.18 \pm 0.43 \pm 0.14$	$-0.80 \pm 0.11 \pm 0.02$	$-0.57 \pm 0.10 \pm 0.07$	$-0.73 \pm 0.17 \pm 0.07$	

$$\sin \Delta_0 = -0.28 \pm 0.13 \pm 0.03$$

- Best precisions on the hadronic weak decay asymmetries
- No theoretical models fully describe the new BESIII results
- The transverse polarization is firstly studied and found to be non-zero with  $2.1\sigma$

- No spin-determination of the  $\Lambda_c$  since first discovery more than 30 years ago
- Currently, the spin half of the  $\Lambda_c$  is inferred to be from the naive quark model
- It would be crucial to test this spin assignment in experiment, to test the quark model hadron classification
- Multi-dimensional angular analysis on the ST samples of  $\Lambda_c \rightarrow pK_S, \Lambda\pi^+, \Sigma^+\pi^0$  and  $\Sigma^0\pi^+$  are carried out to test both hypotheses of  $J=1/2$  and  $3/2$




$J=1/2$  is preferred over  $J=3/2$   
with a significance of  $6\sigma$

- consistent with the expectation of the naive quark model.
- a cornerstone in the extraction of the properties of heavier charmed and beauty baryons

# Inclusive decay $\Lambda_c^+ \rightarrow e^+ X$

PRL121, 251801(2018)

- Difference between  $B(\Lambda_c \rightarrow \Lambda e^+ \nu)$  and  $B(\Lambda_c \rightarrow e^+ X)$  can shed light on searching for new semi-leptonic mode
- Two clean tag modes are used
- PID unfolding is implemented
- Improved results from Mark II's




$\Lambda_c^+ \rightarrow X e^+ \nu_e$	Right sign	Wrong sign
Observed yields		
Tag signal region	$228.0 \pm 15.1$	$26.0 \pm 5.1$
Tag sideband region	$11.0 \pm 3.3$	$2.0 \pm 1.4$
PID unfolding		
Tag signal region	$250.1 \pm 17.1$	$28.3 \pm 6.2$
Tag sideband region	$12.1 \pm 3.8$	$1.7 \pm 1.5$
Sideband subtraction	$240.7 \pm 17.4$	$27.0 \pm 6.3$
Wrong-sign subtraction	$213.7 \pm 18.5$	
Correction of tracking efficiency	$272.1 \pm 23.5$	

✓ **BESIII results:**

$$B(\Lambda_c^+ \rightarrow X e^+ \nu_e) = (3.95 \pm 0.34 \pm 0.09)\%$$

$$\frac{B(\Lambda_c^+ \rightarrow \Lambda e^+ \nu_e)}{B(\Lambda_c^+ \rightarrow X e^+ \nu_e)} = (91.9 \pm 12.5 \pm 5.4)\%$$

$$\frac{\Gamma(\Lambda_c^+ \rightarrow X e^+ \nu_e)}{\Gamma(D \rightarrow X e^+ \nu_e)} = 1.26 \pm 0.12 \quad \text{consistent with theoretical predictions}$$




# Inclusive decay $\Lambda_c^+ \rightarrow K_S^0 X$

- Sum BF for exclusive channels with  $K^0$  or  $\bar{K}^0$  is evaluated to be  $(22.4 \pm 0.9)\%$  with inclusion of guessed modes
- Its difference from the inclusive rate  $B(\Lambda_c^+ \rightarrow K^0/\bar{K}^0 X)$  will help in identifying unknown modes
- Eleven hadronic tag modes are used

EPJC 80, 935 (2020)

Mode	Value (%)	Mode	Value (%)
Observed BF		Extrapolated BF	
$p\bar{K}^0$	$3.18 \pm 0.16$	$n\bar{K}^0\pi^+\pi^0$	$3.07 \pm 0.16$
$p\bar{K}^0\pi^0$	$3.94 \pm 0.26$	$p\bar{K}^0\pi^0\pi^0$	$1.36 \pm 0.07$
$p\bar{K}^0\pi^+\pi^-$	$3.20 \pm 0.24$	$n\bar{K}^0\pi^+\pi^+\pi^-$	$0.14 \pm 0.09$
$n\bar{K}^0\pi^+$	$3.64 \pm 0.50$	$p\bar{K}^0\pi^+\pi^-\pi^0$	$0.22 \pm 0.14$
$p\bar{K}^0\eta$	$1.60 \pm 0.40$	$n\bar{K}^0\pi^+\pi^0\pi^0$	$0.10 \pm 0.06$
$\Lambda K^+\bar{K}^0$	$0.57 \pm 0.11$	$p\bar{K}^0\pi^0\pi^0\pi^0$	$0.03 \pm 0.02$
		in statistical isospin model $(\Sigma K)^+\bar{K}^0$	$0.68 \pm 0.34$
		<i>PRD97, 116015 (2018)</i>	
		$\Xi^0 K^0\pi^+$	$0.62 \pm 0.06$
Total	$16.1 \pm 0.8$	Total	$6.3 \pm 0.4$
Total		Total	$22.4 \pm 0.9$




$$B(\Lambda_c^+ \rightarrow K_S^0 X) = (9.9 \pm 0.6 \pm 0.4)\%$$

$$B(\Lambda_c^+ \rightarrow K^0/\bar{K}^0 X) = (19.8 \pm 1.2 \pm 0.8 \pm 1.0)\%$$

- 2D fit
- signal yields:  $478 \pm 27$
- third error due to isospin breaking for  $\Lambda_c^+$  decaying to  $K_S$  or  $K_L$  in final states
- compatible with the estimated exclusive rate

# Heavier charmed baryons



Energy thresholds

✓	$\Lambda_c^+ \bar{\Sigma}_c^-$	4.74 GeV
✓	$\Lambda_c^+ \bar{\Sigma}_c \pi$	4.88 GeV
✓	$\Sigma_c \bar{\Sigma}_c$	4.91 GeV
✓	$\Xi_c \bar{\Xi}_c$	4.95 GeV
✓	$\Omega_c^0 \bar{\Omega}_c^0$	5.4 GeV


	Structure	$J^P$	Mass, MeV	Width, MeV	Decay
$\Lambda_c^+$	$udc$	$(1/2)^+$	$2286.46 \pm 0.14$	$(200 \pm 6)$ fs	weak
$\Xi_c^+$	$usc$	$(1/2)^+$	$2467.8^{+0.4}_{-0.6}$	$(442 \pm 26)$ fs	weak
$\Xi_c^0$	$dsc$	$(1/2)^+$	$2470.88^{+0.34}_{-0.8}$	$112^{+13}_{-10}$ fs	weak
$\Sigma_c^{++}$	$uuc$	$(1/2)^+$	$2454.02 \pm 0.18$	$2.23 \pm 0.30$	$\Lambda_c^+ \pi^+$
$\Sigma_c^+$	$udc$	$(1/2)^+$	$2452.9 \pm 0.4$	$< 4.6$	$\Lambda_c^+ \pi^0$
$\Sigma_c^0$	$ddc$	$(1/2)^+$	$2453.76 \pm 0.18$	$2.2 \pm 0.4$	$\Lambda_c^+ \pi^-$
$\Xi_c^+$	$usc$	$(1/2)^+$	$2575.6 \pm 3.1$	—	$\Xi_c^+ \gamma$
$\Xi_c^0$	$dsc$	$(1/2)^+$	$2577.9 \pm 2.9$	—	$\Xi_c^0 \gamma$
$\Omega_c^0$	$ssc$	$(1/2)^+$	$2695.2 \pm 1.7$	$(69 \pm 12)$ fs	weak
$\Sigma_c^{*++}$	$uuc$	$(3/2)^+$	$2518.4 \pm 0.6$	$14.9 \pm 1.9$	$\Lambda_c^+ \pi^+$
$\Sigma_c^{*+}$	$udc$	$(3/2)^+$	$2517.5 \pm 2.3$	$< 17$	$\Lambda_c^+ \pi^0$
$\Sigma_c^{*0}$	$ddc$	$(3/2)^+$	$2518.0 \pm 0.5$	$16.1 \pm 2.1$	$\Lambda_c^+ \pi^-$
$\Xi_c^{*+}$	$usc$	$(3/2)^+$	$2645.9^{+0.5}_{-0.6}$	$< 3.1$	$\Xi_c \pi$
$\Xi_c^{*0}$	$dsc$	$(3/2)^+$	$2645.9 \pm 0.5$	$< 5.5$	$\Xi_c \pi$
$\Omega_c^{*0}$	$ssc$	$(3/2)^+$	$2765.9 \pm 2.0$	—	$\Omega_c^0 \gamma$

# BESII New data samples in 2020 and 2021



Two major changes in BEPCII machine:

- max beam energy: 2.30 → 2.35(2018) → 2.48 GeV(2020)
- top-up injection: data taking efficiency increases by 20~30%





## Available data for charmed baryon

- ✓ 0.567 fb<sup>-1</sup> at 4.6 GeV (35 days in 2014)
- ✓ 3.8 fb<sup>-1</sup> scan at 4.61, 4.63, 4.64, 4.66, 4.68, 4.7 GeV (186 days in 2020)
- ✓ 2 fb<sup>-1</sup> scan at 4.74, 4.78, 4.84, 4.91, 4.95 GeV (99 days in 2021)

~ 10x  $\Lambda_c$  data than those at 4.6 GeV; accessible to  $\Sigma_c/\Xi_c$  prod. & decays

- optimized energy at 2.35 GeV with luminosity 3 times higher than the current BEPCII.



4.95 ~ 5.6 GeV: new energy coverage of BEPCII-upgrade





- No absolute BF's have been measured/calculated until 2019
- Belle measured abs. BF's in 2019, but uncertainties are large:  $\delta B \sim 30\%$

## $\Xi_c^+$ : relative to the decay of $\Xi^- 2\pi^+$

No absolute branching fractions have been measured. The following are branching to  $\Xi^- \pi^+$ . Cabibbo-favored ( $S = -2$ ) decays – relative to  $\Xi^- \pi^+$

Mode	Fraction ( $\Gamma_i / \Gamma$ )
$\Gamma_1$ $p 2 K_S^0$	$0.087 \pm 0.021$
$\Gamma_2$ $\Lambda \bar{K}^0 \pi^+$	
$\Gamma_3$ $\Sigma(1385)^+ \bar{K}^0$	$1.0 \pm 0.5$
$\Gamma_4$ $\Lambda K^- 2 \pi^+$	$0.323 \pm 0.033$
$\Gamma_5$ $\Lambda \bar{K}^*(892)^0 \pi^+$	$< 0.16$
$\Gamma_6$ $\Sigma(1385)^+ K^- \pi^+$	$< 0.23$
$\Gamma_7$ $\Sigma^+ K^- \pi^+$	$0.94 \pm 0.10$
$\Gamma_8$ $\Sigma^+ \bar{K}^*(892)^0$	$0.81 \pm 0.15$
$\Gamma_9$ $\Sigma^0 K^- 2 \pi^+$	$0.27 \pm 0.12$
$\Gamma_{10}$ $\Xi^0 \pi^+$	$0.55 \pm 0.16$
$\Gamma_{11}$ $\Xi^- 2 \pi^+$	<b>DEFINED AS 1</b>
$\Gamma_{12}$ $\Xi(1530)^0 \pi^+$	$< 0.10$
$\Gamma_{13}$ $\Xi^0 \pi^+ \pi^0$	$2.3 \pm 0.7$
$\Gamma_{14}$ $\Xi^0 \pi^- 2 \pi^+$	$1.7 \pm 0.5$
$\Gamma_{15}$ $\Xi^0 e^+ \nu_e$	$2.3^{+0.7}_{-0.8}$
$\Gamma_{16}$ $\Omega^- K^+ \pi^+$	$0.07 \pm 0.04$
<b>Cabibbo-suppressed</b>	
$\Gamma_{17}$	
$\Gamma_{18}$	
$\Gamma_{19}$	
$\Gamma_{20}$	
$\Gamma_{21}$ $\Sigma^+ K^+ K^-$	$0.15 \pm 0.06$

## Decay Modes $\Xi_c^0$

Mode	Fraction ( $\Gamma_i / \Gamma$ )
<b>Cabibbo-favored (<math>S = -2</math>) decays</b>	
$\Gamma_1$ $p K^- K^- \pi^+$	$(4.8 \pm 1.2) \times 10^{-3}$
$\Gamma_2$ $p K^- \bar{K}^*(892)^0, \bar{K}^{*0} \rightarrow K^- \pi^+$	$(2.0 \pm 0.6) \times 10^{-3}$
$\Gamma_3$ $p K^- K^- \pi^+$ (no $\bar{K}^{*0}$ )	$(3.0 \pm 0.9) \times 10^{-3}$
$\Gamma_4$ $\Lambda K_S^0$	$(3.0 \pm 0.8) \times 10^{-3}$
$\Gamma_5$ $\Lambda K^- \pi^+$	$(1.45 \pm 0.33)\%$
$\Gamma_6$ $\Lambda \bar{K}^0 \pi^+ \pi^-$	seen
$\Gamma_7$ $\Lambda K^- \pi^+ \pi^+ \pi^-$	seen
$\Gamma_8$ $\Xi^- \pi^+$	$(1.43 \pm 0.32)\%$
$\Gamma_9$ $\Xi^- \pi^+ \pi^+ \pi^-$	$(4.8 \pm 2.3)\%$
$\Gamma_{10}$ $\Omega^- K^+$	$(4.2 \pm 1.0) \times 10^{-3}$
$\Gamma_{11}$ $\Xi^- e^+ \nu_e$	$(1.8 \pm 1.2)\%$
<b>Cabibbo-suppressed decays</b>	
$\Gamma_{12}$ $\Xi^- K^+$	$(3.9 \pm 1.2) \times 10^{-4}$
$\Gamma_{13}$ $\Lambda K^+ K^-$ (no $\phi$ )	$(4.1 \pm 1.4) \times 10^{-4}$
$\Gamma_{14}$ $\Lambda \phi$	$(4.9 \pm 1.5) \times 10^{-4}$

Very limited knowledge on their decays  
 We have opportunity to systematically study more decays

# Studies on the $\Omega_c^0$

*Mode*

*Fraction ( $\Gamma_i / \Gamma$ )*

▼ **No absolute branching fractions have been measured. The following are branching ratios Cabibbo-favored ( $S = -3$ ) decays – relative to  $\Omega^- \pi^+$**

$\Gamma_i$	Mode	Fraction ( $\Gamma_i / \Gamma$ )
$\Gamma_1$	$\Omega^- \pi^+$	<b>DEFINED AS 1</b>
$\Gamma_2$	$\Omega^- \pi^+ \pi^0$	$1.80 \pm 0.33$
$\Gamma_3$	$\Omega^- \rho^+$	$> 1.3$
$\Gamma_4$	$\Omega^- \pi^- 2 \pi^+$	$0.31 \pm 0.05$
$\Gamma_5$	$\Omega^- e^+ \nu_e$	$2.4 \pm 1.2$
$\Gamma_6$	$\Xi^0 \bar{K}^0$	$1.64 \pm 0.29$
$\Gamma_7$	$\Xi^0 K^- \pi^+$	$1.20 \pm 0.18$
$\Gamma_8$	$\Xi^0 \bar{K}^{*0}, \bar{K}^{*0} \rightarrow K^- \pi^+$	$0.68 \pm 0.16$
$\Gamma_9$	$\Xi^- \bar{K}^0 \pi^+$	$2.12 \pm 0.28$
$\Gamma_{10}$	$\Xi^- K^- 2 \pi^+$	$0.63 \pm 0.09$
$\Gamma_{11}$	$\Xi(1530)^0 K^- \pi^+, \Xi^{*0} \rightarrow \Xi^- \pi^+$	$0.21 \pm 0.06$
$\Gamma_{12}$	$\Xi^- \bar{K}^{*0} \pi^+$	$0.34 \pm 0.11$
$\Gamma_{13}$	$\Sigma^+ K^- K^- \pi^+$	$< 0.32$
$\Gamma_{14}$	$\Lambda \bar{K}^0 \bar{K}^0$	$1.72 \pm 0.35$

# Studies on most of the $\Xi_c / \Omega_c$ weak decays are missing in experiment (I)

## BFs of CF decays

RQM Pole Pole RQM Pole Pole (in units of %)

Decay	Körner, Krämer ('92)	Xu, Kamal ('92)	Cheng, Tseng ('93)	Ivanov et al. ('98)	Żenczykowski ( '94)	Sharma, Verma ('99)	Expt.
$\Xi_c^+ \rightarrow \Sigma^+ \bar{K}^0$	6.45	0.44	0.84	3.08	1.56	0.04	
$\Xi_c^+ \rightarrow \Xi^0 \pi^+$	3.54	3.36	3.93	4.40	1.59	0.53	$0.55 \pm 0.16^a$
$\Xi_c^0 \rightarrow \Lambda \bar{K}^0$	0.12	0.37	0.27	0.42	0.35	0.54	seen
$\Xi_c^0 \rightarrow \Sigma^0 \bar{K}^0$	1.18	0.11	0.13	0.20	0.11	0.07	
$\Xi_c^0 \rightarrow \Sigma^+ K^-$	0.12	0.12		0.27	0.36	0.12	
$\Xi_c^0 \rightarrow \Xi^0 \pi^0$	0.03	0.56	0.28	0.04	0.69	0.87	
$\Xi_c^0 \rightarrow \Xi^0 \eta$	0.24			0.28	0.01	0.22	
$\Xi_c^0 \rightarrow \Xi^0 \eta'$	0.85			0.31	0.09	0.06	
$\Xi_c^0 \rightarrow \Xi^- \pi^+$	1.04	1.74	1.25	1.22	0.61	2.46	seen
$\Omega_c^0 \rightarrow \Xi^0 \bar{K}^0$	1.21		0.09	0.02			

# Studies on most of the $\Xi_c / \Omega_c^0$ weak decays are missing in experiment (II)



## Decay asymmetry $\alpha$ for CF decays

Longitudinal pol. of daughter baryon from unpol. parent baryon

⇒ information on the relative sign between s- and p-waves

Decay	Körner, Krämer ('92)	Xu, Kamal ('92)	Cheng, Tseng ('93)	Ivanov et al. ('98)	Żenczykowski ( '94)	Sharma, Verma ('99)	Expt.
$\Xi_c^+ \rightarrow \Sigma^+ \bar{K}^0$	-1.0	0.24	-0.09	-0.99	1.00	0.54	
$\Xi_c^+ \rightarrow \Xi^0 \pi^+$	-0.78	-0.81	-0.77	-1.0	1.00	-0.27	
$\Xi_c^0 \rightarrow \Lambda \bar{K}^0$	-0.76	1.0	-0.73	-0.75	-0.29	-0.79	
$\Xi_c^0 \rightarrow \Sigma^0 \bar{K}^0$	-0.96	-0.99	-0.59	-0.55	-0.50	0.48	
$\Xi_c^0 \rightarrow \Sigma^+ K^-$	0	0		0	0	0	
$\Xi_c^0 \rightarrow \Xi^0 \pi^0$	0.92	0.92	-0.54	0.94	0.21	-0.80	
$\Xi_c^0 \rightarrow \Xi^0 \eta$	-0.92			-1.0	-0.04	0.21	
$\Xi_c^0 \rightarrow \Xi^0 \eta'$	-0.38			-0.32	-1.00	0.80	
$\Xi_c^0 \rightarrow \Xi^- \pi^+$	-0.38	-0.38	-0.99	-0.84	-0.79	-0.97	$-0.6 \pm 0.4$
$\Omega_c^0 \rightarrow \Xi^0 \bar{K}^0$	0.51		-0.93	-0.81			

Studies on most of the  $\Xi_c / \Omega_c^0$  weak decays are missing in experiment (III)

## Charm-flavor-conserving weak decays

- Light quarks undergo weak transitions, while c quark behaves as a “spectator” e.g.  $\Xi_c \rightarrow \Lambda_c \pi$  ( $s \rightarrow W^- u$ ). Can be studied using HHChPT.

$$\text{Br}(\Xi_c^0 \rightarrow \Lambda_c^+ \pi^-) = 2.9 \times 10^{-4} \quad \mathcal{B}(\Xi_c^0 \rightarrow \pi^- \Lambda_c^+) = (0.55 \pm 0.02 \pm 0.18)\% \quad \text{Larger than theoretical predictions}$$

$$\text{Br}(\Xi_c^+ \rightarrow \Lambda_c^+ \pi^0) = 6.7 \times 10^{-4} \quad \text{[LHCb, PRD 102, 071101 (2020)]}$$

Cheng, Cheung, Lin, Lin, Yan, Yu ('92)

These can be further tested at BESIII

## Semileptonic decays



Process	Pérez-Marcial et al. [85]	Singleton [86]	Cheng, Tseng [81]	Ivanov et al. [87]	Luo [88]	Marques de Carvalho et al. [89]	Huang, Wang [90]	Expt. [3]
$\Xi_c^0 \rightarrow \Xi^- e^+ \nu_e$	18.1 (12.5)	8.5	7.4	8.16	9.7			seen
$\Xi_c^+ \rightarrow \Xi^0 e^+ \nu_e$	18.4 (12.7)	8.5	7.4	8.16	9.7			seen

in units of  $10^{10} \text{ s}^{-1}$

$$\mathcal{B}(\Xi_c^0 \rightarrow \Xi^- e^+ \nu_e) = (1.72 \pm 0.10 \pm 0.12 \pm 0.50)\% \quad \text{[Belle, arXiv:2103.06496]}$$

$$\mathcal{B}(\Xi_c^0 \rightarrow \Xi^- \mu^+ \nu_\mu) = (1.71 \pm 0.17 \pm 0.13 \pm 0.50)\%$$

$$\mathcal{B}_{\text{exp}}(\Xi_c^0 \rightarrow \Xi^- e^+ \nu_e) = 2.43(0.25)(0.35)(0.72)\% \quad \text{[ALICE, PoS ICHEP 2020, 524(2021)]}$$

# Summary

- BESIII has been playing significant role in studying  $\Lambda_c$  decays
- Many new results of  $\Lambda_c$  decays have been published in Charm2018
- BEPCII energy upgrade during 2020-2021 has improved the BESIII capability in  $\Lambda_c$  physics by accumulating more statistics at different energy points and pose opportunity to study  $\Sigma_c/\Xi_c$  physics
- Proposal of BEPCII upgrade (3x luminosity and energy up to 5.6 GeV) will greatly extend the physics opportunities in  $c$ -baryon sector

Thank you!  
谢谢!