

Beyond s-wave interactions of two- and three-meson systems with maximal isospin from lattice QCD

Andrew Hanlon

Brookhaven National Laboratory

In collaboration with Tyler D. Blanton, Ben Hörz,
Colin Morningstar, Fernando Romero-López, and
Stephen R. Sharpe

Outline

1. Motivation


- Test and push the limits of three-particle quantization condition

2. Setup and technical details

- Ensembles, code, analysis
- Three-particle quantization condition

3. Results for pions and kaons

- Three pion masses: 200, 280, 340 MeV
- d-wave interactions constrained



Motivation

- Most QCD resonance decays involve three or more particles
 - $\omega(782) \rightarrow \pi\pi\pi$, $a_1(1260) \rightarrow \pi\pi\pi$, $N(1440) \rightarrow N\pi\pi$
- Many recent developments on the theoretical side (and their applications)
- Three competing formalisms to interpret three-particle finite-volume energies
- Provide real lattice data to test and push the limits of various three-particle formalisms

Lattice setup

- $N_F = 2 + 1$ $O(a)$ -improved Wilson-clover fermions generated by CLS
- Three pion masses allow study of chiral dependence
 - Trace of bare quark masses held fixed
- One lattice spacing, $a = 0.06426(76)$ fm
- Consider constituent momenta up to and including $\mathbf{d}^2 = L^2/(2\pi)^2 \mathbf{P}^2 = 9$




	$(L/a)^3 \times (T/a)$	M_π [MeV]	M_K [MeV]	N_{cfg}	t_{src}	N_{ev}
N203	$48^3 \times 128$	340	440	771	32, 52	192
N200	$48^3 \times 128$	280	460	1712	32, 52	192
D200	$64^3 \times 128$	200	480	2000	35, 92	448

Procedure

1. Calculate matrices of two-point correlation functions
 - a. Use stochastic LapH for quark propagation
 - b. Construct operators to transform in irreps of little group
 - c. Optimize contractions (https://github.com/laphnn/contraction_optimizer)
2. Extract finite-volume energies from correlation matrices
 - a. Solve Generalized EigenValue Problem (GEVP) for correlator matrices
 - b. Fit ratio of rotated correlators to single-exponential to extract shifts from non-interacting
 - c. Reconstruct energies and boost to center-of-momentum frame
3. Obtain K-matrices from spectrum
 - a. Adjust K-matrix parameters until lattice energies match predictions from quantization condition

Single-Meson Energies



- Single-exponential fits to correlators of momentum-projected kaon operators
- Continuum dispersion relation works well up to $d^2 = 9$
- No sign of cutoff effects here
- Similar situation for pions



Spectrum Results on N200

Single-exponential fits to

$$R_n(t) \equiv \frac{v_n^\dagger(\tau_0, \tau_D) C(t) v_n(\tau_0, \tau_D)}{\prod_i C^{(\text{sh})}(\mathbf{p}_i^2, t)}$$



Two- and Three-particle Quantization Conditions

Two-particle QC

$$\det \left[F(E_2, \mathbf{P}, L)^{-1} + \mathcal{K}_2(E_2^*) \right] = 0$$

- F is a purely kinematic known finite-volume function
- $\mathcal{K}_2(E_2^*)_{\ell' m'; \ell m} = \delta_{\ell' \ell} \delta_{m' m} \mathcal{K}_2^{(\ell)}(E_2^*)$ is an infinite-volume quantity with algebraic relation to two-particle scattering amplitude

Three-particle QC

$$\det \left[F_3(E, \mathbf{P}, L)^{-1} + \mathcal{K}_{\text{df},3}(E^*) \right] = 0$$

- F_3 contains both kinematic functions and the two-particle K-matrix
- $\mathcal{K}_{\text{df},3}$ is an infinite-volume quantity but is scheme-dependent
- Must solve integral equation to obtain three-particle scattering amplitude

Fitting the Spectrum

- Parameterization of two-particle K-matrix
 - For s-wave, use the effective range expansion or a form that explicitly includes the Adler zero
 - Use the d-wave scattering length
- Parameterization of $\mathcal{K}_{\text{df},3}$ given by threshold expansion to quadratic order

$$\mathcal{K}_{\text{df},3} = \mathcal{K}_{\text{df},3}^{\text{iso},0} + \mathcal{K}_{\text{df},3}^{\text{iso},1} \Delta + \mathcal{K}_{\text{df},3}^{\text{iso},2} \Delta^2 + \underbrace{\mathcal{K}_A \Delta_A + \mathcal{K}_B \Delta_B}_{\text{Two-particle d-wave contributions}}$$


(see arXiv:1901.07095 for details)

- Parameters $\{p_n\}$ determined from minimum of

$$\chi^2(\{p_n\}) = \sum_{ij} \left(E_i - E_i^{\text{QC}}(\{p_n\}) \right) C_{ij}^{-1} \left(E_j - E_j^{\text{QC}}(\{p_n\}) \right)$$



Testing the Limits of the Formalism

- QC is valid up to first threshold with more than three particles (depending on allowed transitions)
- Transition to $3K\pi$ expected to be NNLO in ChPT, leading to suppressed coupling near threshold
- Fits describe data well above rigorous applicability of QC



Inclusion of d-wave terms

- \mathcal{K}_B and d-wave in \mathcal{K}_2 essential for good fit quality
- Only \mathcal{K}_B contributes to non-trivial irreps, making it easier to constrain
- Can only appear at NLO in ChPT
- Larger error on D200 from large $M_K L$, leading to suppression of energy shifts



Conclusions and Outlooks

- Three-particle quantization condition for simple systems
 - Hundreds of energies extracted
 - d-wave terms in both two- and three-particle K -matrix improve fit quality (substantially in some cases)
 - First calculation showing strong indication that non-zero three-particle interactions are needed
- Future work
 - Spectra for mixed-flavor systems (e.g. $\pi\pi K$ and πKK)
 - Systems with non-maximal isospin, resonances, and/or bound states [GWU arXiv:2107.03973]
 - Integral equations for d-wave

Thanks!

