

# Simultaneous Monte Carlo analysis of parton densities and fragmentation functions

Eric Moffat  
Hadron 2021  
7-14-2021

Collaborators: Wally Melnitchouk, Ted Rogers, Nobuo Sato

*Phys.Rev.D* 104 (2021) 1, 016015

 **Jefferson Lab**  
Thomas Jefferson National Accelerator Facility

 **Jefferson Lab Angular  
Momentum Collaboration**

# Introduction

- Significant tension between large transverse momentum data and Fixed Order (FO) predictions using existing collinear Parton Distribution Functions (PDFs) and Fragmentation Functions (FFs)
- Resolving this tension is crucial for the study of Transverse Momentum Dependent (TMD) PDFs and FFs.
- To facilitate exploring the reasons for this tension in SIDIS, performed a new fit using Jefferson Lab Angular Momentum Collaboration (JAM) methodology:
  - Multi-Step Monte Carlo fit utilizing Bayesian Inference
  - Simultaneously fit PDFs and charged pion, kaon, and unidentified hadron FFs.
    - First such fit involving charged hadrons

# JAM fit

- Simultaneously fit PDFs and charged pion, kaon, and unidentified hadron FFs
  - Functional form:

$$T(x; \mathbf{a}) = \mathcal{M} \frac{x^\alpha (1-x)^\beta (1 + \gamma\sqrt{x} + \delta x)}{\int_0^1 dx x^{\alpha+1} (1-x)^\beta (1 + \gamma\sqrt{x} + \delta x)}$$

- Unidentified charged hadron fragmentation function:

$$D_i^{h^+} = D_i^{\pi^+} + D_i^{K^+} + D_i^{\text{res}^+}$$

# JAM fit

- Multi-Step Monte Carlo fit utilizing Bayesian Inference:

$$\mathcal{P}(\mathbf{a}|\text{data}) \sim \mathcal{L}(\mathbf{a}, \text{data}) \pi(\mathbf{a}) \quad \mathcal{L}(\mathbf{a}, \text{data}) = \exp\left(-\frac{1}{2}\chi^2(\mathbf{a}, \text{data})\right)$$

$$\chi^2(\mathbf{a}) = \sum_{i,e} \left( \frac{d_{i,e} - \sum_k r_e^k \beta_{i,e}^k - T_{i,e}(\mathbf{a})/N_e}{\alpha_{i,e}} \right)^2 + \sum_k (r_e^k)^2 + \left( \frac{1 - N_e}{\delta N_e} \right)^2$$

$$\mathbb{E}[\mathcal{O}] = \int d^d \mathbf{a} \mathcal{P}(\mathbf{a}|\text{data}) \mathcal{O}(\mathbf{a}),$$


$$\mathbb{E}[\mathcal{O}] = \frac{1}{n} \sum_{k=1}^n \mathcal{O}(\mathbf{a}_k),$$

$$\mathbb{V}[\mathcal{O}] = \int d^d \mathbf{a} \mathcal{P}(\mathbf{a}|\text{data}) (\mathcal{O}(\mathbf{a}) - \mathbb{E}[\mathcal{O}])^2$$

$$\mathbb{V}[\mathcal{O}] = \frac{1}{n} \sum_{k=1}^n (\mathcal{O}(\mathbf{a}_k) - \mathbb{E}[\mathcal{O}])^2$$

- Use least squares to obtain maximum likelihood (minimum chi squared) for each replica.

# Multi-step process



# Data sets



- Data Sets:
  - Inclusive Deep Inelastic Scattering (DIS)
    - BCDMS, NMC, SLAC, HERA
  - Semi-Inclusive DIS (SIDIS)
    - COMPASS
  - Single-Inclusive  $e^+/e^-$  Annihilation (SIA)
    - TASSO, TPC, TOPAZ, BELLE, BABAR, ARGUS, DELPHI, ALEPH, OPAL, SLD
  - Drell-Yan Scattering (DY)
    - E866

$\chi_{\text{red}}^2$ 


reaction	$\chi_{\text{red}}^2$	$N_{\text{dat}}$
DIS	1.29	2680
DY	1.52	250
SIDIS $\pi^\pm$	1.39	498
$K^\pm$	1.38	494
$h^\pm$	0.85	498
SIA $\pi^\pm$	1.09	231
$K^\pm$	1.37	213
$h^\pm$	1.15	120
<b>total</b>	1.26	4984

$$\chi_{\text{red}}^2 = \frac{1}{N} \sum_{i,e} \frac{1}{\alpha_{i,e}^2} \left( d_{i,e} - \text{E} \left[ \sum_k r_e^k \beta_{i,e}^k + T_{i,e}/N_e \right] \right)^2$$


$$\text{residual}(e, i) = \frac{1}{\alpha_{i,e}} \left( d_{i,e} - \text{E} \left[ \sum_k r_e^k \beta_{i,e}^k + T_{i,e}/N_e \right] \right)$$




# SIDIS Data and theory comparison: pions




# SIDIS Data and theory comparison: kaons




# SIDIS Data and theory comparison: hadrons




# SIA Data over theory comparison: pions




# SIA Data over theory comparison: kaons




# SIA Data over theory comparison: hadrons



# PDFs




# FFs




## Strange PDF suppression

- Best fits to kaon SIA data favor smaller strange PDFs
- Consistent with JAM19 findings



# Transverse momentum dependent SIDIS predictions

- FO predictions for large transverse momentum SIDIS
- Discrepancy between FO predictions and data can be reduced significantly by including the transverse momentum dependent data in the fit.



$0.24 < z < 0.3$   
 $0.3 < z < 0.4$   
 $0.4 < z < 0.5$   
 $0.65 < z < 0.7$

# Conclusions

- Successfully performed simultaneous global extraction of PDFs and pion, kaon, and hadron FFs.
- Observed similar strange PDF suppression as was observed in JAM19.
- Comparison of FO predictions using these results to COMPASS transverse momentum dependent data shows there is still a large discrepancy
  - This discrepancy is reduced when transverse momentum dependent data is included in the fit.