

Holographic Model in Anisotropic Hot Dense QGP with External Magnetic Field

K.A. Rannu

Peoples Friendship University of Russia (RUDN)
Steklov Mathematical Institute (MI RAS)

19th International Conference
on Hadron Spectroscopy and Structure



Mexico City
28.07.2021




with I.Ya. Aref'eva and P. Slepov
JHEP 07(2021)161 [arXiv:2011.07023]

The Expected QCD Phase Diagram

Goal of Holographic QCD — describe QCD phase diagram


Requirements:

- reproduce the QCD results from perturbative theory at short distances
- reproduce Lattice QCD results at large distances (~ 1 fm) and **small μ_B**




QCD Phase Diagram: Lattice

Phase diagram
on quark mass



Main problem with $\mu \neq 0$
Imaginary chemical potential method




Columbia plot
Brown et al., PRL (1990)


Philipsen, Pinke, PRD (2016)

“Heavy” and “Light” Quarks from Columbia Plot

Light quarks



Heavy quarks



Anisotropy in QGP




- Origins of anisotropy

- Longitudinal and 2 transversal directions

- Multiplicity dependencies *ALICE*
 - *Aref'eva, Golubtsova, JHEP (2014)*

$$\mathcal{M}(s) \sim s^{0.155(4)}$$
$$\mathcal{M}(s) \sim s^{1/(\nu+2)} \Rightarrow \nu = 4.5$$

- Full anisotropy (in strong magnetic field, $eB \sim 5 - 10 m_\pi^2$, m_π – pion mass)



Peripheral HIC

Fully Anisotropic Background

$$\mathcal{L} = R - \frac{f_1(\phi)}{4} \textcolor{red}{F}_{(1)}^2 - \frac{f_2(\phi)}{4} F_{(2)}^2 - \frac{f_B(\phi)}{4} \textcolor{blue}{F}_{(B)}^2 - \frac{1}{2} \partial_\mu \phi \partial^\mu \phi - V(\phi)$$

$$A_\mu^{(1)} = A_t(z) \delta_\mu^0 \quad F_{\mu\nu}^{(2)} = q \ dy^1 \wedge dy^2 \quad F_{\mu\nu}^{(B)} = q_B \ dx \wedge dy^1$$

$$A_t(0) = \textcolor{red}{\mu} \quad g(0) = 1 \quad \textcolor{blue}{Dudal et al., (2019)}$$

$$A_t(z_h) = 0 \quad g(z_h) = 0 \quad \phi(\textcolor{violet}{z}_0) = 0 \rightarrow \sigma_{\text{string}}$$

$$ds^2 = \frac{L^2}{z^2} \textcolor{magenta}{b}(z) \left[- \textcolor{red}{g}(z) dt^2 + dx^2 + \left(\frac{z}{L} \right)^{2-\frac{2}{\nu}} dy_1^2 + e^{c_B z^2} \left(\frac{z}{L} \right)^{2-\frac{2}{\nu}} dy_2^2 + \frac{dz^2}{\textcolor{red}{g}(z)} \right]$$

I.A., A.G. (2014), Giataganas (2013) Gürsoy, Järvinen et al., (2019)

$\mathfrak{b}(z) = e^{2\mathcal{A}(z)} \rightarrow$ quarks mass “Bottom-up approach”

$\mathcal{A}(z) = -cz^2/4 \rightarrow$ heavy quarks background (**b, t**) *Andreev, Zakharov (2006)*
 $\mathcal{A}(z) = -a \ln(bz^2 + 1) \rightarrow$ light quarks background (**d, u**) *Li, Yang, Yuan (2020)*

Anisotropic Solution for “Heavy” Quarks

$$A_t = \mu \frac{e^{(c-2c_B)z^2/4} - e^{(c-2c_B)z_h^2/4}}{1 - e^{(c-2c_B)z_h^2/4}}, \quad \rho = -\frac{\mu(c-2c_B)}{4 \left(1 - e^{(c-2c_B)z_h^2/4}\right)}$$

$$f_B = -\frac{2c_B z^{1-\nu/2} g}{q_B^2 e^{cz^2/2}} \left(\frac{3cz}{2} + \frac{2}{\nu z} - c_B z - \frac{g'}{g} \right)$$

$$\begin{aligned} g = e^{c_B z^2} & \left\{ 1 - \frac{\Gamma\left(1 + \frac{1}{\nu}\right) - \Gamma\left(1 + \frac{1}{\nu}; \frac{3}{4}(2c_B - c)z^2\right)}{\Gamma\left(1 + \frac{1}{\nu}\right) - \Gamma\left(1 + \frac{1}{\nu}; \frac{3}{4}(2c_B - c)z_h^2\right)} - \right. \\ & - \frac{\mu^2 (2c_B - c)^{-\frac{1}{\nu}}}{4 \left(1 - e^{(c-2c_B)\frac{z_h^2}{4}}\right)^2} \left(\Gamma\left(1 + \frac{1}{\nu}\right) - \Gamma\left(1 + \frac{1}{\nu}; \frac{3}{4}(2c_B - c)z^2\right) \right) \times \\ & \times \left[1 - \frac{\Gamma\left(1 + \frac{1}{\nu}\right) - \Gamma\left(1 + \frac{1}{\nu}; \frac{3}{4}(2c_B - c)z^2\right)}{\Gamma\left(1 + \frac{1}{\nu}\right) - \Gamma\left(1 + \frac{1}{\nu}; \frac{3}{4}(2c_B - c)z_h^2\right)} \frac{\Gamma\left(1 + \frac{1}{\nu}\right) - \Gamma\left(1 + \frac{1}{\nu}; (2c_B - c)z_h^2\right)}{\Gamma\left(1 + \frac{1}{\nu}\right) - \Gamma\left(1 + \frac{1}{\nu}; (2c_B - c)z^2\right)} \right] \end{aligned}$$

$$\phi = \int_{z_0}^z \frac{1}{\nu \xi} \sqrt{4\nu - 4 + (4\nu c_B + 3(3c - 2c_B)\nu^2) \xi^2 + \left(\frac{3}{2} \nu^2 c^2 - 2c_B^2\right) \xi^4} d\xi, \quad z_0 \neq 0$$

Thermodynamics in Holography

$$T = \frac{|g'|}{4\pi} \Big|_{z=z_h} \longrightarrow \text{QGP temperature (Maldacena)}$$



$$s = \frac{1}{4} \sqrt{\prod_{i=1}^3 g_{ii}} \Big|_{z=z_h} \longrightarrow \text{multiplicity of process (Landau)}$$

$$F = \int_{z_h}^{\infty} s \, dT \longrightarrow \text{background BH-BH phase transition}$$


$$A_t = \mu - \rho z^2 + \dots \longrightarrow \text{chemical potential, density}$$

BH-BH Phase Transition

BH-BH describes transition from small black holes \rightarrow large black holes





BH-BH Phase Transition: “Heavy” Quarks in Magnetic Field



Light quarks *Aref'eva, K.R., Slepov, JHEP 06 (2021) 090 [arXiv:2009.05562]*

Temporal Wilson Loop

$$W[C_\vartheta] = e^{-S_{\vartheta,t}} \quad \vec{n}: \quad n_x = \cos \vartheta, \quad n_{y_1} = \sin \vartheta, \quad n_{y_2} = 0$$



Two special cases:

- $\vartheta = 0$ WL (longitudinal)
- $\vartheta = \pi/2$ WT (transversal)

$$\ell \rightarrow \infty \quad S \sim \sigma_{DW} \ell$$



the string tension



$$\sigma_{DW} = \frac{b(z) e^{\sqrt{\frac{2}{3}} \phi(z, z_0)}}{z^2} \sqrt{g(z) \left(z^{2-\frac{2}{\nu}} \sin^2(\vartheta) + \cos^2(\vartheta) \right)} \Big|_{z=z_{DW}}, \quad \left. \frac{\partial \sigma}{\partial z} \right|_{z=z_{DW}} = 0$$

Aref'eva, K.R., Slepov, PLB 792 (2019) 470 [arXiv:1808.05596]


Phase Transition for “Heavy” Quarks

JHEP 05(2018)206
[arXiv:1802.05652]


$$c_B = 0$$



$$c_B = -0.0096$$




$$c_B = -0.015$$



- We see the **inverse** magnetic catalysis

Our String Tension vs Lattice

- We take $z_0 = 10 \exp(-z_h/4) + 0.1$ (#)



- The green curve shows the string tension as function of temperature for $z_0 = (\#)$, $\mu = 0$, $\nu = 1$, $a = 4.046$, $b = 0.01613$, $c = 0.227$
- The dots with different decorations Lattice *Cardoso, Bicudo, PRD (2012)*
- The thin green dotted line shows the WL phase transition

Spatial Wilson Loops



Nesting of the 2-dimensional world sheet in 5-dimensional space-time

$$X^0(\xi) = \text{const}$$

$$X^i(\xi) = \sum_{\alpha=1,2} a_{i\alpha}(\phi, \theta, \psi) \xi^\alpha, \quad i = 1, 2, 3, \quad \alpha = 1, 2$$



$$X^4(\xi) = z(\xi^1)$$

- x^i – spatial coordinates
- $a_{ij}(\phi, \theta, \psi)$ – entries of the rotation matrix



*brick's orientation is equivalent
to rectangle's orientation*

Drag Forces Phase Transitions



Phase transition surfaces – surfaces stretched on blue/green curves

- Under variation of (T , μ , magnetic field) the spatial string tension undergoes the phase transition *Aref'eva, K.R., Slepov [arXiv:2012.05758]*
- The light green and blue lines represent to the end of the phase transitions
- Drag forces and energy losses undergo the phase transition as well

Conclusion

- We have an interplay between two types of phase transitions
 - Crossover
 - 1-st order phase transition
- Magnetic field influences the phase diagram for “heavy” quarks model
 - Inverse magnetic catalysis
 - External magnetic field suppresses BH-BH transition
 - For larger anisotropy $\nu > 1$ this suppression weakens
 - Drag forces and energy losses undergo the phase transition depending on anisotropy and magnetic field
- What to do next
 - “Light” quarks model in magnetic field
 - “Hybrid” model corresponding to realistic quark masses
 - Other characteristics (susceptibility, transport coefficients, eta/s, direct-photon spectra, jet quenching, thermalization time, etc)

Thank you
for your attention