

# ATLAS results on charmonium production and $B_c^+$ production and decays

Tatiana Lyubushkina<sup>1</sup>  
*on behalf of ATLAS Collaboration*

<sup>1</sup>Joint Institute for Nuclear Research  
*Tatiana.Lyubushkina@cern.ch*




19th International Conference on Hadron Spectroscopy and Structure  
26-31 July 2021

# Introduction



## B-physics at ATLAS

- ▶ Possible to measure heavy flavour (HF) production at *high energy*
- ▶ Much *higher HF yields* in *pp* environment compared to B-factories
- ▶ Extended spectroscopy studies are possible
- ▶ ATLAS and CMS, although not specially optimized for B-physics, provide *complementary kinematic region to LHCb*
  - ▶ Benefit from *higher statistics* in certain analyses
- ▶ This talk covers a selection of ATLAS results
  - ▶ High- $p_T$   $J/\psi$  and  $\psi(2S)$  production at 13 TeV [ATLAS-CONF-2019-047](#)
  - ▶ Relative  $B_c/B^+$  production measurement at 8 TeV – [Phys. Rev. D 104, 012010](#)
  - ▶ Study of the  $B_c^+ \rightarrow J/\psi D_s^+$  and  $B_c^+ \rightarrow J/\psi D_s^{*+}$  decays in  $pp$  collisions at  $\sqrt{s} = 13$  TeV with the ATLAS detector [ATLAS-CONF-2021-046](#)






# High- $p_T$ $J/\psi$ and $\psi(2S)$ production measurement

- ▶ Two different mechanisms for charmonium production:
  - ▶ *Prompt*: directly in  $pp$  interaction or via feed-down from heavier states
  - ▶ *Non-prompt*: from decays of  $b$  hadrons
    - ▶ Can be distinguished by fitting the pseudo proper lifetime
  - ▶ FONLL calculations within the framework of perturbative QCD have been reasonably successful in describing the non-prompt contributions
  - ▶ Satisfactory understanding of the prompt production mechanisms is still to be achieved
- ▶ Previous ATLAS measurement:  $J/\psi$  and  $\psi(2S)$  differential x-sections at 7, 8 TeV – *Eur. Phys. J. C 76 (2016) 283*
  - ▶ Overall reasonable agreement with theoretical predictions for prompt and non-prompt production
  - ▶ Di-muon triggers with low thresholds – could not reach beyond  $p_T$  of  $\sim 100$  GeV
- ▶ New measurement focuses on high- $p_T$  charmonia
  - ▶ Help to discriminate various theoretical models
  - ▶ Use *single-muon* triggers (50 GeV muon  $p_T$  threshold) to cover the range **60–360 GeV**
  - ▶ Full Run-2 dataset,  $139 \text{ fb}^{-1}$  at 13 TeV



# High- $p_T$ $J/\psi$ and $\psi(2S)$ measurement

- Measured are
  - Prompt and non-prompt  $J/\psi$  and  $\psi(2S)$  double-differential x-sections
  - Non-prompt fraction for  $J/\psi$  and  $\psi(2S)$
  - $\psi(2S)/J/\psi$  production ratio for prompt and non-prompt



- $p_T$  ranges extend significantly
- FONLL consistent at low- $p_T$ , over-estimates high- $p_T$  production

# Relative $B_c^+ / B^+$ production measurement at 8 TeV

- ▶  $B_c^+$  is the only known weakly decaying particle made of two heavy quarks

## ▶ Motivation:





- ▶ test of the QCD prediction;
- ▶ important input for heavy quark production models;
- ▶ complements CMS and LHCb measurements;

- ▶ Measure the ratio: 
$$\frac{\sigma(B_c^+) \cdot \mathcal{B}(B_c^+ \rightarrow J/\psi \pi^+) \cdot \mathcal{B}(J/\psi \rightarrow \mu^+ \mu^-)}{\sigma(B^+) \cdot \mathcal{B}(B^+ \rightarrow J/\psi K^+) \cdot \mathcal{B}(J/\psi \rightarrow \mu^+ \mu^-)}$$

- ▶ Common systematic uncertainties mostly cancel

- ▶ Fiducial region of the measurement:

- ▶  $p_T(B) > 13 \text{ GeV}$ ,  $|y(B)| < 2.3$
- ▶ In addition to the full bin two bins in  $p_T$  ( $13 < p_T(B) < 22 \text{ GeV}$  and  $p_T > 22 \text{ GeV}$ ) and two bins in rapidity ( $|y| < 0.75$  and  $0.75 < |y| < 2.3$ ) were defined



# $B_c^+ / B^+$ production: results



- Production ratio in the fiducial region

$$\frac{\sigma(B_c^+) \cdot \mathcal{B}(B_c^+ \rightarrow J/\psi\pi^+) \cdot \mathcal{B}(J/\psi \rightarrow \mu^+\mu^-)}{\sigma(B^+) \cdot \mathcal{B}(B^+ \rightarrow J/\psi K^+) \cdot \mathcal{B}(J/\psi \rightarrow \mu^+\mu^-)} =$$

$$(0.34 \pm 0.04(\text{stat.})_{-0.02}^{+0.06}(\text{syst.}) \pm 0.01(\text{lifetime}))\%$$


- Lower than the **LHCb result** for more forward and lower- $p_T$  fiducial volume:  $(0.683 \pm 0.018 \pm 0.009)\%$ ,  $0 < p_T(B) < 20 \text{ GeV}$ ,  $2 < y(B) < 4.5$
- Fairly consistent with the **CMS result** in a similar (but not identical) fiducial volume:  $(0.48 \pm 0.05(\text{stat.}) \pm 0.03(\text{syst.}) \pm 0.05(\text{lifetime}))\%$  at  $\sqrt{s} = 7 \text{ TeV}$ ,  $p_T(B) > 15 \text{ GeV}$ ,  $|y(B)| < 1.6$
- $B_c^+$  production decreases faster with  $p_T$  than that for  $B^+$
- No evident rapidity dependence

Analysis bin	$\sigma(B_c^\pm)/\sigma(B^\pm) \times \mathcal{B}(B_c^\pm \rightarrow J/\psi\pi^\pm)/\mathcal{B}(B^\pm \rightarrow J/\psi K^\pm)$
$p_T(B) > 13 \text{ GeV}$ , $ y(B)  < 2.3$	$(0.34 \pm 0.04_{\text{stat}}^{+0.06}_{-0.02} \text{syst} \pm 0.01_{\text{lifetime}})\%$
$13 < p_T(B) < 22 \text{ GeV}$ , $ y(B)  < 2.3$	$(0.44 \pm 0.07_{\text{stat}}^{+0.09}_{-0.04} \text{syst} \pm 0.01_{\text{lifetime}})\%$
$p_T(B) > 22 \text{ GeV}$ , $ y(B)  < 2.3$	$(0.24 \pm 0.04_{\text{stat}}^{+0.05}_{-0.01} \text{syst} \pm 0.01_{\text{lifetime}})\%$
$p_T(B) > 13 \text{ GeV}$ , $ y(B)  < 0.75$	$(0.38 \pm 0.06_{\text{stat}}^{+0.05}_{-0.04} \text{syst} \pm 0.01_{\text{lifetime}})\%$
$p_T(B) > 13 \text{ GeV}$ , $0.75 <  y(B)  < 2.3$	$(0.29 \pm 0.05_{\text{stat}}^{+0.07}_{-0.02} \text{syst} \pm 0.01_{\text{lifetime}})\%$



## New: Study of the $B_c^+ \rightarrow J/\psi D_s^{(*)+}$


- ▶ Decays  $B_c^+ \rightarrow J/\psi D_s^{(*)+}$  can occur through  $b$  decay with  $c$  as spectator, or through annihilation diagram



- ▶ Only  $B_c^+ \rightarrow J/\psi D_s^{(*)+}$  decays observed earlier by LHCb ([PRD 87 \(2013\) 112012](#)) and ATLAS ([EPJC 76 \(2016\) 4](#)).
- ▶ This analysis aims at more precise measurement of  $B_c^+ \rightarrow J/\psi D_s^{(*)+}$  branching fraction and polarization with full Run-2 data
  - ▶ Test various approaches predicting these ([perturbative QCD calculation](#), [relativistic potential models](#), [sum rules calculations](#)...)

# Study of the $B_c^+ \rightarrow J/\psi D_s^{(*)+}$ decays

- ▶  $B_c^+ \rightarrow J/\psi(\mu^+\mu^-)D_s^+(\rightarrow \phi(\rightarrow K^+K^-)\pi^+)$
- ▶  $B_c^+ \rightarrow J/\psi(\mu^+\mu^-)D_s^{*+}(\rightarrow D_s^+\gamma/\pi^0)$ 
  - ▶ Same reconstructed final state, soft neutral particle escapes detection
- ▶ **Reference channel:**  $B_c^+ \rightarrow J/\psi\pi^+$ 
  - ▶ Use it for  $\mathcal{B}$  measurement
- ▶ Define fiducial range of the measurement:  
 $p_T(B_c^+) > 15 \text{ GeV}, |\eta(B_c^+)| < 2.0$
- ▶ Measured quantities:
  - ▶ Ratios b/w  $\mathcal{B}$  of signal channels and  $\mathcal{B}(B_c^+ \rightarrow J/\psi\pi^+)$ :  $R_{D_s^{(*)+}/\pi^+}$
  - ▶ Ratios b/w  $\mathcal{B}$ 's of signal channels (to cancel some of the uncertainties):  $R_{D_s^{*+}/D_s^+}$
  - ▶ Transverse polarisation fraction  $\Gamma_{\pm\pm}/\Gamma$  for  $B_c^+ \rightarrow J/\psi D_s^{*+}$



# Study of the $B_c^+ \rightarrow J/\psi D_s^{(*)+}$ decays





- ▶ *Dataset 1*: candidates in the events collected by the standard dimuon or three-muon triggers without requirements on additional ID track.

- ▶ can be safely used to measure

$$R_{D_s^+/\pi^+}, R_{D_s^{*+}/\pi^+}$$

- ▶ *Dataset 2*: candidates collected only by the dedicated  $B_s^0 \rightarrow \mu^+ \mu^- \phi$  triggers and not by other ones used in the analysis.

- ▶ improve sensitivity to  $R_{D_s^{*+}/D_s^+}$ ,  $\Gamma_{\pm\pm}/\Gamma$




# Study of the $B_c^+ \rightarrow J/\psi D_s^{(*)+}$ decays: results

$$R_{D_s^+/\pi^+} = 2.76 \pm 0.33(\text{stat.}) \pm 0.29(\text{syst.}) \pm 0.16(\text{br.f.})$$

$$R_{D_s^{*+}/\pi^+} = 5.33 \pm 0.61(\text{stat.}) \pm 0.67(\text{syst.}) \pm 0.32(\text{br.f.})$$

$$R_{D_s^{*+}/D_s^+} = 1.93 \pm 0.24(\text{stat.}) \pm 0.10(\text{syst.})$$

$$\Gamma_{\pm\pm}/\Gamma = 0.70 \pm 0.10(\text{stat.}) \pm 0.04(\text{syst.})$$



- ▶ All results are consistent with the earlier measurements of ATLAS and LHCb.
- ▶ The precision of the measurement exceeds that of all previous studies of these decays




$R_{D_s^+/\pi^+}$	$R_{D_s^{*+}/\pi^+}$	$R_{D_s^{*+}/D_s^+}$	$\Gamma_{\pm\pm}/\Gamma$	Ref.
$2.76 \pm 0.47$	$5.33 \pm 0.96$	$1.93 \pm 0.26$	$0.70 \pm 0.11$	ATLAS Run 2
$2.90 \pm 0.62$	–	$2.37 \pm 0.57$	$0.52 \pm 0.20$	LHCb Run 1
$3.8 \pm 1.2$	$10.4 \pm 3.5$	$2.8^{+1.2}_{-0.9}$	$0.38 \pm 0.24$	ATLAS Run 1
2.6	4.5	1.7	–	QCD potential model
1.3	5.2	3.9	–	QCD sum rules
2.0	5.7	2.9	–	RCQM
$1.29 \pm 0.26$	$5.09 \pm 1.02$	$3.96 \pm 0.80$	$0.46 \pm 0.09$	CCQM
2.2	–	–	–	BSW
$2.06 \pm 0.86$	–	$3.01 \pm 1.23$	–	LFQM
$3.45^{+0.49}_{-0.17}$	–	$2.54^{+0.07}_{-0.21}$	$0.48 \pm 0.04$	pQCD
–	–	–	0.410	RIQM
$3.257 \pm 0.293$	–	–	–	FNCM

# Summary

- ▶ A selection of ATLAS results on heavy flavour production was presented
  - ▶ Production of hidden charm
  - ▶ Physics of  $B_c^+$  mesons
- ▶ More can be found here:
  - ▶ [ATLAS B-physics public results page](#) 
- ▶ Many interesting results are still to come!

Backup slides



# ATLAS detector and trigger



# $J/\psi$ and $\psi(2S)$ fits



i	Type	P/NP	$f_i(m)$	$h_i(\tau)$
1	$J/\psi$	P	$\omega G_1(m) + (1 - \omega)CB_1(m)$	$\delta(\tau)$
2	$J/\psi$	NP	$\omega G_1(m) + (1 - \omega)CB_1(m)$	$E_1(\tau)$
3	$\psi(2S)$	P	$\omega G_2(m) + (1 - \omega)CB_2(m)$	$\delta(\tau)$
4	$\psi(2S)$	NP	$\omega G_2(m) + (1 - \omega)CB_2(m)$	$E_2(\tau)$
5	Bkg	P	$B$	$\delta(\tau)$
6	Bkg	NP	$E_4(m)$	$E_5(\tau)$
7	Bkg	NP	$E_6(m)$	$E_7( \tau )$

Notation	Function
$G$	Gaussian
$CB$	Crystal Ball
$E$	Exponential
$B$	Bernstein polynomials




# High- $p_T$ $J/\psi$ and $\psi(2S)$ measurement

- ▶ Measured are
  - ▶ Prompt and non-prompt  $J/\psi$  and  $\psi(2S)$  double-differential x-sections
  - ▶ Non-prompt fraction for  $J/\psi$  and  $\psi(2S)$
  - ▶  $\psi(2S)/J/\psi$  production ratio for prompt and non-prompt
- ▶  $p_T$  ranges extend significantly
- ▶ FONLL consistent at low- $p_T$ , over-estimates high- $p_T$  production



# $B_c^+ \rightarrow J/\psi\pi^+$ fit

Parameter	Value
$m_{B_c^+}$ [MeV]	$6274.5 \pm 1.5$
$\sigma_{B_c^+}$ [MeV]	$47.5 \pm 2.5$
$N_{B_c^+ \rightarrow J/\psi\pi^+}$	$8440^{+550}_{-470}$







## $B_c^+ \rightarrow J/\psi D_s^{(*)+}$ fit PDF

- ▶ 2D unbinned ML fit of  $m(J/\psi D_s^+)$  and  $|\cos\theta'(\mu^+)|$ ; mass and angular PDF are factorized
- ▶ ratio between  $B_c^+ \rightarrow J/\psi D_s^+$  and  $B_c^+ \rightarrow J/\psi D_s^{*+}$  yield and  $f_{\pm\pm}$  are the same in DS1 and DS2
- ▶  $B_c^+ \rightarrow J/\psi D_s^+$  signal
  - ▶ *mass*: modified Gaussian<sup>1</sup>
  - ▶  $|\cos\theta'(\mu^+)|$ : MC kernel template (same in DS1 and DS2)
- ▶  $B_c^+ \rightarrow J/\psi D_s^{*+}$  signals, separately  $A_{\pm\pm}$  and  $A_{00}$  components
  - ▶ *mass*: MC kernel templates (same in DS1 and DS2)
  - ▶  $|\cos\theta'(\mu^+)|$ : MC kernel templates (same in DS1 and DS2)
- ▶ Background
  - ▶ *mass*: exponential (same slope in DS1 and DS2)
  - ▶  $|\cos\theta'(\mu^+)|$ : 2nd order polynomial (same parameters in DS1 and DS2)

---

<sup>1</sup> $Gauss^{mod} \propto \exp(-0.5 \times t^{1+1/(1+t/2)})$ , where  $t = |m(J/\psi D_s^+) - m_{B_c^+}|/\sigma_{B_c^+}$

# $B_c^+ \rightarrow J/\psi D_s^{(*)+}$ fit result



Parameter	Value
$m_{B_c^+}$ [MeV]	$6274.8 \pm 1.4$
$\sigma_{B_c^+}$ [MeV]	$11.5 \pm 1.5$
$r_{D_s^{*+}/D_s^+}$	$1.76 \pm 0.22$
$f_{\pm\pm}$	$0.70 \pm 0.10$
$N_{B_c^+ \rightarrow J/\psi D_s^+}^{\text{DS1}}$	$193 \pm 20$
$N_{B_c^+ \rightarrow J/\psi D_s^+}^{\text{DS2}}$	$49 \pm 10$
$N_{B_c^+ \rightarrow J/\psi D_s^{*+}}^{\text{DS1}}$	$338 \pm 32$
$N_{B_c^+ \rightarrow J/\psi D_s^{*+}}^{\text{DS1\&2}}$	$241 \pm 28$
$N_{B_c^+ \rightarrow J/\psi D_s^{*+}}^{\text{DS1\&2}}$	$424 \pm 46$

# Ratios calculation

$$R_{D_s^{(*)+}/\pi^+} = \frac{\mathcal{B}(B_c^+ \rightarrow J/\psi D_s^{(*)+})}{\mathcal{B}(B_c^+ \rightarrow J/\psi \pi^+)} = \frac{N_{B_c^+ \rightarrow J/\psi D_s^{(*)+}}^{\text{DS1}}}{N_{B_c^+ \rightarrow J/\psi \pi^+}} \times \frac{\varepsilon_{B_c^+ \rightarrow J/\psi \pi^+}}{\varepsilon_{B_c^+ \rightarrow J/\psi D_s^{(*)+}}^{\text{DS1}}} \times \frac{1}{\mathcal{B}(D_s^+ \rightarrow \phi(K^+K^-)\pi^+)}, \quad (1)$$

▶  $\mathcal{B}(D_s^+ \rightarrow \phi(K^+K^-)\pi^+)$  taken as  $m(K^+K^-)$ -dependent, using CLEO measurement, recalculated to  $\pm 7$  MeV

$$R_{D_s^{*+}/D_s^+} = \frac{\mathcal{B}(B_c^+ \rightarrow J/\psi D_s^{*+})}{\mathcal{B}(B_c^+ \rightarrow J/\psi D_s^+)} = \frac{N_{B_c^+ \rightarrow J/\psi D_s^{*+}}^{\text{DS1\&2}}}{N_{B_c^+ \rightarrow J/\psi D_s^+}^{\text{DS1\&2}}} \times \frac{\varepsilon_{B_c^+ \rightarrow J/\psi D_s^+}^{\text{DS1\&2}}}{\varepsilon_{B_c^+ \rightarrow J/\psi D_s^{*+}}^{\text{DS1\&2}}} = r_{D_s^{*+}/D_s^+} \times \frac{\varepsilon_{B_c^+ \rightarrow J/\psi D_s^+}^{\text{DS1\&2}}}{\varepsilon_{B_c^+ \rightarrow J/\psi D_s^{*+}}^{\text{DS1\&2}}}, \quad (2)$$

$$\varepsilon_{B_c^+ \rightarrow J/\psi D_s^{*+}} = \frac{1}{f_{\pm\pm}/\varepsilon_{B_c^+ \rightarrow J/\psi D_s^{*+}, A_{\pm\pm}} + (1 - f_{\pm\pm})/\varepsilon_{B_c^+ \rightarrow J/\psi D_s^{*+}, A_{00}}}, \quad (3)$$

$$\Gamma_{\pm\pm}/\Gamma = f_{\pm\pm} \times \frac{\varepsilon_{B_c^+ \rightarrow J/\psi D_s^{*+}}^{\text{DS1\&2}}}{\varepsilon_{B_c^+ \rightarrow J/\psi D_s^{*+}, A_{\pm\pm}}^{\text{DS1\&2}}}. \quad (4)$$

Mode	$\varepsilon_{B_c^+ \rightarrow J/\psi X}^{\text{DS1}}$ [%]	$\varepsilon_{B_c^+ \rightarrow J/\psi X}^{\text{DS1\&2}}$ [%]
$B_c^+ \rightarrow J/\psi D_s^+$	$0.971 \pm 0.012$	$1.163 \pm 0.013$
$B_c^+ \rightarrow J/\psi D_s^{*+}, A_{00}$	$0.916 \pm 0.012$	$1.088 \pm 0.012$
$B_c^+ \rightarrow J/\psi D_s^{*+}, A_{\pm\pm}$	$0.868 \pm 0.010$	$1.049 \pm 0.011$
$B_c^+ \rightarrow J/\psi \pi^+$	$2.169 \pm 0.018$	–

Source	Uncertainty [%]			
	$R_{D_s^+/\pi^+}$	$R_{D_s^{*+}/\pi^+}$	$R_{D_s^{*+}/D_s^+}$	$\Gamma_{\pm\pm}/\Gamma$
Simulated $p_T(B_c^+)$ spectrum	1.5	1.9	0.4	0.1
Simulated $ \eta(B_c^+) $ spectrum	0.7	0.7	0.1	0.2
$B_c^+$ lifetime	0.1	< 0.1	-	-
$D_s^+$ lifetime	0.4	0.4	-	-
Tracking efficiency	1.0	1.0	< 0.1	< 0.1
Pile-up effects	1.0	1.0	-	-
$\chi^2/ndf$ cut efficiency	3.2	3.2	-	-
Impact parameter cuts efficiency	0.2	0.2	-	-
BDT cut efficiency	1.3	1.3	-	-
Trigger efficiency	1.0	1.0	-	-
$B_c^+ \rightarrow J/\psi D_s^{(*)+}$ signal fit:				
$D_s^+$ signal mass modelling	1.8	0.5	1.3	0.8
$D_s^{*+}$ signal mass modelling	0.6	1.2	1.7	2.7
signal angular modelling	0.4	< 0.1	0.4	0.6
background mass modelling	6.0	9.0	3.2	1.0
background angular modelling	0.9	1.3	2.1	2.4
$B_s^0 \rightarrow \mu^+ \mu^- \phi$ triggers	0.8	0.5	1.3	4.0
$B_c^+ \rightarrow J/\psi \pi^+$ signal fit				
signal modelling	4.2	4.2	-	-
PRD/comb. background modelling	5.8	5.8	-	-
CKM-suppr. background modelling	1.0	1.0	-	-
MC statistics	1.5	1.5	1.7	1.5
Total	10.7	12.6	5.0	5.8
$B(D_s^+ \rightarrow \phi(K^+ K^-)\pi^+)$	5.9	5.9	-	-

- ▶ Effect of muon reconstruction and identification efficiency uncertainty affects individual channel efficiencies by about 1–2%. However, for the measured quantities, due to cancellation in the efficiency ratios, it is found to be negligible.