

Novel QCD Phenomena at the LHC (II)



Stan Brodsky
SLAC

October 1, 2010

HIGH ENERGY
Instituto de Ciencias Nucleares
MÉXICO, CITY
SEPTEMBER 27th- OCTOBER 1st
UNIVERSIDAD NACIONAL AUTÓNOMA DE MÉXICO




5th Workshop on High p-T
Physics at LHC




QCD Myths

- **Anti-Shadowing is Universal**
- **ISI and FSI are higher twist effects and universal**
- **High transverse momentum hadrons arise only from jet fragmentation -- baryon anomaly!**
- **heavy quarks only from gluon splitting**
- **renormalization scale cannot be fixed**
- **QCD condensates are vacuum effects**
- **Infrared Slavery**
- **Nuclei are composites of nucleons only**
- **Real part of DVCS arbitrary**

Deep Inelastic Electron-Proton Scattering



Deep Inelastic Electron-Proton Scattering



*Conventional wisdom:
Final-state interactions of struck quark can be neglected*

Single-spin asymmetries

**Leading Twist
Sivers Effect**


Hwang,
Schmidt, sjb

Collins, Burkardt
Ji, Yuan

*QCD S- and P-
Coulomb Phases
--Wilson Line*

*Leading-Twist
Rescattering
Violates pQCD
Factorization!*

$i \vec{S}_p \cdot \vec{q} \times \vec{p}_q$
Pseudo-T-Odd




*Light-Front Wavefunction
S and P-Waves*

Final-State Interactions Produce Pseudo T-Odd (Sivers Effect)


Hwang, Schmidt, sjb
Collins

- **Leading-Twist Bjorken Scaling!**
- **Requires nonzero orbital angular momentum of quark**
- **Arises from the interference of Final-State QCD Coulomb phases in S- and P-waves;**
- **Wilson line effect -- gauge independent**
- **Relate to the quark contribution to the target proton anomalous magnetic moment and final-state QCD phases**
- **QCD phase at soft scale!**
- **New window to QCD coupling and running gluon mass in the IR**
- **QED S and P Coulomb phases infinite -- difference of phases finite!**
- **Alternate: Retarded and Advanced Gauge: Augmented LFWFs**


$$\mathbf{i} \vec{S} \cdot \vec{p}_{jet} \times \vec{q}$$



Pasquini, Xiao, Yuan, sjb
Mulders, Boer Qiu, Sterman




can interfere with



and produce a T-odd effect!
(also need $L_z \neq 0$)

HERMES coll., A. Airapetian et al., Phys. Rev. Lett. 94 (2005) 012002.

Sivers asymmetry from HERMES



- First evidence for non-zero Sivers function!
- \Rightarrow presence of non-zero **quark orbital angular momentum!**
- **Positive** for π^+ ...
Consistent with zero for π^- ...


Gamberg: Hermes data compatible with BHS model

Schmidt, Lu: Hermes charge pattern follow quark contributions to anomalous moment

Anomalous effect from Double ISI in Massive Lepton Production


Boer, Hwang, sjb

$\cos 2\phi$ correlation



- Leading Twist, valence quark dominated
- Violates Lam-Tung Relation!
- Not obtained from standard PQCD subprocess analysis
- Normalized to the square of the single spin asymmetry in semi-inclusive DIS
- No polarization required
- Challenge to standard picture of PQCD Factorization

Predict Opposite Sign SSA in DY !



Collins;
Hwang,
Schmidt. sjb

Single Spin Asymmetry In the Drell Yan Process

$$\vec{S}_p \cdot \vec{p} \times \vec{q}_{\gamma^*}$$

Quarks Interact in the Initial State

Interference of Coulomb Phases for S and P states

Produce Single Spin Asymmetry [Siver's Effect] Proportional
to the Proton Anomalous Moment and α_s .

Opposite Sign to DIS! No Factorization


Key QCD Experiment

Collins;
Hwang,
Schmidt. sjb

Measure single-spin asymmetry A_N
in Drell-Yan reactions

Leading-twist Bjorken-scaling A_N
from S, P -wave
initial-state gluonic interactions

Predict: $A_N(DY) = -A_N(DIS)$
Opposite in sign!



$$\bar{p} p_{\uparrow} \rightarrow l^{+} l^{-} X$$

$$\vec{S} \cdot \vec{q} \times \vec{p}$$
 correlation

$$f_1 = \text{[Diagram: Yellow circle with a light blue center dot.]}$$

*Unpolarized
Distribution*

$$g_{1L} = \text{[Diagram: Yellow circle with light blue center dot and right-pointing arrow.]} - \text{[Diagram: Yellow circle with light blue center dot and left-pointing arrow.]}$$

Bj Sum Rule

$$h_{1T} = \text{[Diagram: Yellow circle with light blue center dot and up-pointing arrow.]} - \text{[Diagram: Yellow circle with light blue center dot and down-pointing arrow.]}$$

Transversity


$$f_{1T}^\perp = \text{[Diagram: Yellow circle with light blue center dot and up-pointing arrow.]} - \text{[Diagram: Yellow circle with light blue center dot and down-pointing arrow.]}$$

Sivers Function

$$h_1^\perp = \text{[Diagram: Yellow circle with light blue center dot and down-pointing arrow.]} - \text{[Diagram: Yellow circle with light blue center dot and up-pointing arrow.]}$$

*Boer-Mulders
Function*

*T-Odd:
Require ISI or FSI*




DY $\cos 2\phi$ correlation at leading twist from double ISI

Product of Boer - Mulders Functions

$$h_1^\perp(x_1, \mathbf{p}_\perp^2) \times \bar{h}_1^\perp(x_2, \mathbf{k}_\perp^2)$$

Drell-Yan angular distribution

Unpolarized DY



Lam – Tung SR : $1 - \lambda = 2\nu$

NLO pQCD : $\lambda \approx 1 \quad \mu \approx 0 \quad \nu \approx 0$

- Experimentally, a violation of the Lam-Tung sum rule is observed by sizeable $\cos 2\Phi$ moments
- Several model explanations
 - higher twist
 - spin correlation due to non-trivial QCD vacuum
 - Non-zero Boer Mulders function

$$\frac{1}{\sigma} \frac{d\sigma}{d\Omega} = \frac{3}{4\pi} \frac{1}{\lambda + 3} \left(1 + \lambda \cos^2 \theta + \mu \sin 2\theta \cos \phi + \frac{\nu}{2} \sin^2 \theta \cos 2\phi \right)$$

Experiment: $\nu \simeq 0.6$

B. Seitz

Double Initial-State Interactions


generate anomalous $\cos 2\phi$

Boer, Hwang, sjb



Drell-Yan planar correlations

$$\frac{1}{\sigma} \frac{d\sigma}{d\Omega} \propto \left(1 + \lambda \cos^2 \theta + \mu \sin 2\theta \cos \phi + \frac{\nu}{2} \sin^2 \theta \cos 2\phi \right)$$

PQCD Factorization (Lam Tung): $1 - \lambda - 2\nu = 0$



Violates Lam-Tung relation!



Model: Boer,

Stan Brodsky, SLAC


UNAM

October 1, 2010

Novel QCD Phenomena at the LHC

LHC Experiment

Boer, Hwang, sjb




$DY \cos 2\phi$ correlation at leading twist from double ISI

Product of Boer - Mulders Functions

$$h_1^\perp(x_1, \mathbf{p}_\perp^2) \times \bar{h}_1^\perp(x_2, \mathbf{k}_\perp^2)$$

Measurement of Angular Distributions of Drell-Yan Dimuons in $p + d$ Interaction at 800 GeV/c


(FNAL E866/NuSea Collaboration)




Huge Effect in
 $\pi W \rightarrow \mu^+ \mu^- X$
 Negligible Effect
 $pd \rightarrow \mu^+ \mu^- X$

Parameter ν vs. p_T in the Collins-Soper frame for three Drell-Yan measurements. Fits to the data using Eq. 3 and $M_C = 2.4 \text{ GeV}/c^2$ are also shown.

Important Corrections from Initial and Final State Corrections



Sivers & Collins Odd-T Spin Effects, Co-planarity Correlations




Problem for factorization when both ISI and FSI occur


Factorization is violated in production of high-transverse-momentum particles in hadron-hadron collisions

John Collins, [Jian-Wei Qiu](#) . ANL-HEP-PR-07-25, May 2007.

e-Print: [arXiv:0705.2141](#) [hep-ph]




The exchange of two extra gluons, as in this graph, will tend to give non-factorization in unpolarized cross sections.



$\cos 2\phi$ correlation for quarkonium production at leading twist from double ISI

Enhanced by gluon color charge


DDIS




- In a large fraction ($\sim 10\text{--}15\%$) of DIS events, the proton escapes intact, keeping a large fraction of its initial momentum
- This leaves a large *rapidity gap* between the proton and the produced particles
- The t -channel exchange must be *color singlet* \rightarrow a *pomeron*??

Diffractive Deep Inelastic Lepton-Proton Scattering

Remarkable observation at HERA




*10% to 15%
of DIS events
are
diffractive!*



Fraction r of events with a large rapidity gap, $\eta_{\max} < 1.5$, as a function of Q_{DA}^2 for two ranges of x_{DA} . No acceptance corrections have been applied.

M. Derrick et al. [ZEUS Collaboration], Phys. Lett. B 315, 481 (1993)

Diffractive Structure Function F_2^D



Diffractive inclusive cross section


$$\frac{d^3 \sigma_{NC}^{diff}}{dx_{IP} d\beta dQ^2} \propto \frac{2\pi\alpha^2}{xQ^4} F_2^{D(3)}(x_{IP}, \beta, Q^2)$$

$$F_2^D(x_{IP}, \beta, Q^2) = f(x_{IP}) \cdot F_2^{IP}(\beta, Q^2)$$

extract DPDF and $xg(x)$ from scaling violation

Large kinematic domain $3 < Q^2 < 1600 \text{ GeV}^2$



Precise measurements sys 5%, stat 5–20%




Diffractive Deep Inelastic Scattering

Diffractive DIS $ep \rightarrow epX$ where there is a large rapidity gap and the target nucleon remains intact probes the final state interaction of the scattered quark with the spectator system via gluon exchange.

Diffractive DIS on nuclei $eA \rightarrow e'AX$ and hard diffractive reactions such as $\gamma^* A \rightarrow VA$ can occur coherently leaving the nucleus intact.



Final-State Interaction Produces Diffractive DIS



Quark Rescattering

Hoyer, Marchal, Peigne, Sannino, SJB (BHMPS)


Enberg, Hoyer, Ingelman, SJB

Hwang, Schmidt, SJB

1-2005
8711A18


Low-Nussinov model of Pomeron

QCD Mechanism for Rapidity Gaps




Reproduces lab-frame color dipole approach

Final State Interactions in QCD




Feynman Gauge



Light-Cone Gauge


Result is Gauge Independent

Predict: Reduced DDIS/DIS for Heavy Quarks



Kopeliovitch, Schmidt, sjb

Reproduces lab-frame color dipole approach




Integration over on-shell domain produces phase i

Need Imaginary Phase to Generate Pomeron and DDIS

Need Imaginary Phase to Generate
T-Odd Single-Spin Asymmetry

Physics of FSI not in Wavefunction of Target


Nuclear Shadowing in QCD




Shadowing depends on understanding leading twist-diffraction in DIS

Nuclear Shadowing not included in nuclear LFWF !


Dynamical effect due to virtual photon interacting in nucleus



M. Hirai, S. Kumano and T. H. Nagai,
 "Nuclear parton distribution functions
 and their uncertainties,"
 Phys. Rev. C **70**, 044905 (2004)
 [arXiv:hep-ph/0404093].




M. Hirai, S. Kumano and T. H. Nagai,
 "Nuclear parton distribution functions
 and their uncertainties,"
 Phys. Rev. C **70**, 044905 (2004)
 [arXiv:hep-ph/0404093].



The one-step and two-step processes in DIS on a nucleus.


Coherence at small Bjorken x_B :
 $1/Mx_B = 2\nu/Q^2 \geq L_A$.



If the scattering on nucleon N_1 is via pomeron exchange, the one-step and two-step amplitudes are opposite in phase, thus diminishing the \bar{q} flux reaching N_2 .

→ Shadowing of the DIS nuclear structure functions.

Observed HERA DDIS produces nuclear shadowing



Shadowing depends on leading-twist DDIS

Integration over on-shell domain produces phase i


Need Imaginary Phase to Generate Pomeron

Need Imaginary Phase to Generate T-Odd Single-Spin Asymmetry

Physics of FSI not in Wavefunction of Target


Antishadowing (Reggeon exchange) is not universal!

Schmidt, Yang, sjb



The one-step and two-step processes in DIS on a nucleus.

Coherence at small Bjorken x_B :
 $1/Mx_B = 2\nu/Q^2 \geq L_A$.




If the scattering on nucleon N_1 is via ~~pomeron~~ exchange, the one-step and two-step amplitudes are ~~opposite~~ in phase, thus ~~diminishing~~ the \bar{q} flux reaching N_2 .

Reggeon

increasing

→ *Anti-* Shadowing of the DIS nuclear structure functions.



*Non-singlet
Reggeon
Exchange*

*Kuti-Weisskopf
behavior*

Origin of Regge Behavior of Deep Inelastic Structure Functions


$$F_{2p}(x) - F_{2n}(x) \propto x^{1/2}$$

Antiquark interacts with target nucleus at energy $\hat{s} \propto \frac{1}{x_{bj}}$

Regge contribution: $\sigma_{\bar{q}N} \sim \hat{s}^{\alpha_R - 1}$

Nonsinglet Kuti-Weisskoff $F_{2p} - F_{2n} \propto \sqrt{x_{bj}}$ at small x_{bj} .

Shadowing of $\sigma_{\bar{q}M}$ produces shadowing of nuclear structure function.



Landshoff, Polkinghorne, Short

Close, Gunion, sjb

Schmidt, Yang, Lu, sjb

Reggeon Exchange

Phase of two-step amplitude relative to one step:

$$\frac{1}{\sqrt{2}}(1 - i) \times i = \frac{1}{\sqrt{2}}(i + 1)$$


Constructive Interference

Depends on quark flavor!

Thus antishadowing is not universal


Different for couplings of γ^* , Z^0 , W^\pm

Critical test: Tagged Drell-Yan



The one-step and two-step processes in DIS on a nucleus.


Coherence at small Bjorken x_B :
 $1/Mx_B = 2\nu/Q^2 \geq L_A$.







If the scattering on nucleon N_1 is via ~~pomeron~~ exchange, the one-step and two-step amplitudes are ~~opposite~~ in phase, thus ~~diminishing~~ the \bar{q} flux reaching N_2 . *Reggeon* *increasing*

→ *Anti-* Shadowing of the DIS nuclear structure functions.

$$Q^2 = 5 \text{ GeV}^2$$







Scheinbein, Yu, Keppel, Morfin, Olness, Owens



Schmidt, Yang; sjb

Nuclear Antishadowing not universal!


Shadowing and Antishadowing of DIS Structure Functions



S. J. Brodsky, I. Schmidt and J. J. Yang,
 “Nuclear Antishadowing in
 Neutrino Deep Inelastic Scattering,”
 Phys. Rev. D 70, 116003 (2004)
 [arXiv:hep-ph/0409279].

Modifies
NuTeV extraction of
 $\sin^2 \theta_W$

Test in flavor-tagged
lepton-nucleus collisions



Predicted nuclear shadowing and antishadowing at $Q^2 = 1 \text{ GeV}^2$

S. J. Brodsky, I. Schmidt and J. J. Yang,
 “Nuclear Antishadowing in
 Neutrino Deep Inelastic Scattering,”
 Phys. Rev. D 70, 116003 (2004)
 [arXiv:hep-ph/0409279].


Shadowing and Antishadowing in Lepton-Nucleus Scattering

- Shadowing: **Destructive Interference** of Two-Step and One-Step Processes
Pomeron Exchange
- Antishadowing: **Constructive Interference** of Two-Step and One-Step Processes!
Reggeon and Odderon Exchange
- Antishadowing is Not Universal!
Electromagnetic and weak currents:
different nuclear effects !

Jian-Jun Yang
Ivan Schmidt
Hung Jung Lu
sjb

Can explain NuTeV result


$$Q^2 = 5 \text{ GeV}^2$$



Scheinbein, Yu, Keppel, Morfin, Olness, Owens

LHC p-A Collisions

Leading-Twist Contribution to Hadron Production on Nuclei



$$G_{q/A}(x_2, p_\perp^2) \simeq A^{\alpha(x_2)} G_{q/N}(x_2, p_\perp^2)$$

$$\frac{d\sigma}{d^3p/E}(pA \rightarrow \pi X) = A^\alpha(x_2) \frac{d\sigma}{d^3p/E}(pN \rightarrow \pi X)$$

Anti-shadowing is quark specific

A

UNAM

October 1, 2010

Novel QCD Phenomena at the LHC

46


Stan Brodsky, SLAC

Physics of Rescattering

- Sivers Asymmetry and Diffractive DIS: New Insights into Final State Interactions in QCD
- Origin of Hard Pomeron
- Structure Functions not Probability Distributions!
Not square of LFWFs
- T-odd SSAs, Shadowing, Antishadowing
- Diffractive dijets/ trijets, doubly diffractive Higgs
- Novel Effects: Color Transparency, Color Opacity, Intrinsic Charm, Odderon


Static

- Square of Target LFWFs
- No Wilson Line
- Probability Distributions
- Process-Independent
- T-even Observables
- No Shadowing, Anti-Shadowing
- Sum Rules: Momentum and J^z
- DGLAP Evolution; mod. at large x
- No Diffractive DIS



Dynamic

- Modified by Rescattering: ISI & FSI
- Contains Wilson Line, Phases
- No Probabilistic Interpretation
- Process-Dependent - From Collision
- T-Odd (Sivers, Boer-Mulders, etc.)
- Shadowing, Anti-Shadowing, Saturation
- Sum Rules Not Proven
- DGLAP Evolution
- Hard Pomeron and Odderon Diffractive DIS




**Hwang,
Schmidt, sjb,
Mulders, Boer
Qiu, Sterman
Collins, Qiu
Pasquini, Xiao,
Yuan, sjb**

Formation of Relativistic Anti-Hydrogen

Measured at CERN-LEAR and FermiLab

Munger, Schmidt, sjb




Coalescence of off-shell co-moving positron and antiproton.

Wavefunction maximal at small impact separation and equal rapidity


"Hadronization" at the Amplitude Level

Hadronization at the Amplitude Level



Construct helicity amplitude using Light-Front Perturbation theory; coalesce quarks via LFWFs

Hadronization at the Amplitude Level




Construct helicity amplitude using Light-Front Perturbation theory; coalesce quarks via LFWFs


Features of LF T-Matrix Formalism

“Event Amplitude Generator”

- Same principle as antihydrogen production: off-shell coalescence
- coalescence to hadron favored at equal rapidity, small transverse momenta
- leading heavy hadron production: D and B mesons produced at large z
- hadron helicity conservation if hadron LFWF has $L^z = 0$
- Baryon AdS/QCD LFWF has aligned and anti-aligned quark spin



Hadronization at the Amplitude Level




Higher Fock State Coalescence $|uuds\bar{s}\rangle$

Asymmetric Hadronization! $D_{s \rightarrow p}(z) \neq D_{s \rightarrow \bar{p}}(z)$

B-Q Ma, sjb


- Quarks and Gluons:
Fundamental constituents of hadrons and nuclei



- *Quantum Chromodynamics (QCD)*

- New Insights from higher space-time dimensions: *AdS/QCD*


- *Light-Front Holography*



- *Hadronization at the Amplitude Level*

- *Light Front Wavefunctions:* analogous to the
Schrodinger wavefunctions of atomic physics

$$\Psi_n(x_i, \vec{k}_{\perp i}, \lambda_i)$$



*Each element of
flash photograph
illuminated
at same LF time*

$$\tau = t + z/c$$

Evolve in LF time


$$P^- = i \frac{d}{d\tau}$$

Eigenstate -- independent of τ



Light-Front Wavefunctions: rigorous representation of composite systems in quantum field theory

$$x = \frac{k^+}{P^+} = \frac{k^0 + k^3}{P^0 + P^3}$$



$$\Psi_n(x_i, \vec{k}_{\perp i}, \lambda_i)$$

$$\sum_i^n x_i = 1$$

$$\sum_i^n \vec{k}_{\perp i} = \vec{0}_\perp$$

Invariant under boosts! Independent of P^μ

Angular Momentum on the Light-Front

$$J^z = \sum_{i=1}^n s_i^z + \sum_{j=1}^{n-1} l_j^z.$$

Conserved
LF Fock state by Fock State

$$l_j^z = -i \left(k_j^1 \frac{\partial}{\partial k_j^2} - k_j^2 \frac{\partial}{\partial k_j^1} \right)$$

n-1 orbital angular momenta

Nonzero Anomalous Moment --> Nonzero orbital angular momentum

Light-Front Wavefunctions

Dirac's Front Form: Fixed $\tau = t + z/c$

$$\Psi(x, k_{\perp}) \quad x_i = \frac{k_i^+}{P^+}$$

Invariant under boosts. Independent of P^{μ}

$$H_{LF}^{QCD} |\psi\rangle = M^2 |\psi\rangle$$

Direct connection to QCD Lagrangian


*Remarkable new insights from AdS/CFT,
the duality between conformal field theory
and Anti-de Sitter Space*

Goal: an analytic first approximation to QCD

- **As Simple as Schrödinger Theory in Atomic Physics**
- **Relativistic, Frame-Independent, Color-Confining**
- **QCD Coupling at all scales**
- **Hadron Spectroscopy**
- **Light-Front Wavefunctions**
- **Form Factors, Hadronic Observables, Constituent Counting Rules**
- **Insight into QCD Condensates**
- **Systematically improvable**

de Teramond, Deur, Shrock, Roberts, Tandy


Applications of AdS/CFT to QCD




Changes in physical length scale mapped to evolution in the 5th dimension z

in collaboration with Guy de Teramond


$$\phi(z)$$



- *Light-Front Holography*



$$\Psi_n(x_i, \vec{k}_{\perp i}, \lambda_i)$$



- *Light Front Wavefunctions:*

Schrödinger Wavefunctions
of Hadron Physics

String Theory



AdS/CFT

Mapping of Poincare' and Conformal $SO(4,2)$ symmetries of 3+1 space to AdS5 space

Goal: First Approximant to QCD

Counting rules for Hard Exclusive Scattering
Regge Trajectories

AdS/QCD

Conformal behavior at short distances
+ Confinement at large distance

QCD at the Amplitude Level



Semi-Classical QCD / Wave Equations

Holography

Boost Invariant 3+1 Light-Front Wave Equations

$J=0, 1, 1/2, 3/2$ plus L

Integrable!



Hadron Spectra, Wavefunctions, Dynamics

LF(3+1)

AdS₅

$$\psi(x, \vec{b}_\perp)$$




$$\phi(z)$$

$$\zeta = \sqrt{x(1-x)} \vec{b}_\perp^2$$




$$z$$

$$\psi(x, \vec{b}_\perp)$$



x



\vec{b}_\perp



$(1-x)$

$$\psi(x, \zeta) = \sqrt{x(1-x)} \zeta^{-1/2} \phi(\zeta)$$

Light-Front Holography: Unique mapping derived from equality of LF and AdS formula for current matrix elements


Light-Front Holography: Map AdS/CFT to 3+1 LF Theory

Relativistic LF radial equation

Frame Independent

$$\left[-\frac{d^2}{d\zeta^2} + \frac{1 - 4L^2}{4\zeta^2} + U(\zeta) \right] \phi(\zeta) = \mathcal{M}^2 \phi(\zeta)$$

$$\zeta^2 = x(1-x)b_{\perp}^2.$$



$$U(\zeta) = \kappa^4 \zeta^2$$

*soft wall
confining potential:*

G. de Teramond, sjb

$$H_{QED}$$

*QED atoms: positronium
and muonium*

$$(H_0 + H_{int}) |\Psi\rangle = E |\Psi\rangle$$

Coupled Fock states

$$\left[-\frac{\Delta^2}{2m_{\text{red}}} + V_{\text{eff}}(\vec{S}, \vec{r}) \right] \psi(\vec{r}) = E \psi(\vec{r})$$

Effective two-particle equation

Includes Lamb Shift, quantum corrections

$$\left[-\frac{1}{2m_{\text{red}}} \frac{d^2}{dr^2} + \frac{1}{2m_{\text{red}}} \frac{\ell(\ell+1)}{r^2} + V_{\text{eff}}(r, S, \ell) \right] \psi(r) = E \psi(r)$$

Spherical Basis r, θ, ϕ

$$V_{\text{eff}} \rightarrow V_C(r) = -\frac{\alpha}{r}$$

Coulomb potential

Bohr Spectrum

Semiclassical first approximation to QED

$$H_{QCD}^{LF}$$

QCD Meson Spectrum

$$(H_{LF}^0 + H_{LF}^I) |\Psi\rangle = M^2 |\Psi\rangle$$

Coupled Fock states

$$\left[\frac{\vec{k}_\perp^2 + m^2}{x(1-x)} + V_{\text{eff}}^{LF} \right] \psi_{LF}(x, \vec{k}_\perp) = M^2 \psi_{LF}(x, \vec{k}_\perp)$$

Effective two-particle equation

$$\zeta^2 = x(1-x)b_\perp^2$$

$$\left[-\frac{d^2}{d\zeta^2} + \frac{-1 + 4L^2}{\zeta^2} + U(\zeta, S, L) \right] \psi_{LF}(\zeta) = M^2 \psi_{LF}(\zeta)$$

Azimuthal Basis ζ, ϕ

$$U(\zeta, S, L) = \kappa^2 \zeta^2 + \kappa^2 (L + S - 1/2)$$

Confining AdS/QCD potential

Semiclassical first approximation to QCD





Fig: Orbital and radial AdS modes in the soft wall model for $\kappa = 0.6$ GeV .

Soft Wall Model




Light meson orbital (a) and radial (b) spectrum for $\kappa = 0.6$ GeV.

- Effective LF Schrödinger wave equation

$$\left[-\frac{d^2}{dz^2} - \frac{1 - 4L^2}{4z^2} + \kappa^4 z^2 + 2\kappa^2(L + S - 1) \right] \phi_S(z) = \mathcal{M}^2 \phi_S(z)$$

with eigenvalues $\mathcal{M}^2 = 2\kappa^2(2n + 2L + S)$. *Same slope in n and L*

- Compare with Nambu string result (rotating flux tube): $M_n^2(L) = 2\pi\sigma(n + L + 1/2)$.




Kirchbach:
Conformal
symmetry

Vector mesons orbital (a) and radial (b) spectrum for $\kappa = 0.54 \text{ GeV}$.

- Glueballs in the bottom-up approach: (HW) Boschi-Filho, Braga and Carrion (2005); (SW) Colangelo, De Fazio, Jugeau and Nicotri(2007).

Spacelike pion form factor from AdS/CFT



Data Compilation
Baldini, Kloe and Volmer

— Soft Wall: Harmonic Oscillator Confinement

— Hard Wall: Truncated Space Confinement

One parameter - set by pion decay constant

de Teramond, sjb

See also: Radyushkin

- We write the Dirac equation

$$(\alpha\Pi(\zeta) - \mathcal{M})\psi(\zeta) = 0,$$

in terms of the matrix-valued operator Π

$$\nu = L + 1$$

$$\Pi_\nu(\zeta) = -i \left(\frac{d}{d\zeta} - \frac{\nu + \frac{1}{2}}{\zeta} \gamma_5 - \kappa^2 \zeta \gamma_5 \right),$$

and its adjoint Π^\dagger , with commutation relations

$$\left[\Pi_\nu(\zeta), \Pi_\nu^\dagger(\zeta) \right] = \left(\frac{2\nu + 1}{\zeta^2} - 2\kappa^2 \right) \gamma_5.$$

- Solutions to the Dirac equation

$$\psi_+(\zeta) \sim z^{\frac{1}{2}+\nu} e^{-\kappa^2 \zeta^2 / 2} L_n^\nu(\kappa^2 \zeta^2),$$

$$\psi_-(\zeta) \sim z^{\frac{3}{2}+\nu} e^{-\kappa^2 \zeta^2 / 2} L_n^{\nu+1}(\kappa^2 \zeta^2).$$

- Eigenvalues

$$\mathcal{M}^2 = 4\kappa^2(n + \nu + 1).$$

Kirchbach:
Conformal
symmetry

- Baryon: twist-dimension $3 + L$ ($\nu = L + 1$)

$$\mathcal{O}_{3+L} = \psi D_{\{\ell_1 \dots D_{\ell_q} \psi D_{\ell_{q+1}} \dots D_{\ell_m}\}} \psi, \quad L = \sum_{i=1}^m \ell_i.$$

$$\mathcal{M}^2 = 4\kappa^2(n + L + 1).$$



Proton Regge Trajectory $\kappa = 0.49\text{GeV}$

Novel QCD Phenomena at the LHC


- Δ spectrum identical to Forkel and Klempt, Phys. Lett. B 679, 77 (2009)

$$4\kappa^2 \text{ for } \Delta n = 1$$


$$4\kappa^2 \text{ for } \Delta L = 1$$

$$2\kappa^2 \text{ for } \Delta S = 1$$

$$\mathcal{M}^2$$



Parent and daughter 56 Regge trajectories for the N and Δ baryon families for $\kappa = 0.5$ GeV





E. Klempt *et al.*: Δ^* resonances, quark models, chiral symmetry and AdS/QCD

H. Forkel, M. Beyer and T. Frederico, JHEP **0707** (2007) 077.

H. Forkel, M. Beyer and T. Frederico, Int. J. Mod. Phys. E **16** (2007) 2794.

Other Applications of Light-Front Holography

- Light baryon spectrum
- Light meson spectrum
- Nucleon form-factors: space-like region
- Pion form-factors: space and time-like regions
- Gravitational form factors of composite hadrons
- n -parton holographic mapping
- Heavy flavor mesons



9-2007
8757A2

hep-th/0501022
hep-ph/0602252
arXiv:0707.3859
arXiv:0802.0514
arXiv:0804.0452

Space-Like Dirac Proton Form Factor

- Consider the spin non-flip form factors

$$F_+(Q^2) = g_+ \int d\zeta J(Q, \zeta) |\psi_+(\zeta)|^2,$$

$$F_-(Q^2) = g_- \int d\zeta J(Q, \zeta) |\psi_-(\zeta)|^2,$$

where the effective charges g_+ and g_- are determined from the spin-flavor structure of the theory.


- Choose the struck quark to have $S^z = +1/2$. The two AdS solutions $\psi_+(\zeta)$ and $\psi_-(\zeta)$ correspond to nucleons with $J^z = +1/2$ and $-1/2$.
- For $SU(6)$ spin-flavor symmetry

$$F_1^p(Q^2) = \int d\zeta J(Q, \zeta) |\psi_+(\zeta)|^2,$$

$$F_1^n(Q^2) = -\frac{1}{3} \int d\zeta J(Q, \zeta) [|\psi_+(\zeta)|^2 - |\psi_-(\zeta)|^2],$$


where $F_1^p(0) = 1$, $F_1^n(0) = 0$.

- Scaling behavior for large Q^2 : $Q^4 F_1^p(Q^2) \rightarrow \text{constant}$ Proton $\tau = 3$



SW model predictions for $\kappa = 0.424$ GeV. Data analysis from: M. Diehl *et al.* Eur. Phys. J. C **39**, 1 (2005).

- Scaling behavior for large Q^2 : $Q^4 F_1^n(Q^2) \rightarrow \text{constant}$ Neutron $\tau = 3$




SW model predictions for $\kappa = 0.424$ GeV. Data analysis from M. Diehl *et al.* Eur. Phys. J. C **39**, 1 (2005).


Spacelike Pauli Form Factor

Preliminary

From overlap of $L = 1$ and $L = 0$ LFWFs



Prediction from AdS/CFT: Meson LFWF



$$\phi_M(x, Q_0) \propto \sqrt{x(1-x)}$$

Increases PQCD prediction for $F_\pi(Q^2)$ by 16/9

Second Moment of Pion Distribution Amplitude

$$\langle \xi^2 \rangle = \int_{-1}^1 d\xi \xi^2 \phi(\xi)$$

$$\xi = 1 - 2x$$

$$\langle \xi^2 \rangle_{\pi} = 1/5 = 0.20 \quad \phi_{asympt} \propto x(1-x)$$

$$\langle \xi^2 \rangle_{\pi} = 1/4 = 0.25 \quad \phi_{AdS/QCD} \propto \sqrt{x(1-x)}$$

$$\text{Lattice (I)} \quad \langle \xi^2 \rangle_{\pi} = 0.28 \pm 0.03$$

Donnellan et al.


$$\text{Lattice (II)} \quad \langle \xi^2 \rangle_{\pi} = 0.269 \pm 0.039$$

Braun et al.

Chiral Features of Soft-Wall AdS/QCD Model

- **Boost Invariant**
- **Trivial LF vacuum.**
- **Massless Pion**
- **Hadron Eigenstates have LF Fock components of different L^z**
- **Proton: equal probability $S^z = +1/2, L^z = 0; S^z = -1/2, L^z = +1$**
- **Self-Dual Massive Eigenstates: Proton is its own chiral partner.**
- **Label State by minimum L as in Atomic Physics**
- **Minimum L dominates at short distances**
- **AdS/QCD Dictionary: Match to Interpolating Operator Twist at $z \rightarrow 0$**

QCD and the LF Hadron Wavefunctions



Features of Soft-Wall AdS/QCD

- Single-variable frame-independent radial Schrodinger equation
- Massless pion ($m_q = 0$)
- Regge Trajectories: universal slope in n and L
- Valid for all integer J & S .
- Dimensional Counting Rules for Hard Exclusive Processes
- Phenomenology: Space-like and Time-like Form Factors
- LF Holography: LFWFs; broad distribution amplitude
- No large N_c limit required
- Add quark masses to LF kinetic energy
- Systematically improvable -- diagonalize H_{LF} on AdS basis

Result: Soft-Wall LFWF for massive constituents

$$\psi(x, \mathbf{k}_\perp) = \frac{4\pi c}{\kappa \sqrt{x(1-x)}} e^{-\frac{1}{2\kappa^2} \left(\frac{\mathbf{k}_\perp^2}{x(1-x)} + \frac{m_1^2}{x} + \frac{m_2^2}{1-x} \right)}$$

LF WF in impact space: soft-wall model with massive quarks

$$\psi(x, \mathbf{b}_\perp) = \frac{c \kappa}{\sqrt{\pi}} \sqrt{x(1-x)} e^{-\frac{1}{2} \kappa^2 x(1-x) \mathbf{b}_\perp^2 - \frac{1}{2\kappa^2} \left[\frac{m_1^2}{x} + \frac{m_2^2}{1-x} \right]}$$

$$z \rightarrow \zeta \rightarrow \chi$$

$$\chi^2 = b^2 x(1-x) + \frac{1}{\kappa^4} \left[\frac{m_1^2}{x} + \frac{m_2^2}{1-x} \right]$$

J/ψ

$\psi_{J/\psi}(x, b)$

LFWF peaks at

$$x_i = \frac{m_{\perp i}}{\sum_j^n m_{\perp j}}$$


where

$$m_{\perp i} = \sqrt{m^2 + k_{\perp}^2}$$

*minimum of LF
energy
denominator*

$$\kappa = 0.375 \text{ GeV}$$


$$m_a = m_b = 1.25 \text{ GeV}$$



$$|\pi^+\rangle = |u\bar{d}\rangle$$


$$m_u = 2 \text{ MeV}$$

$$m_d = 5 \text{ MeV}$$





$$|K^+\rangle = |u\bar{s}\rangle$$

$$m_s = 95 \text{ MeV}$$



$$|D^+\rangle = |c\bar{d}\rangle$$



$$m_c = 1.25 \text{ GeV}$$



$$|\eta_c\rangle = |c\bar{c}\rangle$$

$$|B^+\rangle = |u\bar{b}\rangle$$


$$m_b = 4.2 \text{ GeV}$$



$$|\eta_b\rangle = |b\bar{b}\rangle$$


$$\kappa = 375 \text{ MeV}$$

Hadronization at the Amplitude Level




Construct helicity amplitude using Light-Front Perturbation theory; coalesce quarks via LFWFs

Hadronization at the Amplitude Level




Construct helicity amplitude using Light-Front Perturbation theory; coalesce quarks via LFWFs

Hadronization at the Amplitude Level



AdS/QCD
Hard Wall
Confinement: Capture if $\zeta^2 = x(1-x)b_\perp^2 > \frac{1}{\Lambda_{QCD}^2}$
 i.e.,
 $\mathcal{M}^2 = \frac{k_\perp^2}{x(1-x)} < \Lambda_{QCD}^2$

Hadronization at the Amplitude Level



Higher Fock State Coalescence $|uuds\bar{s}\rangle$

Asymmetric Hadronization! $D_{s \rightarrow p}(z) \neq D_{s \rightarrow \bar{p}}(z)$

B-Q Ma, sjb


Physics of Rescattering

- Sivers Asymmetry and Diffractive DIS: New Insights into Final State Interactions in QCD
- Origin of Hard Pomeron, DDIS
- Structure Functions not Probability Distributions! *Not square of LFWFs*
- T-odd SSAs, Shadowing, Antishadowing
- Diffractive dijets/ trijets, doubly diffractive Higgs
- Novel Effects: Color Transparency, Color Opacity, Intrinsic Charm, Odderon


Applications of Nonperturbative Running Coupling from AdS/QCD

- **Sivers Effect in SIDIS, Drell-Yan**
- **Double Boer-Mulders Effect in DY**
- **Diffraction DIS**
- **Heavy Quark Production at Threshold**


*All involve gluon exchange at small
momentum transfer*



Analytic, defined at all scales, IR Fixed Point



$$\beta^{AdS}(Q^2) = \frac{d}{d \log Q^2} \alpha_s^{AdS}(Q^2) = \frac{\pi Q^2}{4\kappa^2} e^{-Q^2/4\kappa^2}$$



An analytic first approximation to QCD

AdS/QCD + Light-Front Holography

- **As Simple as Schrödinger Theory in Atomic Physics**
- **LF radial variable ζ conjugate to invariant mass squared**
- **Relativistic, Frame-Independent, Color-Confining**
- **QCD Coupling at all scales: Essential for Gauge Link phenomena**
- **Hadron Spectroscopy and Dynamics from one parameter κ**
- **Wave Functions, Form Factors, Hadronic Observables, Constituent Counting Rules**
- **Insight into QCD Condensates: Zero cosmological constant!**
- **Systematically improvable with DLCQ Methods**

Features of AdS/QCD LF Holography

- **Based on Conformal Scaling of Infrared QCD Fixed Point**
- **Conformal template: Use isometries of AdS₅**
- **Interpolating operator of hadrons based on twist, superfield dimensions**
- **Finite $N_c = 3$: Baryons built on 3 quarks -- Large N_c limit not required**
- **Break Conformal symmetry with dilaton**
- **Dilaton introduces confinement -- positive exponent**
- **Origin of Linear and HO potentials: Stochastic arguments (Glazek); General 'classical' potential for Dirac Equation (Hoyer)**
- **Effective Charge from AdS/QCD at all scales**
- **Conformal Dimensional Counting Rules for Hard Exclusive Processes**

*“One of the gravest puzzles of
theoretical physics”*

**DARK ENERGY AND
THE COSMOLOGICAL CONSTANT PARADOX**

A. ZEE

*Department of Physics, University of California, Santa Barbara, CA 93106, USA
Kavil Institute for Theoretical Physics, University of California,
Santa Barbara, CA 93106, USA
zee@kitp.ucsb.edu*

$$(\Omega_{\Lambda})_{QCD} \sim 10^{45}$$

$$\Omega_{\Lambda} = 0.76(\text{expt})$$

$$(\Omega_{\Lambda})_{EW} \sim 10^{56}$$

*QCD Problem Solved if Quark and Gluon condensates reside
within hadrons, not vacuum!*

R. Shrock, sjb

arXiv:0905.1151 [hep- th], Proc. Nat'l. Acad. Sci., (in press);
“Condensates in Quantum Chromodynamics and the Cosmological Constant”

Chiral magnetism (or magnetohydrochironics)

Aharon Casher and Leonard Susskind

Tel Aviv University Ramat Aviv, Tel-Aviv, Israel

(Received 20 March 1973)

I. INTRODUCTION


The spontaneous breakdown of chiral symmetry in hadron dynamics is generally studied as a vacuum phenomenon.¹ Because of an instability of the chirally invariant vacuum, the real vacuum is “aligned” into a chirally asymmetric configuration.

On the other hand an approach to quantum field theory exists in which the properties of the vacuum state are not relevant. This is the parton or constituent approach formulated in the infinite-momentum frame.² A number of investigations have indicated that in this frame the vacuum may be regarded as the structureless Fock-space vacuum. Hadrons may be described as nonrelativistic collections of constituents (partons). In this framework the spontaneous symmetry breakdown must be attributed to the properties of the hadron’s wave function and not to the vacuum.³


*Light-Front
Formalism*

Simple physical argument for "in-hadron" condensate

Roberts, Shrock, Tandy, sjb



Gribov pairs



B-Meson

*Use Dyson-Schwinger Equation for bound-state quark propagator:
find confined condensate*


$$\langle B | \bar{q}q | B \rangle \text{ not } \langle 0 | \bar{q}q | 0 \rangle$$

Bethe-Salpeter Analysis

Maris,
Roberts, Tandy

$$f_H P^\mu = Z_2 \int^\Lambda \frac{d^4 q}{(2\pi)^4} \frac{1}{2} [T_H \gamma_5 \gamma^\mu \mathcal{S}(\frac{1}{2}P + q) \Gamma_H(q; P) \mathcal{S}(\frac{1}{2}P - q)]$$

f_H Meson Decay Constant
 T_H flavor projection operator,
 $Z_2(\Lambda), Z_4(\Lambda)$ renormalization constants
 $\mathcal{S}(p)$ dressed quark propagator
 $\Gamma_H(q; P) = F.T. \langle H | \psi(x_a) \bar{\psi}(x_b) | 0 \rangle$
 Bethe-Salpeter bound-state vertex amplitude.




$$i\rho_\zeta^H \equiv \frac{-\langle q\bar{q} \rangle_\zeta^H}{f_H} = Z_4 \int^\Lambda \frac{d^4 q}{(2\pi)^4} \frac{1}{2} [T_H \gamma_5 \mathcal{S}(\frac{1}{2}P + q) \Gamma_H(q; P) \mathcal{S}(\frac{1}{2}P - q)]$$

In-Hadron Condensate!


$$f_H m_H^2 = -\rho_\zeta^H \mathcal{M}_H \quad \mathcal{M}_H = \sum_{q \in H} m_q$$

$$m_\pi^2 \propto (m_q + m_{\bar{q}}) / f_\pi \quad \text{GMOR}$$

Higher Light-Front Fock State of Pion Simulates DCSB




$$f_\pi P^+ = \langle 0 | \bar{q} \gamma^5 \gamma^+ q | \pi \rangle$$



Instantaneous quark propagator contribution to ρ_π derived from higher Fock state

$$i\rho_\pi = \langle 0 | \bar{q} \gamma^5 q | \pi \rangle$$



Higher Fock state acts like mass insertion

PHYSICAL REVIEW C **82**, 022201(R) (2010)**New perspectives on the quark condensate**Stanley J. Brodsky,^{1,2} Craig D. Roberts,^{3,4} Robert Shrock,⁵ and Peter C. Tandy⁶¹*SLAC National Accelerator Laboratory, Stanford University, Stanford, California 94309, USA*²*Centre for Particle Physics Phenomenology: CP³-Origins, University of Southern Denmark, Odense 5230 M, Denmark*³*Physics Division, Argonne National Laboratory, Argonne, Illinois 60439, USA*⁴*Department of Physics, Peking University, Beijing 100871, China*⁵*C.N. Yang Institute for Theoretical Physics, Stony Brook University, Stony Brook, New York 11794, USA*⁶*Center for Nuclear Research, Department of Physics, Kent State University, Kent, Ohio 44242, USA*

(Received 25 May 2010; published 18 August 2010)

We show that the chiral-limit vacuum quark condensate is qualitatively equivalent to the pseudoscalar meson leptonic decay constant in the sense that they are both obtained as the chiral-limit value of well-defined gauge-invariant hadron-to-vacuum transition amplitudes that possess a spectral representation in terms of the current-quark mass. Thus, whereas it might sometimes be convenient to imagine otherwise, neither is essentially a constant mass-scale that fills all spacetime. This means, in particular, that the quark condensate can be understood as a property of hadrons themselves, which is expressed, for example, in their Bethe-Salpeter or light-front wave functions.

Determinations of the vacuum Gluon Condensate

$$\langle 0 | \frac{\alpha_s}{\pi} G^2 | 0 \rangle [\text{GeV}^4]$$


-0.005 ± 0.003 from τ decay.

Davier et al.

$+0.006 \pm 0.012$ from τ decay. Geshkenbein, Ioffe, Zyablyuk

$+0.009 \pm 0.007$ from charmonium sum rules

Ioffe, Zyablyuk



*Consistent with zero
vacuum condensate*

Quark and Gluon condensates reside within hadrons, not vacuum

Casher and Susskind

Maris, Roberts, Tandy

Shrock and sjb

- **Bound-State Dyson Schwinger Equations**
- **AdS/QCD**
- **Analogous to finite size superconductor**
- **Implications for cosmological constant --
Eliminates 45 orders of magnitude conflict**

R. Shrock, sjb

ArXiv:0905.1151

*“One of the gravest puzzles of
theoretical physics”*

**DARK ENERGY AND
THE COSMOLOGICAL CONSTANT PARADOX**

A. ZEE

*Department of Physics, University of California, Santa Barbara, CA 93106, USA
Kavil Institute for Theoretical Physics, University of California,
Santa Barbara, CA 93106, USA
zee@kitp.ucsb.edu*

$$(\Omega_{\Lambda})_{QCD} \sim 10^{45}$$

$$\Omega_{\Lambda} = 0.76(\text{expt})$$

$$(\Omega_{\Lambda})_{EW} \sim 10^{56}$$

*QCD Problem Solved if Quark and Gluon condensates reside
within hadrons, not vacuum!*

R. Shrock, sjb

arXiv:0905.1151 [hep- th], Proc. Nat'l. Acad. Sci., (in press);
“Condensates in Quantum Chromodynamics and the Cosmological Constant”

- **Color Confinement: Maximum Wavelength of Quark and Gluons**
- **Conformal symmetry of QCD coupling in IR**
- **Conformal Template (BLM, CSR, BFKL scale)**
- **Motivation for AdS/QCD**
- **QCD Condensates inside of hadronic LFWFs**
- **Technicolor: confined condensates inside of technihadrons -- alternative to Higgs**
- **Simple physical solution to cosmological constant conflict with Standard Model**

Roberts, Shrock, Tandy, and sjb

QCD Myths

- **Anti-Shadowing is Universal**
- **ISI and FSI are higher twist effects and universal**
- **High transverse momentum hadrons arise only from jet fragmentation -- baryon anomaly!**
- **heavy quarks only from gluon splitting**
- **renormalization scale cannot be fixed**
- **QCD condensates are vacuum effects**
- **Infrared Slavery**
- **Nuclei are composites of nucleons only**
- **Real part of DVCS arbitrary**