

Observation of the $K^+ \rightarrow \pi^+ \nu \bar{\nu}$ decay and measurement of its branching ratio by the NA62 experiment at CERN



Jürgen Engelfried

Instituto de Física
Universidad Autónoma de San Luis Potosí, Mexico
on behalf of the NA62 Collaboration



XV Latin American Symposium on High Energy Physics
(SILAFEA 2024)

CINVESTAV, Mexico City

November 4, 2024



- The Golden Modes $K \rightarrow \pi \nu \bar{\nu}$ in the SM and beyond
- NA62 after LS2: detector upgrades and performance
- New measurement of $\mathcal{B}(K^+ \rightarrow \pi^+ \nu \bar{\nu})$



Presentation follows the talk by Joel Swallow at the CERN EP Seminar on Sept 24



$K \rightarrow \pi \nu \bar{\nu}$: Precision test of the Standard Model



SM: Z-penguin & box diagrams



• $\mathcal{B}(K \rightarrow \pi \nu \bar{\nu})$ highly suppressed in SM

• GIM mechanism & maximum CKM suppression $s \rightarrow d$ transition: $\sim \frac{m_t}{m_W} \left| V_{ts}^* V_{td} \right|$

• Theoretically clean \Rightarrow high precision SM predictions

• Dominated by short distance contributions.

• Hadronic matrix element extracted from $\mathcal{B}(K \rightarrow \pi^0 \ell^+ \nu_\ell)$ decays via isospin rotation.

Mode	SM Branching Ratio [1]	SM Branching Ratio [2]	Experimental Status
$K^+ \rightarrow \pi^+ \nu \bar{\nu}$	$(8.60 \pm 0.42) \times 10^{-11}$	$(7.86 \pm 0.61) \times 10^{-11}$	$(10.6 \pm 4.0) \times 10^{-11}$ NA62 16–18
$K_L \rightarrow \pi^0 \nu \bar{\nu}$	$(2.94 \pm 0.15) \times 10^{-11}$	$(2.68 \pm 0.30) \times 10^{-11}$	$< 2 \times 10^{-9}$ KOTO (2021 data)

¹Recent SM calculations [1: [Buras et al. EPJC 82 \(2022\) 7, 615](#)] [2: [D'Ambrosio et al. JHEP 09 \(2022\) 148](#)]
 (Differences in SM calculations from choice of CKM parameters: see [Eur.Phys.J.C 84 \(2024\) 4, 377](#))



$K \rightarrow \pi \nu \bar{\nu}$: Beyond the Standard Model




- Correlations between BSM contributions to BRs of K^+ and K_L modes [JHEP 11 (2015) 166].
 - Must measure both to discriminate between BSM scenarios.
- Correlations with other observables (ϵ'/ϵ , ΔM_B , B-decays) [JHEP 12 (2020) 097][PLB 809 (2020) 135769].
- Leptoquarks [EPJ.C 82 (2022) 4, 320], Interplay between CC and FCNC [JHEP 07 (2023) 029], NP in neutrino sector [EPJ.C 84 (2024) 7, 680] and additional scalar/tensor contributions [JHEP 12 (2020) 186][arXiv:2405.06742] ...


- **Green:** CKM-like flavour structure
 - Models with Minimal Flavour Violation
- **Blue:** new flavour-violating interactions where LH or RH currents dominate
 - Z' models with pure LH/RH couplings
- **Red:** general NP models without above constraints
- **Grossman-Nir Bound:** model-independent relation

[PLB 398 (1997) 163-168]

$$\frac{\mathcal{B}(K_L \rightarrow \pi^0 \nu \bar{\nu}) \tau_{K^+}}{\mathcal{B}(K^+ \rightarrow \pi^+ \nu \bar{\nu}) \tau_{K_L}} \lesssim 1$$

$$\Rightarrow \mathcal{B}(K_L \rightarrow \pi^0 \nu \bar{\nu}) \lesssim 4.3 \mathcal{B}(K^+ \rightarrow \pi^+ \nu \bar{\nu})$$





Charged Kaons at the Main Injector

April 15, 1998

**A Proposal for a Precision Measurement of the Decay
 $K^+ \rightarrow \pi^+ \nu \bar{\nu}$ and Other Rare K^+ Processes at Fermilab
 Using the Main Injector**

R. Coleman, P.S. Cooper*, M. Crisler, C. Milstene†, E. Ramberg, R. S. Tschirhart
 Fermi National Accelerator Laboratory, Batavia, IL, USA

A. Kozhevnikov, V. Kubarovsky, V. Kurshetsov, L. Landsberg, V. Molchanov, V. Obraztsov, V. Polyakov,
 D. Vavilov, O. Yushchenko
 Institute of High Energy Physics, Serpukhov, Russia

J. Engelfried, A. Morelos
 Instituto de Fisica, Universidad Autonoma de San Luis Potosi, Mexico

K. Lang, J.L. Ritchie*
 University of Texas at Austin, Austin, Texas 78712

† Resident at Fermilab, on leave from Tel Aviv University.

* Cospokesmen: P.S. Cooper, pcooper@fnal.gov, (630) 840-2629
 J.L. Ritchie, ritchie@hep.utexas.edu, (512) 471-4488

VIII MEXICAN WORKSHOP ON PARTICLES AND FIELDS

November 14-20, 2001

Zacatecas, México

Invited Speakers

J. Bonn	Mainz, Germany
A. J. Buras	Munich, Germany
R. Cashmore	CERN, Switzerland
J. Conway	Rutgers, USA
G. Contreras	Mérida, México
P. S. Cooper	FNAL, USA
J. Dos Anjos	CBPF, Brazil
Robin Earbacher	BNL, USA
Ada Farilla	INFN-Rome, Italy
R. Flores-Mendieta	San Luis Potosí, México
N. Fornengo	Torino, Italy
Elizabeth Jenkins	San Diego, USA
W. C. Louis	LANL, USA
O. G. Miranda	CINVESTAV-IPN, México
A. Mondragón	UNAM, México
Giulia Panzeri	INFN, Italy
S. Pastor	Munich, Germany
J. S. Russ	Carnegie Mellon, USA
A. Santoro	LAFEX/CBPF, Brazil
N. Tarnopolski	Helsinki, Finland
A. Wagner	DESY, Germany
J. Wambach	Darmstadt, Germany
H.-J. Weber	Charlottesville, USA
A. Wirzba	FZ Jülich, Germany
J. Wudka	Riverside, USA
C. P. Yuan	Michigan, USA

Topics

Heavy Quarks and CP
Neutrino Physics
Chiral Dynamics
Higgs Particle

National Organizing Committee

J. L. Díaz Cruz	IF-BUAP (Chair)
G. Herrera	CINVESTAV
M. Mondragón	IE-UNAM
N. Nellen	ICN-UNAM
M. A. Pérez	CINVESTAV

Local Organizing Committee

J. Engelfried	IF-UASLP
A. A. Espinoza	EE-UAZ
A. Gutiérrez	EE-UAZ
M. Kerchbach	EE-UAZ (Chair)
J. Ontz Saavedra	EE-UAZ
J. M. Rivera Juárez	EE-UAZ

Organized by the División Partículas
and Fields of the Sociedad Mexicana
de Física

taller2001@dpyc.smf.mx
<http://dpyc.smf.mx/Workshop/>

Design: Comunic


Sponsored by



Centro Latinoamericano de Física
Centro Latinoamericano de Física
Latin American Center of Physics



Andrzej J. Buras (Munich, Germany): *CP Violation, Rare Decays and the CKM Matrix*
Peter S. Cooper (Fermilab): *Rare Kaon Decays*



Andrzej Buras (right, with John Conway), Restaurant “Mamá Inés”, Zacatecas, Mexico, November 2001


NA62: The K^+ factory at the CERN north area



The NA62 Experiment at CERN



~200 collaborators from ~30 institutions.



- **Primary goal:** measurement of $\mathcal{B}(K^+ \rightarrow \pi^+ \nu \bar{\nu})$
- **New Technique:** K^+ decay-in-flight
- **Results:** [\[PLB 791 \(2019\) 156\]](#) [\[JHEP 11 \(2020\) 042\]](#) [\[JHEP 06 \(2021\) 093\]](#)
- **Broader physics programme:**
 - Rare K^+ decays (e.g. $K^+ \rightarrow \pi^+ \gamma \gamma$ [\[PLB 850 \(2024\) 138513\]](#))
 - LNV/LFV decays (e.g. $K^+ \rightarrow \pi^- (\pi^0) e^+ e^+$ [\[PLB 830 \(2022\) 137172\]](#))
 - Exotics (e.g. Dark photon [\[PRL 133 \(2024\) 11, 111802\]](#))
- **Data taking**
 - 2016 Commissioning + Physics run (45 days).
 - 2017 Physics run (160 days).
 - 2018 Physics run (217 days).
 - 2021 Physics run (85 days [10 beam dump]).
 - 2022 Physics run (215 days).
 - 2023 Physics run (150 days [10 beam dump]).
 - 2024 Physics run ongoing ...



The NA62 Collaboration

- Birmingham
- Bratislava
- Bristol
- Bucharest
- CERN
- Dubna
- George Mason
- Ferrara
- Firenze
- Frascati
- Glasgow
- Louvain-la-Neuve
- EPFL Lausanne
- Mainz
- IN2P3 Marseille
- INR Moscow
- Max Planck München
- Napoli
- Perugia
- Pisa
- Prague
- Protvino
- Roma I
- Roma II
- San Luis Potosí
- SLAC
- Torino
- TRIUMF
- Vancouver

~ 230 collaborators



The Mexicans in NA62




$K^+ \rightarrow \pi^+ \nu \bar{\nu}$ at NA62

NA62 Strategy:

- Tag K^+ and measure momentum.
- Identify π^+ and measure momentum.
- Match K^+ and π^+ in time & form vertex.
 - Determine $m_{miss}^2 = (P_K - P_\pi)^2$
- Reject any additional activity.

NA62 Performance Keystones:

- $\mathcal{O}(100)$ ps timing between detectors
- $\mathcal{O}(10^4)$ background suppression from kinematics
- $> 10^7$ muon rejection
- $> 10^7$ rejection of π^0 from $K^+ \rightarrow \pi^+ \pi^0$ decays



Decay mode	Branching Ratio [PDG]
$K^+ \rightarrow \mu^+ \nu_\mu$	$(63.56 \pm 0.11) \%$
$K^+ \rightarrow \pi^+ \pi^0$	$(20.67 \pm 0.08) \%$
$K^+ \rightarrow \pi^+ \pi^+ \pi^-$	$(5.583 \pm 0.024) \%$
$K^+ \rightarrow \pi^+ \pi^- e^+ \nu_e$	$(4.247 \pm 0.024) \times 10^{-5}$




$K^+ \rightarrow \pi^+ \nu \bar{\nu} \quad (8.60 \pm 0.42) \times 10^{-11}$ [SM]

[Buras et al. EPJC 82 \(2022\) 7, 615](#)




Kinematic constraints & signal regions

$$m_{miss}^2 = (P_{K^+} - P_{\pi^+})^2$$

$\mathcal{O}(10^4)$ background suppression from kinematics



The story so far: $K^+ \rightarrow \pi^+ \nu \bar{\nu}$ with 2016–18 data



Data-taking year	[Reference]	N_{bg}	$N_{\pi\nu\bar{\nu}}^{SM,exp}$	N_{obs}
2016	[PLB 791 (2019) 156]	$0.152^{+0.093}_{-0.035}$	0.267 ± 0.020	1
2017	[JHEP 11 (2020) 042]	1.46 ± 0.33	2.16 ± 0.13	2
2018	[JHEP 06 (2021) 093]	$5.42^{+0.99}_{-0.75}$	7.58 ± 0.40	17
2016–18	[JHEP 06 (2021) 093]	$7.03^{+1.05}_{-0.82}$	10.01 ± 0.42	20

Statistical combination:

$$\mathcal{B}(K^+ \rightarrow \pi^+ \nu \bar{\nu}) = (10.6^{+4.0}_{-3.4} |_{\text{stat}} \pm 0.9_{\text{syst}}) \times 10^{-11} \text{ at } 68\% \text{ CL}$$

$$\text{In background-only hypothesis: } p = 3.4 \times 10^{-4} \Rightarrow \text{significance} = 3.4\sigma.$$




NA62 beamline & detector

[JINST 12 (2017) 05, P05025]



Secondary 75 GeV/c beam:

- π^+ : 70 % , p : 23 % , K^+ : 6 %



- Designed & optimised for study of $K^+ \rightarrow \pi^+ \nu \bar{\nu}$:
 - Particle tracking: beam particle (GTK) & downstream tracks (STRAW)
 - PID: K^+ - KTAG, π^+ - RICH, Calorimeters (LKr, MUV1,2), MUV3 (μ detector)
 - Comprehensive veto systems: CHANTI (beam interactions), LAV, LKr, IRC, SAC (γ)



11




Particle ID performance : 2021–22 data




Calorimeters

- Use BDT classifier for LKr & MUV1,2
- + MUV3 (fast μ^+ detector)



RICH

Designed to distinguish between π^+/μ^+ with 15 – 35 GeV/c.





$$\varepsilon(\pi \text{ ID}) = (73.00 \pm 0.01) \%$$

$$P(\mu^+ \text{ misID as } \pi^+) = (1.3 \pm 0.2) \times 10^{-8}$$

12



Comprehensive photon veto system: 2021–22 **NA62**



- Probability of $K^+ \rightarrow \pi^+ \pi^0$, $\pi^0 \rightarrow \gamma\gamma$ events passing all photon veto conditions: $\eta_{\pi^0} = (1.72 \pm 0.07) \times 10^{-8}$



- Meets target: combined γ/π^0 rejection of $\mathcal{O}(10^8)$.

Upgrading NA62



- 2016–18 analysis proved NA62 technique.
- Limitation: tight cuts to reject backgrounds \Rightarrow reduces signal efficiency.
- To improve: need new tools to control background.




Upgrading NA62



- 2016–18 analysis proved NA62 technique.
- Limitation: tight cuts to reject backgrounds \Rightarrow reduces signal efficiency.
- To improve: need new tools to control background.

Background	N(exp) 2018 (S2)
Upstream	$2.76^{+0.90}_{-0.70}$
$K^+ \rightarrow \pi^+\pi^0$	0.52 ± 0.05
$K^+ \rightarrow \mu^+\nu$	0.45 ± 0.06
$K^+ \rightarrow \pi^+\pi^-e^+\nu$	0.41 ± 0.10
$K^+ \rightarrow \pi^+\pi^+\pi^-$	0.17 ± 0.08
Total	$4.31^{+0.91}_{-0.72}$



Largest backgrounds:

1. Upstream
2. $K^+ \rightarrow \pi^+\pi^0$


Veto by detecting previously missed particles...



New upstream vetos: VetoCounter & ANTIO



[FELIX readout: [Streaming Readout Workshop talk 2021](#)]



VetoCounter

- Detect particles from decays upstream of final collimator.
- **Factor ~3 rejection** with ~2% accidental veto.

ANTIO

- Detect particles up to ~1 m from beam line.
- **Reject ~20% of upstream background** with <1% signal loss.




[JINST 15 (2020) C07007]

[SPSC report 2023][EP Newsletter, Dec21]



4th GTK station

- Si Pixel detector exposed to $\sim 1\text{GHz}$ beam.
- Essential for $K^+ - \pi^+$ matching.
 - Measures K^+ 3-mom. & time
- 4th GTK station improves efficiency & pileup resilience.




More info: [\[Kleimenova_VERTEX2023\]](#)




Summary of NA62 upgrades

- New detectors, installed during LS2:
 - 4th GTK (Kaon beam tracker) & rearrange GTK achromat (GTK2 upstream of scraper).
 - New upstream veto (**VetoCounter**) & veto hodoscope (**ANTI0**) upstream of decay volume.
 - Additional veto detector (**HASC2**) at end of beam-line.
- Intensity increased by $\sim 35\%$ with respect to 2018 [450 \rightarrow 600 MHz].
- Improvements to the trigger configuration.

New detectors
installed in 2021:




Beam intensity: 2018 vs 2022



- Average beam intensity increased.
- NA62 “Full intensity” with 4.8s spill = 600 MHz

2021 instantaneous beam intensity



- Remove events in first 1s of 4.8s spill for 2021 data only.
- DAQ overwhelmed by instantaneous rates up to 10x higher than design.

Analysis strategy



Triggers:

- **Minimum Bias:** $K^+ \rightarrow \mu^+ \nu$
- **Normalisation:** $K^+ \rightarrow \pi^+ \pi^0$
- **Signal:** $K^+ \rightarrow \pi^+ \nu \bar{\nu}$ candidates

- RICH multiplicity (reference time)
- Signal in CHODs
- No signal in MUV3 (μ veto)
- Tag K^+ (≥ 5 KTAG sectors)
- < 40 GeV in LKr ($\pi^0/\gamma/e$ veto)
- LAV veto (downstream of vertex).

Common conditions

+ add more conditions

Selection:

- **Normalisation** $K^+ \rightarrow \pi^+ \pi^0$: 1 downstream track (only); identified as π^+ ; $K^+ - \pi^+$ matching (space & time); upstream vetos.
- **Signal** $K^+ \rightarrow \pi^+ \nu \bar{\nu}$ candidates: same as normalisation selection + full photon and detector multiplicity cuts (reject all extra activity).



27




Bayesian classifier for $K^+ - \pi^+$ matching

Example of selection update





- **Inputs:** spatial (CDA) & time (ΔT_+) matching, intensity/pileup (N_{GTK}) [prior]
- Models for PDFs/Prior from $K^+ \rightarrow \pi^+ \pi^+ \pi^-$ data.



- **Output:** posterior probability of GTK track = true K^+
 - Use likelihoods of kaons (K) and pileup (P)
 - Likelihood ratio used to select true match when $N_{GTK} > 1$

$\epsilon = 80\%$ P^P mistag = 6% P^K mistag = 2%



- Efficiency improved (+10%) and mistagging probability maintained.



Signal sensitivity

- Normalisation channel: $K^+ \rightarrow \pi^+\pi^0$, momentum range $p \in [15,45]$ GeV/c.

Effective number of K^+ decays, N_K :

$$N_K = \frac{N_{\pi\pi} D_0}{\mathcal{B}_{\pi\pi} A_{\pi\pi}}$$

Number of normalisation events $\rightarrow N_{\pi\pi} D_0$
 Downscaling factor of normalisation trigger (generally 400) $\rightarrow D_0$
 Branching ratio of $K^+ \rightarrow \pi^+\pi^0$ decay $\rightarrow \mathcal{B}_{\pi\pi}$
 Acceptance of normalisation selection $\rightarrow A_{\pi\pi}$

Single event sensitivity:

(Branching ratio corresponding to expectation of 1 event)

$$\mathcal{B}_{SES} = \frac{1}{N_K \epsilon_{RV} \epsilon_{trig} A_{\pi\nu\bar{\nu}}}$$

Random veto efficiency $\rightarrow \epsilon_{RV}$
 Trigger efficiency (ratio) $\rightarrow \epsilon_{trig}$
 Signal selection acceptance $\rightarrow A_{\pi\nu\bar{\nu}}$

Number of expected SM events:

(For comparison to previous results use $\mathcal{B}_{\pi\nu\bar{\nu}}^{SM} = 8.4 \times 10^{-11}$ [JHEP 11 (2015) 166], but results are independent of this choice)


$$N_{\pi\nu\bar{\nu}}^{SM} = \frac{\mathcal{B}_{\pi\nu\bar{\nu}}^{SM}}{\mathcal{B}_{SES}}$$

Acceptances

Analysis is performed in (5 GeV/c) bins of momentum:

$$N_{\pi\nu\bar{\nu}}^{\text{exp}}(p_i) = \frac{\mathcal{B}_{\pi\nu\bar{\nu}}^{\text{SM}}}{\mathcal{B}_{\pi\pi}^{\text{SES}}(p_i)} = \frac{\mathcal{B}_{\pi\nu\bar{\nu}}^{\text{SM}}}{\mathcal{B}_{\pi\pi}} \frac{A_{\pi\nu\bar{\nu}}(p_i)}{A_{\pi\pi}(p_i)} D_0 N_{\pi\pi}(p_i) \varepsilon_{\text{trig}}(p_i) \varepsilon_{\text{RV}}$$

Acceptances



Case	OLD 2018 (S2)	NEW 2021–22	
Norm.	11.8%	13.4%	+15%
Signal	(6.37±0.64)%	(7.61±0.18)%	+20%

- Increased selection efficiencies.
 - New K-pi matching technique.
 - Re-tuned vertex conditions.
 - Relaxation of some vetos.
- Improved precision (plus improved systematic uncertainty evaluation).




Acceptances evaluated at 0 intensity.
Intensity dependence captured in ε_{RV}

30

Trigger efficiencies

Analysis is performed in (5 GeV/c) bins of momentum:

$$N_{\pi\nu\bar{\nu}}^{\text{exp}}(p_i) = \frac{\mathcal{B}_{\pi\nu\bar{\nu}}^{\text{SM}}}{\mathcal{B}_{\text{SES}}(p_i)} = \frac{\mathcal{B}_{\pi\nu\bar{\nu}}^{\text{SM}} A_{\pi\nu\bar{\nu}}(p_i)}{\mathcal{B}_{\pi\pi} A_{\pi\pi}(p_i)} D_0 N_{\pi\pi}(p_i) \epsilon_{\text{trig}}(p_i) \epsilon_{\text{RV}}$$



$$\epsilon_{\text{trig}} = \frac{\epsilon_{\text{sig}}}{\epsilon_{\text{norm}}}$$

$$\epsilon_{\text{trig}}(\text{new}) = (85.9 \pm 1.4) \%$$

$$\epsilon_{\text{trig}}(\text{2018}) = (89 \pm 5) \%$$

- Trigger efficiency ratio:
 - **New:** several components in both normalisation & signal triggers: **partial cancellation.**
 - **Old:** in 2016–18 data normalise with fully independent min bias trigger (**no cancellation**).
- Improved precision by factor 3 with reduced systematic uncertainty.




31



Random veto

ϵ_{RV} is independent of track momentum
(related to additional activity only)

$$N_{\pi\nu\bar{\nu}}^{exp}(p_i) = \frac{\mathcal{B}_{\pi\nu\bar{\nu}}^{SM}}{\mathcal{B}_{SES}(p_i)} = \frac{\mathcal{B}_{\pi\nu\bar{\nu}}^{SM} A_{\pi\nu\bar{\nu}}(p_i)}{\mathcal{B}_{\pi\pi} A_{\pi\pi}(p_i)} D_0 N_{\pi\pi}(p_i) \epsilon_{trig}(p_i) \epsilon_{RV}$$



- ϵ_{RV} = Random Veto Efficiency:
 - $1 - \epsilon_{RV}$ = Probability of rejecting a signal event due to additional activity.
- Balance:
 - Strict vetos \Rightarrow lower efficiency
 - Loose vetos \Rightarrow higher background
- Operational intensity higher but re-tuning vetos means ϵ_{RV} is comparable:

$$\epsilon_{RV}(\text{new}, \overline{\lambda_{21-22}} \approx 600 \text{ MHz}) = (63.6 \pm 0.6) \%$$

$$\epsilon_{RV}(\text{old}, \overline{\lambda_{2018}} \approx 400 \text{ MHz}) = (66 \pm 1) \%$$

Signal sensitivity results

$$N_K = \frac{N_{\pi\pi} D_0}{\mathcal{B}_{\pi\pi} A_{\pi\pi}}$$

$$\mathcal{B}_{SES} = \frac{1}{N_K \epsilon_{RV} \epsilon_{trig} A_{\pi\nu\bar{\nu}}}$$

- Display integrals (15–45 GeV/c, 2021+22) for summary tables.
- * Acceptances evaluated at 0 intensity.

$N_{\pi\pi}$	Normalisation $K^+ \rightarrow \pi^+ \pi^0$	2.0×10^8
$A_{\pi\pi}$	Normalisation acceptance	$(13.410 \pm 0.005)\%$
N_K	Effective K^+ decays	2.9×10^{12}
$A_{\pi\nu\bar{\nu}}$	Signal acceptance	$(7.6 \pm 0.2)\%$
ϵ_{trig}	Trigger efficiency	$(85.9 \pm 1.4)\%$
ϵ_{RV}	Random veto efficiency	$(63.6 \pm 0.6)\%$
\mathcal{B}_{SES}	Single event sensitivity	$(0.84 \pm 0.03) \times 10^{-11}$


$$N_{\pi\nu\bar{\nu}}^{exp} = \frac{\mathcal{B}_{\pi\nu\bar{\nu}}^{SM}}{\mathcal{B}_{SES}}$$

Assuming $\mathcal{B}_{\pi\nu\bar{\nu}}^{SM} = 8.4 \times 10^{-11}$:
 2021–22: $N_{\pi\nu\bar{\nu}} = 10.00 \pm 0.34$
 c.f. 2016–18 : $N_{\pi\nu\bar{\nu}} = 10.01 \pm 0.42$

↓
 Double expected signal by including 21–22 data.

- Significant improvement in SES uncertainty:
 - old: 6.3% → new: 3.5%. Due to:
 - trigger efficiency cancellations
 - improved procedures for evaluation of acceptances and ϵ_{RV}


Optimum NA62 intensity



- Saturation of expected signal yield with intensity. Mainly due to:
 - Paralyzable effects from TDAQ dead time and trigger veto windows.
 - Offline selection, due to veto conditions.
- Main sources of uncertainty for model:
 - Online time-dependent mis-calibrations.
 - Fit uncertainty.
- **From August 2023 operate at optimal intensity (~75% of full) to maximise $\pi V V$ sensitivity**
 - Maximise signal yield
 - lower expected background
 - Higher DAQ efficiency
- **Studies of 2021–22 data at high intensity were crucial to establish optimal intensity.**



Background regions & background estimations



- Backgrounds from kinematic misconstruction tails in m_{miss}^2

Number of events passing signal selection in background region



Control sample events in Signal Regions

$$N_{bg} = N_{bkgR} \cdot f_{tail} = N_{bkgR} \cdot \frac{N_{SR}^{CS}}{N_{bkgR}^{CS}}$$

Kinematic tail fraction: measured in control sample

Control sample events in Background Region

Backgrounds from kinematic tails



$K^+ \rightarrow \pi^+ \pi^+ \pi^-$



- Use MC to measure f_{tail} :

$N_{bg}(K^+ \rightarrow \pi^+ \pi^+ \pi^-) = 0.11 \pm 0.03$

Radiative decays: $K^+ \rightarrow \pi^+ \pi^0 \gamma$ & $K^+ \rightarrow \mu^+ \nu \gamma$



- $K^+ \rightarrow \pi^+ \pi^0 \gamma$: included with “kinematic tails” estimation.
 - Suppression: photon vetos, rejection with additional γ is 30x stronger.
 - Estimation: MC + measured single photon rejection efficiency: $N_{bg}(K^+ \rightarrow \pi^+ \pi^0 \gamma) = 0.07 \pm 0.01$
 - Validation: m_{miss}^2 control regions (CR1,2 - see later)
- $K^+ \rightarrow \mu^+ \nu \gamma$: not included in “kinematic tails” estimation if γ overlaps μ^+ at LKr (leading to misID as π^+)
 - Suppression: based on $(P_K - P_\mu - P_\gamma)^2$ and E_γ with $\gamma =$ LKr cluster (mis)associated to muon.
 - Necessary for 2021–22 data, since Calorimetric PID degraded at higher intensities.
 - Estimation: min. Bias data control sample with signal in MUV3: $N_{bg}(K^+ \rightarrow \mu^+ \nu \gamma) = 0.8 \pm 0.4$
 - Validation: data sample without $K^+ \rightarrow \mu^+ \nu \gamma$ veto and PID = “less pion-like” (Calo BDT bins below π^+ bin).



38



Other backgrounds

- $K^+ \rightarrow \pi^+ \pi^- e^+ \nu$ (K_{e4})

- No clean control samples for K_{e4} in data: use 2×10^9 simulated decays.

$$N_{bg}(K^+ \rightarrow \pi^+ \pi^- e^+ \nu) = N_K \epsilon_{RV} \epsilon_{trig} \mathcal{B}_{K_{e4}} A_{K_{e4}}$$

Effective # of K^+ → N_K
 Random veto & trigger efficiencies → $\epsilon_{RV} \epsilon_{trig}$
 Acceptance: $A_{K_{e4}} = \frac{N_{MC}^{sel}}{N_{MC}^{gen}} = (1.3 \pm 0.3_{stat}) \times 10^{-8}$
 Branching ratio of K_{e4} (from PDG) → $\mathcal{B}_{K_{e4}}$

$N_{bg}(K^+ \rightarrow \pi^+ \pi^- e^+ \nu) = 0.89^{+0.34}_{-0.28}$

- $K^+ \rightarrow \pi^0 \ell^+ \nu$ and $K^+ \rightarrow \pi^+ \gamma \gamma$:

- Evaluated with simulations.
- Negligible contributions to total background.

$N_{bg}(K^+ \rightarrow \pi^0 \ell^+ \nu) < 1 \times 10^{-3}$

$N_{bg}(K^+ \rightarrow \pi^+ \gamma \gamma) = 0.01 \pm 0.01$

Upstream background evaluation

$$N_{bg} = \sum_i N_i f_{cda} P_i^{match}$$

N Upstream Reference Sample:
signal selection but invert CDA cut (CDA > 4mm)

f_{cda} Scaling factor : bad cda \rightarrow good cda

P_{match} Probability to pass $K^+ - \pi^+$ matching

Calculate using bins (i) of $(\Delta T_+, N_{GTK})$

[Updated to fully data-driven procedure]

$$N = 51 \quad f_{CDA} = 0.20 \pm 0.03 \quad \langle P_{match} \rangle = 73\%$$


$$N_{bg}(\text{Upstream}) = 7.4^{+2.1}_{-1.8}$$

- Upstream reference sample contains all known upstream mechanisms.

- N provides normalisation.


- f_{CDA} depends only on geometry.

- P_{match} depends on $(\Delta T_+, N_{GTK})$.



Upstream background validation

- Invert & loosen upstream vetos to enrich with different mechanisms:
 - Interaction-enriched: Val1,2,7,8
 - Accidental-enriched: Val3,4,5,6,9,10.
- All independent.
- Expectations and observations are in good agreement.
- Number of events rejected by VetoCounter:
 - (i.e. events in signal region with associated VC signal)
 - $N_{exp}^{VC rej.} = 6.9 \pm 1.4$, $N_{obs}^{VC rej.} = 9$
- VetoCounter is essential to control upstream background.



Summary of expectations

Backgrounds

$K^+ \rightarrow \pi^+ \pi^0 (\gamma)$	0.83 ± 0.05
$K^+ \rightarrow \pi^+ \pi^0$	0.76 ± 0.04
$K^+ \rightarrow \pi^+ \pi^0 \gamma$	0.07 ± 0.01
$K^+ \rightarrow \mu^+ \nu (\gamma)$	1.70 ± 0.47
$K^+ \rightarrow \mu^+ \nu$	0.87 ± 0.19
$K^+ \rightarrow \mu^+ \nu \gamma$	0.82 ± 0.43
$K^+ \rightarrow \pi^+ \pi^+ \pi^-$	0.11 ± 0.03
$K^+ \rightarrow \pi^+ \pi^- e^+ \nu$	$0.89^{+0.34}_{-0.28}$
$K^+ \rightarrow \pi^0 \ell^+ \nu$	< 0.001
$K^+ \rightarrow \pi^+ \gamma \gamma$	0.01 ± 0.01
Upstream	$7.4^{+2.1}_{-1.8}$
Total	$11.0^{+2.1}_{-1.9}$

Signal Sensitivity

$$\mathcal{B}_{SES} = (0.84 \pm 0.03) \times 10^{-11}$$

$$N_{\pi\nu\bar{\nu}}^{SM,exp} = \frac{\mathcal{B}_{\pi\nu\bar{\nu}}^{SM}}{\mathcal{B}_{SES}}$$

Assuming $\mathcal{B}_{\pi\nu\bar{\nu}}^{SM} = 8.4 \times 10^{-11}$:

2021–22: $N_{\pi\nu\bar{\nu}} = 10.00 \pm 0.34$



c.f. 2016–18 : $N_{\pi\nu\bar{\nu}} = 10.01 \pm 0.42$

Expected signal doubled by including 2021–22 data

- $N_{\pi\nu\bar{\nu}}^{SM}$ per SPS spill: 2.5×10^{-5} in 2022
 - c.f. 1.7×10^{-5} in 2018. \Rightarrow signal yield increased by 50%.
- Sensitivity for $BR \sim \sqrt{S+B}/S = 0.5$
 - Similar but improved with respect to 2018 analysis for same amount of data.

Control regions

2021 – 22 data




- Good agreement in control regions validates background expectations.



Signal regions



2021 – 22 data




Expected SM signal, $N_{\pi^+\nu\bar{\nu}}^{SM} \approx 10$

Expected background, $N_{bg} = 11.0^{+2.1}_{-1.9}$

Observed, $N_{obs} = 31$



1D projection with differential background predictions & SM signal expectation [not a fit]:



Results: 2021–22 Data



- Measure $\mathcal{B}_{\pi\nu\bar{\nu}}$ and 68% (1σ) confidence interval using a profile likelihood ratio test statistic $q(\theta)$.
- Use 6 (momentum bin) categories



$$\mathcal{B}_{21-22}(K^+ \rightarrow \pi^+ \nu \bar{\nu}) = (16.0^{+5.0}_{-4.5}) \times 10^{-11} = \left(16.0 \begin{matrix} +4.8 \\ -4.2 \end{matrix} \text{stat} \begin{matrix} +1.4 \\ -1.3 \end{matrix} \text{syst} \right) \times 10^{-11}$$



Evaluate statistical-only component by repeating procedure assuming exact knowledge of signal and background expectations.


47



Combining NA62 results: 2016–22




- Integrating 2016–22 data: $N_{bg} = 18_{-2}^{+3}$, $N_{obs} = 51$.



Combining NA62 results: 2016–22





- Integrating 2016–22 data: $N_{bg} = 18_{-2}^{+3}$, $N_{obs} = 51$.
- Background-only hypothesis **p-value = 2×10^{-7}** \Rightarrow **significance $Z > 5$**



Combining NA62 results: 2016–22

- Integrating 2016–22 data: $N_{bg} = 18_{-2}^{+3}$, $N_{obs} = 51$.

- Background-only hypothesis **p-value = 2×10^{-7} \Rightarrow significance $Z > 5$**



$$\mathcal{B}_{16-22}(K^+ \rightarrow \pi^+ \nu \bar{\nu}) = (13.0_{-2.9}^{+3.3}) \times 10^{-11} = \left(13.0_{-2.7}^{+3.0} \text{stat} \left[\begin{matrix} +1.3 \\ -1.2 \end{matrix} \right] \text{syst} \right) \times 10^{-11}$$

Results in context

BNL E787/E949 experiment
[Phys.Rev.D 79 (2009) 092004]


$$\mathcal{B}_{\pi\nu\bar{\nu}}^{16-18} = (10.6^{+4.1}_{-3.5}) \times 10^{-11}$$

[JHEP 06 (2021) 093]

$$\mathcal{B}_{\pi\nu\bar{\nu}}^{21-22} = (16.0^{+5.0}_{-4.5}) \times 10^{-11}$$


$$\mathcal{B}_{\pi\nu\bar{\nu}}^{16-22} = (13.0^{+3.3}_{-2.9}) \times 10^{-11}$$

- NA62 results are consistent
- Central value moved up (now 1.5–1.7 σ above SM)
- Fractional uncertainty decreased: 40% to 25%
- Bkg-only hypothesis rejected with significance $Z > 5$



- Study of $K^+ \rightarrow \pi^+ \nu \bar{\nu}$ decay using NA62 2021-2022 data
 - Improved signal yield per SPS spill by 50 %
 - $N_{\text{bg}} = 11.0_{-1.9}^{+2.1}$, $N_{\text{obs}} = 31$
 - $\mathcal{B}_{21-22}(K^+ \rightarrow \pi^+ \nu \bar{\nu}) = (16.0_{-4.5}^{+5.0}) \times 10^{-11}$
- Combining with 2016-2018 data for full 2016-22 result
 - $N_{\text{bg}} = 18_{-2}^{+3}$, $N_{\text{obs}} = 51$
 - $\mathcal{B}_{16-22}(K^+ \rightarrow \pi^+ \nu \bar{\nu}) = (13.0_{-2.9}^{+3.3}) \times 10^{-11}$
 - Background-only hypothesis rejected with **significance** > 5
- **First observation of $K^+ \rightarrow \pi^+ \nu \bar{\nu}$ decay**
 \mathcal{B} consistent with SM prediction within 1.7σ
- Need full NA62 data-set to clarify SM agreement or tension
 - Analysis of 2023 data in progress
 - Processing of 2024 data in progress
 - Taking more data in 2025 and 2026(?)

Supplemental




54

Mid 2018 - installed TCX Collimator

Much improved shielding - blocking almost all upstream decay paths.


Sketch only





2021 - addition of VetoCounter

Upstream decays can be detected and actively vetoed.

Sketch only








HASC2 veto



- $K^+ \rightarrow \pi^+\pi^0$ was 2nd largest background for 2018 analysis.

- Addition of HASC2:

- 30% less $K^+ \rightarrow \pi^+\pi^0$
- 18% less $K^+ \rightarrow \pi^+\pi^+\pi^-$
- 3.5% less $K^+ \rightarrow \mu^+\nu$
- with only 1.5% signal loss.




$K^+ \rightarrow \mu^+ \nu \gamma$ Background

- Kinematically select $K^+ \rightarrow \mu^+ \nu \gamma$ events:


$$m_{miss}^2(K_{\mu 2\gamma}) = (P_K - P_\mu - P_\gamma)^2$$

- P_K : 4-momentum of K^+ from GTK (as normal)
- P_μ : 4-momentum of track with μ^+ mass hypothesis.
- P_γ : reconstructed from energy and position of LKr cluster (and position of $K^+ - \mu^+$ vertex).

Validation: data sample with PID = “less pion-like” (Calo BDT bins below π^+ bin).



Evaluate background expectation using $\mu \nu \gamma$ control sample from MinimumBias trigger, not applying Calorimetric BDT classifier and MUV3 signal:




- Before $K^+ \rightarrow \mu^+ \nu \gamma$ veto: found excess of events at $p > 35$ GeV/c in Region 2 relative to 2016–18 data.
- Additional background identified and studied in data control samples & MC.
- $K^+ \rightarrow \mu^+ \nu \gamma$ veto added to selection criteria for final analysis.

39

Results in context





- Fractional uncertainty: 25%
- Bkg-only hypothesis rejected with significance $Z > 5$
- **Observation of the $K^+ \rightarrow \pi^+ \nu \bar{\nu}$ decay with BR consistent with SM prediction, within 1.7σ**
- Need full NA62 data-set to clarify SM agreement or tension



$$\mathcal{B}_{\pi\nu\bar{\nu}}^{16-22} = (13.0^{+3.3}_{-2.9}) \times 10^{-11}$$


$$2\sigma \text{ range : } [7.4 - 19.7] \times 10^{-11}$$

Cedar & KTAG : K^+ tagging with threshold Cherenkov counter **NA62**



- 75 GeV Unseparated hadron beam
 $\pi^+ : 70\%$, $p : 23\%$, $K^+ : 6\%$.

- Use fixed diaphragm to select ONLY Cherenkov light from K^+ (adjust diaphragm width and gas pressure in CEDAR to ensure powerful K^+/π^+ discrimination).





New Cedar-H : installed in 2023




- CEDAR-W filled with N_2 at 1.7 bar was biggest contributor to material in beam line ($39 \times 10^{-3} X_0$).
- New CEDAR-H filled with H_2 at 3.8 bar:
 - Reduces material to $7.3 \times 10^{-3} X_0$: reducing multiple scattering.
 - But new optics required to account for different optical properties of H_2 .
- Successful test beam in 2022 (at CERN,H6) and installation in NA62 in early 2023.

• Cedar-H Performance at NA62:

- **>99.5% efficiency** for 5-fold coincidence.
- **π^+ mistag probability: 10^{-4}**
- **~65ps time resolution**
- **30% reduction** in elastically scattered beam particles.



Photon veto performance



- Probability of $K^+ \rightarrow \pi^+\pi^0$ events with $\pi^0 \rightarrow \gamma\gamma$ passing full photon vetos:

Number of events passing full $\pi^+\nu\bar{\nu}$ selection in $\pi^+\pi^0$ region

$$\eta_{\pi^0} = \frac{N_{\pi^+\pi^0 R}}{N_{\pi\pi} D_0 \epsilon_{trig} \epsilon_{RV}}$$

Number of selected normalisation events

Random veto efficiency




Normalisation trigger downscaling and efficiency

$$\eta_{\pi^0} = (1.72 \pm 0.07) \times 10^{-8}$$

- Combined γ/π^0 rejection of $\mathcal{O}(10^8)$.

Bayesian classifier for $K^+ - \pi^+$ matching


- Inputs: spatial (CDA) & time (ΔT_+) matching, intensity/pileup (N_{GTK}) [prior]
 - Models for PDFs/Prior from $K^+ \rightarrow \pi^+ \pi^+ \pi^-$ data.



Example of selection update



- **Output:** posterior probability of GTK track = true K^+
 - Use likelihoods of kaons (K) and pileup (P)
 - Likelihood ratio used to select true match when $N_{GTK} > 1$
 - Efficiency improved (+10%) and mistagging probability maintained.

Kinematic regions



- **Signal regions:**
- **Control regions:**
 - Used to validate background predictions.
- **Background regions:**
 - Used as “reference samples” for some background estimates.

3D signal regions definition



CR3D: control region for events in SR in 2 out of 3 dimensions.

$$m_{miss}^2 = (P_K - P_\pi)^2$$

Default: GTK

Alternative: Nominal beam = $m_{miss,nom}^2$

Default: STRAW



Alternative: $|p|$ from RICH (use as a velocity spectrometer) = $m_{miss,RICH}^2$

Background mechanism: $K^+ \rightarrow \mu^+ \nu \gamma$


- $K^+ \rightarrow \mu^+ \nu \gamma$ decay with fairly energetic photon ($E_\gamma > 5 \text{ GeV}$) and high momentum μ^+ ($p \gtrsim 35 \text{ GeV}/c$).
- γ and μ^+ hit LKr together and are misidentified as a π^+ .
- No rejection power from photon vetos (LKr γ cluster associated to track).
- Additional γ naturally shifts $m_{miss}^2 = (P_K - P_\pi)^2$ towards higher values (i.e. towards signal regions).



Example event (2022 data):




Sketch only



Background evaluation: $K^+ \rightarrow \mu^+ \nu \gamma$


- Evaluate background expectation using $\mu \nu \gamma$ control sample from MinimumBias (MB) trigger.
 - Not applying Calorimetric BDT classifier and a signal in MUV3.



- Kinematically select $K^+ \rightarrow \mu^+ \nu \gamma$ events:

$$m_{miss}^2(K_{\mu\gamma}) = (P_K - P_\mu - P_\gamma)^2$$
 - P_K : 4-momentum of K^+ from GTK (as normal)
 - P_μ : 4-momentum of track with μ^+ mass hypothesis.
 - P_γ : reconstructed from energy (subtracting MIP energy deposit) and position of LKr cluster (and position of $K^+ - \mu^+$ vertex).

$$N_{bg}(K^+ \rightarrow \mu^+ \nu \gamma) = N_{\mu\nu\gamma}^{MB} D_{MB} \frac{\epsilon_{signal}}{\epsilon_{MB}} P_{misID}$$



Not included in kinematic tails calculation because the tails sample imposes Calorimetric PID= μ^+ , while here there is misID of $\mu^+ \gamma \Rightarrow \pi^+$.

Background rejection: $K^+ \rightarrow \mu^+ \nu \gamma$




Minimum Bias Data
Events with MUV3 association and
 $|m_{miss}^2(K_{\mu 2\gamma})|^2 < 0.01 \text{ GeV}^2/c^4$

veto $K^+ \rightarrow \mu^+ \nu \gamma$ events with:

- $|m_{miss}^2(K_{\mu 2\gamma})|^2 < 0.01 \text{ GeV}^2/c^4$ → c.f. resolution $\sim 0.0025 \text{ GeV}^2/c^4$
- $E_\gamma > 5 \text{ GeV}$
- μ^+ -like RICH PID.

- Veto conditions established using data control samples and MC.
- $K^+ \rightarrow \mu^+ \nu \gamma$ Veto \Rightarrow 20x background suppression with 0.4% signal loss.



- Why different to 2016–18 analysis?
 - Calorimetric PID degraded:
 - Higher intensity in 2021–22 data (in particular, affects MUV1,2).
 - Training of BDT classifier.

Upstream background evaluation

$$N_{bg} = \sum_i N_i f_{cda} P_{match}$$

N Upstream Reference Sample:
 signal selection but invert CDA cut (CDA > 4mm)

f_{cda} Scaling factor : bad cda \rightarrow good cda


P_{match} Probability to pass $K^+ - \pi^+$ matching

Calculate using bins (i) of $(\Delta T_+, N_{GTK})$
 [Updated to fully data-driven procedure]

$$N = 51 \quad f_{CDA} = 0.20 \pm 0.03 \quad \langle P_{match} \rangle = 73\%$$

$$N_{bg}(\text{Upstream}) = 7.4^{+2.1}_{-1.8}$$

- Upstream reference sample contains all known upstream mechanisms.
- N provides normalisation.
- f_{CDA} depends only on geometry.
- P_{match} depends on $(\Delta T_+, N_{GTK})$.



Results in context: the long story of $K^+ \rightarrow \pi^+ \nu \bar{\nu}$

Experimental measurements:

- Camerini et al. [PRL 23 (1969) 326-329]
- Klems et al. [PRD 4 (1971) 66-80]
- Ljung et al. [PRD 8 (1973) 1307-1330]
- Cable et al. [PRD 8 (1973) 3807-3812]
- Asano et al. [PLB 107 (1981) 159]

E787 :

- [PRL 64 (1990) 21-24]
- [PRL 70 (1993) 2521-2524]
- [PRL 76 (1996) 1421-1424]
- [PRL 79 (1997) 2204-2207]
- [PRL 84 (2000) 3768-3770]
- [PRL 88 (2002) 041803]

E949 (+E787)


- [PRL 93 (2004) 031801]
- [PRL 101 (2008) 191802]

NA62:

- 2016 data: [PLB 791 (2019) 156]
- 2016+17 data: [JHEP 11 (2020) 042]
- 2016-18 data: [JHEP 06 (2021) 093]
- 2016-22 data: this result.

Theory:

- [Phys.Rev. 163 (1967) 1430-1440]
- [PRD 10 (1974) 897]
- [Prog.Theor.Phys. 65 (1981)]
- [PLB 133 (1983) 443-448]
- [PLB 192 (1987) 201-206]
- [Nucl.Phys.B 304 (1988) 205-235]
- [PRD 54 (1996) 6782-6789]
- [PRD 76 (2007) 034017]
- [PRD 78 (2008) 034006]
- [PRD 83 (2011) 034030]
- [JHEP 11 (2015) 033]
- [JHEP 09 (2022) 148]



$K_L \rightarrow \pi^0 \nu \bar{\nu}$ at KOTO



- Located at J-Park 30 GeV main ring.
- KOTO continues data-taking to reach sensitivity $< 10^{-10}$
- Planned future program (KOTO-2) key part of high priority hadron hall extension plans at J-PARC.

[K.shiomi : Kaons @ CERN 2023]

[T.Nomura : Kaons @ J-PARC 2024]




Signature of $K_L \rightarrow \pi^0 \nu \bar{\nu} = "2\gamma + \text{Nothing} + P_t"$



Assuming 2γ from π^0 ,
 Calculate z vertex on the beam axis
 $M^2(\pi^0) = 2E_1 E_2 (1 - \cos \theta)$
 Calculate π^0 transverse momentum

Branching ratio (BR)




Grossman-Nir bound:
 indirect limit from relation to $BR(K^+ \rightarrow \pi^+ \nu \bar{\nu})$;
 Calc'd from NA62 results (2021) with 1σ region



Latest results:

Results of the 2021 data analysis

Final PT vs Z plot Black: observed Single Event Sensitivity = $(9.26 \pm 0.03_{\text{stat}} \pm 0.75_{\text{sys}}) \times 10^{-10}$
 Red: expected BG
 Contour: signal MC



$N_{\text{observed}} = 0$
BR($K_L \rightarrow \pi^0 \nu \bar{\nu}$) < 2.1 × 10⁻⁹ (90% C.L.)

Obtained the world best limit.

We will submit the paper on this result soon.



physics programme

$K^+ \rightarrow \pi^+ \nu \bar{\nu}$

Rare Decays

Forbidden Decays

Exotics

- $K^+ \rightarrow \pi^+ \nu \bar{\nu}$: [PLB 791 (2019) 156] [JHEP 11 (2020) 042] [JHEP 06 (2021) 093][this talk]
- $K^+ \rightarrow \pi^+ X$: [JHEP 03 (2021) 058] [JHEP 06 (2021) 093]
- $(K^+ \rightarrow \pi^+ \pi^0, \pi^0) \rightarrow$ invisible [JHEP 02 (2021) 201]

- $K^+ \rightarrow \pi^+ \pi^0, \pi^0 \rightarrow e^+ e^-$ [prelim. Spring 2024]
- $K^+ \rightarrow \pi^+ \gamma \gamma$ [PLB 850 (2024) 138513]
- Tagged neutrino [prelim. 2023]

- $K^+ \rightarrow \pi^0 \pi \mu e$ [prelim. Spring 2024]
- $K^+ \rightarrow (\pi^0) \pi^- e^+ e^+$ [PLB 830 (2022) 137172]
- $K^+ \rightarrow \mu^- \nu e^+ e^+$ [PLB838 (2023) 137679]
- $K^+ \rightarrow \pi \mu e, \pi^0 \rightarrow \mu^- e^+$ [PRL 127 (2021) 13, 131802]

- Beam dump dark photon searches:
 - $A' \rightarrow \ell^+ \ell^-$ [PRL 133 (2024) 11, 111802] [JHEP 09 (2023) 035]
 - $A' \rightarrow$ hadrons [prelim. Spring 2024]