

# Lepton number violating/conserving heavy baryon decays, in presence of two almost degenerated heavy neutrinos

Fabiola Fortuna

Work done in collaboration with: Gerardo Hernández, Diego Portillo and Genaro Toledo

PRD 112, 015010 (2025)

XIX Mexican Workshop on Particles and Fields, 2025



# Contents

1 Motivation

2 LNC & LNV baryon decays

3 Conclusions

# Motivation

# Motivation

Several extensions of the Standard Model (SM) aiming at explaining **oscillation phenomena** invoke the introduction of **right-handed (RH) neutrinos**.


The embedding of the seesaw mechanism onto the SM is one of the most economical mechanisms for the **generation of neutrino masses** and **lepton mixings**.



The presence of relatively light RH neutrinos (sterile fermions from the gauge point of view), which have non-negligible mixings with the active ones, leads to the modification of the **charged and neutral lepton currents**.

# Lepton Number Violation

A typical example of LNV is the neutrinoless double beta decay,  $0\nu\beta\beta$ .



This process is only possible if neutrinos are Majorana.

# LNC & LNV baryon decays

# Framework

- Simplified SM extensions that involve the addition of extra neutral Majorana fermions.

The leptonic charged current is modified as follows:

$$\mathcal{L}_{c.c.} = -\frac{g}{\sqrt{2}} \textcolor{violet}{U}_{\alpha i} \bar{\ell}_\alpha \gamma_\mu P_L \nu_i W_\mu^- + \text{h.c.}, \quad (1)$$

where  $P_L = (1 - \gamma_5)/2$  is the left-handed chirality projector, the subindex  $i$  refers to the physical neutrino states (3 light plus 2 heavy states), and the subindex  $\alpha$  represents the flavor of the charged leptons.

Matrix elements for the heavy-light mixings are denoted by


$$U_{\alpha i} = e^{i\phi_{\alpha i}} |U_{\alpha i}|, \quad \alpha = e, \mu, \tau, \quad i = 4, 5 \quad (2)$$

where  $\phi_{\alpha i}$  is the phase of the associated mixing element.

\*More realistic models (motivated to accommodate the mass generation mechanism <sup>1</sup>) invoke two almost degenerate Majorana neutrinos, with masses in the GeV region.


<sup>1</sup>Eur. Phys. J. C 82, 1030 (2022), J. High Energy Phys. 07 (2024) 060.

# LNC & LNV baryon decays



LNC

$$m_P + m_{\ell_\alpha} \leq m_N \leq m_A - m_B - m_{\ell_\beta}$$



LNV

$$m_P + m_{\ell_\beta} \leq m_N \leq m_A - m_B - m_{\ell_\alpha}$$

## Two almost degenerate case

The total squared amplitude can be recast into a single Majorana neutrino process, provided the following conditions are satisfied:

- the masses are  $m_4 \simeq m_5 \equiv m_N$  and  $\Delta m_N \equiv m_5 - m_4 \gtrsim 0$
- the decay widths are  $\Gamma_4 \simeq \Gamma_5 \equiv \Gamma_N$
- the mixing parameters are thus also required to fulfill  
 $|U_{\alpha 4}| |U_{\beta 4}| = |U_{\alpha 5}| |U_{\beta 5}| = |U_{\alpha N}| |U_{\beta N}|$

# Full LNC & LNV results, in the two almost degenerate case

In the LNC decays, we focus on lepton flavor violating (LFV) decays , i.e., with  $\ell_\alpha \neq \ell_\beta$  , to isolate the sterile neutrino contributions.

The average square amplitudes become:

$$\overline{|\mathcal{M}_{\text{LNC}}(2N)|^2} = \overline{|\mathcal{M}_{\text{LNC}}(1N)|^2} R[y, \psi_-], \quad (3)$$


$$\overline{|\mathcal{M}_{\text{LNV}}(2N)|^2} = \overline{|\mathcal{M}_{\text{LNV}}(1N)|^2} R[y, \psi_+], \quad (4)$$

where  $y = \Delta m_N / \Gamma_N$ ,  $\psi_\alpha \equiv \phi_{\alpha 5} - \phi_{\alpha 4}$  and  $R[y, \psi]$  is the recast function

$$R[y, \psi_\mp] \equiv 2 \left\{ 1 + \kappa(y) (\cos(\psi_\alpha \mp \psi_\beta) - y \sin(\psi_\alpha \mp \psi_\beta)) \right\}, \quad (5)$$

with  $\kappa(y) = 1/(1 + y^2)$ .

# Recast function



# Detector length considerations

- We take into account the probability that the on-shell neutrino will decay inside the detector.

This effect is incorporated in the following **probability weight**:

$$P_\nu = 1 - \text{Exp} \left( -L_{\text{det}} \Gamma_N \frac{m_N}{|p_N|} \right), \quad (6)$$

where

- $L_{\text{det}}$  is the detector length
- $\Gamma_N$  the total decay width of the heavy neutrino
- $p_N$  the heavy neutrino three-momentum in the laboratory frame of the decaying baryon

Therefore, the differential decay rate can be written as:

$$d\Gamma = P_\nu |\overline{\mathcal{M}}|^2 dPS, \quad (7)$$





being  $dPS$  the corresponding differential Phase Space according to the transition.

# Considerations for calculation

- The neutrino decay width is taken as dependent on
  - the mass of the heavy neutrinos,
  - the heavy-light mixing parameters,
  - and the corresponding open channels.
- The exclusion regions were obtained under the assumption of an upper limit for the branching fraction of the four-body  $\Lambda_b$  decays to be  $\text{BR} \leq \mathcal{O}(10^{-8})$ , motivated by the expected sensitivity to the process  $\Lambda_b \rightarrow \Lambda_c(p)\pi\mu\mu$  at CMS and LHCb experiments <sup>2</sup>.



<sup>2</sup>Jhovanny Mejia-Guisao, et al. Exploring GeV-scale Majorana neutrinos in lepton-number-violating  $\Lambda_b^0$  baryon decays. *Phys. Rev. D* 96 (2017) 1, 015039

# Exclusion region: heavy-light mixing vs heavy neutrino mass




- \* LNV processes with only one Majorana neutrino.
- \* The dotted lines include the detector effects.

# Scenario with two quasi-degenerate Majorana neutrinos



# Branching ratios



\*The dotted lines include detector effects.

# Conclusions

# Conclusions

- We have studied the LNC and LNV four-body decays of heavy baryons.
- Our analysis extends beyond the simplified assumption of a single heavy neutrino mixing with the active sector, by including **two nearly degenerate heavy neutrinos**.
- We profit from the so-called **recast function**, to illustrate the effect of the two-almost degenerate heavy neutrinos, as compared to the single case.
- A large set of heavy hadron transitions was considered.
- We determined the potential exclusion region for the mass and heavy-light mixing parameters of the neutrinos driving the decay.

# Thank you!

# Framework

- Simplified SM extensions that involve the addition of  $N$  extra neutral Majorana fermions.
- We don't make any assumptions about the mechanism responsible for neutrino mass generation (i.e., treating neutrino masses and lepton mixings as independent).

The leptonic charged current is modified as follows:

$$\mathcal{L}_{c.c.} = -\frac{g}{\sqrt{2}} U_{\alpha i} \bar{\ell}_\alpha \gamma_\mu P_L \nu_i W_\mu^- + \text{h.c.}, \quad (8)$$

where  $P_L = (1 - \gamma_5)/2$  is the left-handed chirality projector, the subindex  $i$  refers to the physical neutrino states (3 light plus  $N$  heavy states), and the subindex  $\alpha$  represents the flavor of the charged leptons.

Matrix elements for the heavy-light mixings are denoted by

$$U_{\alpha i} = e^{i\phi_{\alpha i}} |U_{\alpha i}|, \quad \alpha = e, \mu, \tau, \quad i = 4, 5 \quad (9)$$

where  $\phi_{\alpha i}$  is the phase of the associated mixing element.

\*More realistic models (motivated to accommodate the mass generation mechanism <sup>3</sup>) invoke two almost degenerate Majorana neutrinos, with masses in the GeV region.

<sup>3</sup>Eur. Phys. J. C 82, 1030 (2022), J. High Energy Phys. 07 (2024) 060.

# LNC $B_A^{\{0,\mp\}} \rightarrow B_B^{\{\pm,0\}} P^\mp \ell_\alpha^\mp \ell_\beta^\pm$ Decays

We focus on lepton flavor violating (LFV) decays, i.e., with  $\ell_\alpha \neq \ell_\beta$ , to isolate the sterile neutrino contributions. We obtain the amplitude for

$$B_A^{\{0,-\}}(p_A) \rightarrow B_B^{\{+,0\}}(p_B) P^-(p_P) \ell_\beta^-(p_1) \ell_\alpha^+(p_2)$$

$$\mathcal{M}_{\text{LNC}} = G p_P^\nu \sum_{i=4,5} U_{\beta i} U_{\alpha i}^* \ell_{\mu\nu}^{\text{LNC}} P_{1i} H^\mu(p_B, p_A), \quad (10)$$

where we have defined  $G \equiv G_F^2 V_{AB} V_P f_P$ , with  $V_P$  and  $V_{AB}$  the mixing quark elements of the CKM matrix, and  $f_P$  the decay constant of the meson state. The leptonic part is given by

$$\ell_{\mu\nu}^{\text{LNC}} \equiv \bar{u}(p_1) \gamma_\mu \not{a}_1 \gamma_\nu (1 - \gamma_5) v(p_2), \quad (11)$$

where  $a_1 \equiv p_A - p_B - p_1$  is the momentum carried out by either of the heavy neutrinos, and we have defined

$$P_{1i} \equiv \frac{1}{a_1^2 - m_i^2 + i m_i \Gamma_i}. \quad (12)$$

# Hadronic part

- The meson production coming from the  $W$  is parameterized by  $if_P p_P^\nu$ .
- Baryon transition matrix element

$$H^\mu(p_B, p_A) \equiv \langle B_B(p_B) | J^\mu | B_A(p_A) \rangle, \quad (13)$$

in the most general form is parameterized by six form factors, as follows

$$\begin{aligned} \langle B_B(p_B) | J_\mu | B_A(p_A) \rangle = \\ \bar{u}(p_B) \left[ f_1(q^2) \gamma_\mu + if_2(q^2) \frac{\sigma_{\mu\nu} q^\nu}{M_A} + \frac{q_\mu f_3(q^2)}{M_A} \right. \\ \left. + g_1(q^2) \gamma_\mu \gamma_5 + ig_2(q^2) \frac{\sigma_{\mu\nu} q^\nu \gamma_5}{M_A} + \frac{q_\mu g_3(q^2) \gamma_5}{M_A} \right] u(p_A), \end{aligned} \quad (14)$$

with  $q^2 = (p_A - p_B)^2$  the squared momentum transferred in the baryonic transition.

# Interference terms

After squaring the amplitude for the LNC case, the interference terms can be stated in terms of the relative phases as

$$U_{\alpha 4}^* U_{\beta 4} U_{\alpha 5} U_{\beta 5}^* P_{14} P_{15}^* + U_{\alpha 4} U_{\beta 4}^* U_{\alpha 5}^* U_{\beta 5} P_{14}^* P_{15} = \\ |U_{\alpha 4}| |U_{\beta 4}| |U_{\alpha 5}| |U_{\beta 5}| \left( e^{i(\psi_\alpha - \psi_\beta)} P_{14} P_{15}^* + e^{-i(\psi_\alpha - \psi_\beta)} P_{14}^* P_{15} \right)$$


where  $\psi_\alpha \equiv \phi_{\alpha 5} - \phi_{\alpha 4}$ .

This squared amplitude can be recast into a single Majorana neutrino process, provided the masses satisfy  $m_4 \simeq m_5 \equiv m_N$  and  $\Delta m_N \equiv m_5 - m_4 \gtrsim 0$ , the decay widths are  $\Gamma_4 \simeq \Gamma_5 \equiv \Gamma_N$ , the mixing parameters are thus also required to fulfill  $|U_{\alpha 4}| |U_{\beta 4}| = |U_{\alpha 5}| |U_{\beta 5}| = |U_{\alpha N}| |U_{\beta N}|$  and the relation between the  $P_{1i} P_{1j}^*$

$$P_{14} P_{15}^* = \kappa(y) (1 + iy) \frac{\pi}{m_N \Gamma_N} \delta(a_1^2 - m_N^2), \quad (15)$$

with  $y = \Delta m_N / \Gamma_N$  and  $\kappa(y) = 1/(1 + y^2)$ .

# LNV $B_A^{\{0,\mp\}} \rightarrow B_B^{\{\pm,0\}} P^\pm \ell_\alpha^\mp \ell_\beta^\mp$ Decays



We obtain the amplitude for the specific decay mode  
 $B_A^{\{0,+}\}(p_A) \rightarrow B_B^{\{-,0}\}(p_B) P^-(p_P) \ell_\alpha^+(p_1) \ell_\beta^+(p_2)$

$$\mathcal{M}_{\text{LNV}} = G p_P^\nu \sum_i U_{\alpha i}^* U_{\beta i}^* m_i H^\mu(p_B, p_A) \left( \ell_{\mu\nu}^{\text{LNV}}(p_1, p_2) P_{1i} + \ell_{\nu\mu}^{\text{LNV}}(p_1, p_2) P_{2i} \right), \quad (16)$$

where the leptonic part in this case is written as

$$\ell_{\mu\nu}^{\text{LNV}}(p_1, p_2) \equiv \bar{u}(p_1) \gamma_\mu \gamma_\nu (1 + \gamma_5) v(p_2), \quad (17)$$

We used  $\ell_{\mu\nu}^{\text{LNV}}(p_2, p_1) = -\ell_{\nu\mu}^{\text{LNV}}(p_1, p_2)$ , obtained by applying charge-conjugation relations.

# Detector length considerations

- We take into account the probability that the on-shell neutrino will decay inside the detector.
- The detector consideration will directly affect the constraint on the branching fraction (and consequently to the heavy-light mixings), mainly for neutrinos with masses smaller than  $\sim 500$  MeV (due to the large values of their lifetimes).

This effect is incorporated in the following probability weight:

$$P_\nu = 1 - \text{Exp} \left( -L_{\text{det}} \Gamma_N \frac{m_N}{|p_N|} \right), \quad (18)$$

where  $L_{\text{det}}$  is the detector length,  $\Gamma_N$  the total decay width of the heavy neutrino and  $p_N$  its three-momentum in the laboratory frame of the decaying baryon.

Therefore, the differential decay rate can be written as:

$$d\Gamma = P_\nu |\bar{\mathcal{M}}|^2 dPS, \quad (19)$$

being  $dPS$  the corresponding Phase Space integral according to the transition.

# Considerations for calculation

- For the estimations of the form factors, we use those obtained in the light-front model.
- The neutrino decay width is taken as dependent on
  - the mass of the heavy neutrinos,
  - the heavy-light mixing parameters,
  - and the corresponding open channels.
- In order to simplify the numerical evaluation, we take the universal coupling assumption, i.e., we consider  $|U_{eN}| = |U_{\mu N}| = |U_{\tau N}|$ .
- The exclusion regions were obtained under the assumption of an upper limit for the branching fraction of the four-body  $\Lambda_b$  decays to be  $\text{BR} \leq \mathcal{O}(10^{-8})$ , motivated by the expected sensitivity to the process  $\Lambda_b \rightarrow \Lambda_c(p)\pi\mu\mu$  at CMS and LHCb experiments <sup>4</sup>.

---

<sup>4</sup>Jhovanny Mejia-Guisao, et al. Exploring GeV-scale Majorana neutrinos in lepton-number-violating  $\Lambda_b^0$  baryon decays. *Phys. Rev. D* 96 (2017) 1, 015039