

# Energy and centrality dependence of thermodynamical observables from multiplicity in Pb+Pb and Au+Au collisions

## Content

Using a statistical model, we analyze published multiplicity distributions for identified hadrons produced in heavy ion collisions in the energy range from 2.7-200 GeV. Our analysis enables the prediction of the multiplicity distributions for multi-strangeness hadrons that have not yet been measured in lowerenergy experiments. Furthermore, we obtain freeze-out parameters, including temperature, baryon chemical and strangeness potentials, the strangeness suppression factor, and the system radius, as functions of centrality and collision energy. Additionally, we computed and discussed the Skewness and its behavior at lower collision energies, highlighting the importance of the freeze-out parameters in determining the liquid-gas phase transition in nuclear matter.

## Summary

Using a statistical model, we analyze published multiplicity distributions for identified hadrons produced in heavy ion collisions in the energy range from 2.7-200 GeV. Our analysis enables the prediction of the multiplicity distributions for multi-strangeness hadrons that have not yet been measured in lowerenergy experiments. Furthermore, we obtain freeze-out parameters, including temperature, baryon chemical and strangeness potentials, the strangeness suppression factor, and the system radius, as functions of centrality and collision energy. Additionally, we computed and discussed the Skewness and its behavior at lower collision energies, highlighting the importance of the freeze-out parameters in determining the liquid-gas phase transition in nuclear matter.

**Primary author(s)** : REYES RODRIGUEZ, Jose Francisco (Instituto de Ciencias Nucleares UNAM)

**Co-author(s)** : Dr. CUAUTLE, Eleazar (ICN-UNAM); CUAUTLE FLORES, Eleazar (ICN-UNAM)

**Presenter(s)** : REYES RODRIGUEZ, Jose Francisco (Instituto de Ciencias Nucleares UNAM)