

Effect of K^* meson magnetic dipole moment on the $e^+e^- \rightarrow K^+K^-\pi^0\pi^0$ cross section

Content

We explore the sensitivity of the $e^+e^- \rightarrow K^+K^-2\pi^0$ cross section to the magnetic dipole moment (MDM) of the K^* meson. We describe the $\gamma^* \rightarrow 2K^* \rightarrow 2K^0\pi^0$ vertex using a vector meson dominance model, including the $\rho \rightarrow \pi\pi$ transition. We emphasize the need for higher precision data to provide a first data-driven determination of this parameter to confront it with theoretical predictions.

Summary

We explore the sensitivity of the $e^+e^- \rightarrow K^+K^-2\pi^0$ cross section to the magnetic dipole moment (MDM) of the K^* meson. We describe the $\gamma^* \rightarrow 2K^* \rightarrow 2K^0\pi^0$ vertex using a vector meson dominance model, including the $\rho \rightarrow \pi\pi$ transition. We emphasize the need for higher precision data to provide a first data-driven determination of this parameter to confront it with theoretical predictions.

Primary author(s) : Mr. JIMÉNEZ PÉREZ, Luis Alberto (Instituto de Física)

Co-author(s) : Dr. TOLEDO, Genaro (Instituto de Física, UNAM); Mr. ROJAS RAMOS, Antonio (Instituto de Física UNAM)

Presenter(s) : Mr. JIMÉNEZ PÉREZ, Luis Alberto (Instituto de Física)