



Contribution ID : 463

Type : Oral

## Simultaneous H.E.S.S. and Chandra observations of Sgr A\* during an X-ray flare

Wednesday, 4 July 2007 09:06 (0:12)

### Abstract content

The rapidly varying non-thermal X-ray emission observed from Sgr A points to particle acceleration taking place close to the supermassive black hole. The TeV gamma-ray source HESS J1745-290 is coincident with Sgr A and may be closely related to the X-ray emission. Simultaneous X-ray and TeV observations are required to elucidate the relationship between these two objects. Here we report on joint H.E.S.S./Chandra observations in July 2005, during which an X-ray flare was detected.

### If this paper is presented for a collaboration, please specify the collaboration

H.E.S.S. Collaboration

### Summary

### Reference

Proceedings of the 30th International Cosmic Ray Conference; Rogelio Caballero, Juan Carlos D'Olivo, Gustavo Medina-Tanco, Lukas Nellen, Federico A. Sánchez, José F. Valdés-Galicia (eds.); Universidad Nacional Autónoma de México, Mexico City, Mexico, 2008; Vol. 2 (OG part 1), pages 633-636

**Primary author(s) :** Dr. HINTON, Jim (University of Leeds)

**Co-author(s) :** VIVIER, Matthieu (DAPNIA/DSM/CEA, CE Saclay); BUEHLER, Rolf (Max-Planck-Institut fuer Kernphysik, Heidelberg); PUEHLHOFER, Gerd (Landessternwarte, Universitaet Heidelberg); WAGNER, Stefan (Landessternwarte, Universitaet Heidelberg)

**Presenter(s) :** Dr. HINTON, Jim (University of Leeds)

**Session Classification :** OG 2.1, OG 2.2

**Track Classification :** OG.2.2