30th International Cosmic Ray Conference



Contribution ID : 67

Type : Oral

TeV gamma-rays from Galactic sources: Tycho's SNR, Geminga and Crab Nebula

Wednesday, 4 July 2007 11:42 (0:12)

Abstract content

The gamma-quantum spectra produced by the electronic and hadronic components of cosmic rays have similar shapes at the energies from 1GeV to 1 TeV due to the synchrotron losses of the electrons. So, the only observational possibility to discriminate between leptonic and hadronic contributions is to measure the gamma-quantum spectrum at energies higher than 1 TeV, where these two spectra are expected to be essentially different. The gamma-quantum emitting objects in our Galaxy are the supernova remnants and binary. According to the theoretical prediction about 20 Supernova Remnants should be visible in the TeV gamma-rays whereas only two were detected up to now by SHALON in northern hemisphere, namely Tycho's SNR and Geminga. The observation results of gamma-quantum sources Tycho Brage and Geminga by SHALON gamma-telescope are presented. The energy spectra of Geminga supernova remnants and Tycho's SNR $F(E_O > 0.8 \text{TeV})$ \propto E^k are found to be harder than Crab Nebula spectrum. The integral energy spectrum of Crab Nebula is well described by the single power law $I(> E_{\gamma})$ propto $E_{\gamma}^{-1.44}$ Geminga is one of the brightest sources of MeV – GeV gamma-ray. The value Geminga flux obtained by SHALON is lower than the upper limits published before. Its integral gamma-ray flux is found to be (0.48 pm0.17) times 10^{-12} at energies of > 0.8 TeV. Within the range 0.8 - 5 TeV, the integral energy spectrum is well described by the single power law $I(> E\gamma) \propto E\gamma^{-0.58\pm0.11}$. The integral gamma-ray flux of Tycho.s SNR above 0.8 TeV by SHALON was estimated as (0.52 \pm0.09)\times10^{-12}. The energy spectrum of Tycho's SNR at 0.8 - 20 TeV can be approximated by the power law I(>E γ)\proptoE^{k_ γ }, with k $\gamma = -1.00 \text{ pm 0.06}$. The expected {\pi}^O-decay gamma-quantum flux F_ $\gamma \ E_{\gamma^{-1}} extends up to ~ 30 TeV, whereas the Inverse Compton$ gamma-ray flux has a cutoff above the few TeV. So, the detection of gamma-rays at energies of ~ 10 - 40 TeV by SHALON is the evidence of hadron origin.

If this papers is presented for a collaboration, please specify the collaboration

Summary

Reference

Proceedings of the 30th International Cosmic Ray Conference; Rogelio Caballero, Juan Carlos D'Olivo, Gustavo Medina-Tanco, Lukas Nellen, Federico A. Sánchez, José F. Valdés-Galicia (eds.); Universidad Nacional Autónoma de México, Mexico City, Mexico, 2008; Vol. 2 (OG part 1), pages 543-546

Primary author(s): Dr. SINITSYNA, Vera Georgievna (Lebedev Physical Institute)

Co-author(s): NIKOLSKY, Sergey (Lebedev Physical Institute); SINITSYNA, Vera Yurievna (Lebedev Physical Institute); ARSOV, Todor (Lebedev Physical Institute); MUSIN, Feiruz (Lebedev Physical Institute); MALYSHKO, Alexandr (Lebedev Physical Institute)

Presenter(s) : SINITSYNA, Vera Yurievna (Lebedev Physical Institute)

Session Classification : OG 2.2

Track Classification : OG.2.2