



Contribution ID : 446

Type : Oral

## Measurement of Cosmic Ray Electrons with H.E.S.S.

*Wednesday, 4 July 2007 11:18 (0:12)*

### Abstract content

Due to energy losses in the interstellar medium, cosmic ray electrons at TeV energies carry information on local (within a few hundred parsecs) accelerators. However, measurements of the spectrum of the cosmic ray electrons beyond 1 TeV are extremely difficult due to the rapidly declining flux and the much more numerous background of nucleonic cosmic rays. The very large collection area of Cherenkov telescope arrays makes them promising instruments with which to measure these high energy electrons. While Cherenkov telescopes solve the problem of low fluxes of cosmic ray electrons in the TeV range, they still have to deal with the problem of distinguishing them from the nucleonic background. Here we report on first results towards a measurement of the cosmic ray electron spectrum with the High Energy Stereoscopic System (H.E.S.S.). The improved background suppression that is needed for such a measurement is achieved by an event classification with the 'Random Forest' algorithm based on decision trees.

### If this papers is presented for a collaboration, please specify the collaboration

H.E.S.S. collaboration

### Summary

### Reference

Proceedings of the 30th International Cosmic Ray Conference; Rogelio Caballero, Juan Carlos D'Olivo, Gustavo Medina-Tanco, Lukas Nellen, Federico A. Sánchez, José F. Valdés-Galicia (eds.); Universidad Nacional Autónoma de México, Mexico City, Mexico, 2008; Vol. 2 (OG part 1), pages 35-38

**Primary author(s) :** Mrs. EGBERTS, Kathrin (Max-Planck-Institut fuer Kernphysik)

**Co-author(s) :** Dr. HINTON, Jim (School of Physics and Astronomy, University of Leeds)

**Presenter(s) :** Mrs. EGBERTS, Kathrin (Max-Planck-Institut fuer Kernphysik)

**Session Classification :** OG 1.1

**Track Classification :** OG.1.1