



Contribution ID : 1280

Type : **Poster**

Multiwavelength search for the first blazars

Friday, 6 July 2007 14:45 (0:00)

Abstract content

High redshift massive protogalaxies, the sites of early star formation in the Universe, provided a suitable environment for the formation of the first supermassive black holes. Mass accretion causes the black hole to grow and become a gamma-ray blazar for those observers privileged by a line of sight parallel to the jet. Possible evolutionary sequences are explored in terms of their visibility in the millimeter, optical and gamma-ray regimes.

If this papers is presented for a collaboration, please specify the collaboration

Summary

Reference

Proceedings of the 30th International Cosmic Ray Conference; Rogelio Caballero, Juan Carlos D'Olivo, Gustavo Medina-Tanco, Lukas Nellen, Federico A. Sánchez, José F. Valdés-Galicia (eds.); Universidad Nacional Autónoma de México, Mexico City, Mexico, 2008; Vol. 3 (OG part 2), pages 1093-1096

Primary author(s) : Dr. CARRAMINANA, Alberto (INAOE)

Co-author(s) : Mr. RODRIGUEZ, Gustavo (INAOE)

Presenter(s) : Dr. CARRAMINANA, Alberto (INAOE)

Session Classification : Posters 2 + Coffee

Track Classification : OG.2.3