



Contribution ID : 129

Type : **Poster**

## Discovery of TeV Gamma-Ray Emission in the W28 Region from HESS Observations, and Multiwavelength Comparisons

*Friday, 6 July 2007 14:45 (0:00)*

### Abstract content

H.E.S.S. observations of the old-age (more than 10000 yrs; about 0.5 deg diameter) composite SNR W28 reveal TeV emission situated at its northeastern and southern boundaries. The northeastern TeV source is in an area where W28 is interacting with a dense molecular cloud, containing OH masers, local radio and X-ray peaks. The southern TeV sources are found in a region occupied by several HII regions, including the ultra- compact HII region W28-A2. Our analysis of NANTEN CO data also reveals a dense molecular cloud enveloping this southern region. Further details concerning H.E.S.S., NANTEN, and other multiwavelength results will be presented at the conference.

### If this papers is presented for a collaboration, please specify the collaboration

HESS, NANTEN

### Summary

### Reference

Proceedings of the 30th International Cosmic Ray Conference; Rogelio Caballero, Juan Carlos D'Olive, Gustavo Medina-Tanco, Lukas Nellen, Federico A. Sánchez, José F. Valdés-Galicia (eds.); Universidad Nacional Autónoma de México, Mexico City, Mexico, 2008; Vol. 2 (OG part 1), pages 563-566

**Primary author(s) :** Dr. ROWELL, Gavin (School of Chemistry and Physics, University of Adelaide)

**Co-author(s) :** Dr. BRION, Elizabeth (DAPNIA/DSM/CEA CE Saclay); Dr. REIMER, Olaf (HEPL and KIPAC, Stanford University); Dr. AHARONIAN, Felix (Dublin Institute for Advanced Studies); Dr. GORET, Philippe (DAPNIA/DSM/CEA, CE Saclay); Dr. DJANNATI-ATAI, Arache (APC, Paris University VII); Dr. MORIGUCHI, Yoshiaki (Dept. of Astrophysics, Nagoya University); Prof. FUKUI, Yasuo (Dept. of Astrophysics, Nagoya University)

**Presenter(s) :** Dr. ROWELL, Gavin (School of Chemistry and Physics, University of Adelaide)

**Session Classification :** Posters 2 + Coffee

**Track Classification :** OG.2.2