30th International Cosmic Ray Conference



Contribution ID : 332

Type : Poster

Long-Term Modulation Of Cosmic Ray Intensity In Different Energy Range

Abstract content

Galactic cosmic rays in the energy range from several hundred MeV to tens of GeV are subjected to heliospheric modulation, which charges their intensity and spectrum during 11-year cycle. Since the drift modulation processes are charge/polarity dependent, the 22-year solar magnetic field cycle is visible in cosmic ray data, e.g. in the different shapes of maxima of galactic cosmic rays intensity cycles. In this work, we have calculated the cross correlation function for three stations with different geomagnetic cut-off rigidities (R) for the last three solar cycles. We have calculated the running cross correlation function for high latitude Oulu (R < 1 Gv), mid latitude climax (R ~ 3 Gv) and equatorial Huancayo (R~3 Gv) neutron monitors. It is found that all three curves coincide feairly well with each other within 95% confidence interval,for the most of period of study. This good overall coincidence means that the general behaviour of CR modulation is similar for particles with different energies (within the energy range of neutron monitor sensitivity), even if the depth of modulation changes with particle energy.

If this papers is presented for a collaboration, please specify the collaboration

Summary

Reference

Primary author(s): Dr. MISHRA, Sampati kumar (Y.P.S.PG.College Semariya RewaM.P.)India)

Co-author(s): Dr. RASTOGI, Vinod Kumar (Departmentof Physica, Ch. Charan Singh University Meerut, (U.P)India); Prof. TIWARI, Dadan Prasad (Departmentof Physica, A.P.S. University Rewa, M.P.)India); Dr. SHRIVASTAVA, Pankaj Kumar (Govt.Model Science College Rewa(M.P)India)

Presenter(s): Dr. MISHRA, Sampati kumar (Y.P.S.PG.College Semariya RewaM.P.)India)

Session Classification : Posters 3 + Coffee

Track Classification : SH.3.2