



Galactic Cosmic Ray Modulation in the Heliosphere at Solar Maximum

O. G. MORALES-OLIVARES¹, R. A. CABALLERO-LOPEZ²

¹*ESIA, Instituto Politécnico Nacional, México*

²*Instituto de Geofísica, UNAM, 04510, México*

rogelioc@geofisica.unam.mx

Abstract: The spatial distribution of galactic cosmic rays in the heliosphere at solar maximum of cycles 21, 22 and 23 are studied, using a one dimensional model of the cosmic ray transport equation. We investigated the radial intensity gradients from 1 AU to the distant heliosphere and interpreted the data from IMP8, Voyagers 1 and 2, Pioneer 10 and balloon experiment BESS. In our model we considered three of the physical processes that affect the cosmic radiation: diffusion, convection and adiabatic energy loss. Our analysis indicates that adiabatic energy loss plays an important role in the radial distribution of galactic cosmic ray in the inner heliosphere, while in the outer region the diffusion and convection are the relevant processes.

Introduction

Over the last 30 years some space missions have been exploring the heliosphere. Most of them closed to the ecliptic plane. Pioneers 10 and 11, Voyagers 1 and 2 were launched to investigate the outer region of the heliosphere. They have set, together with IMP missions at 1 AU, the unique network for observing spatial and temporal cosmic ray variations. Voyagers are moving toward the nose of the heliosphere, Voyager 1 is at the heliolatitude of 34°N and Voyager 2 is at 24°S.

Many studies have been done using the cosmic ray data from those missions. Some of them have analyzed the relation between cosmic ray variations and solar activity cycle. In the present work we study the galactic cosmic ray gradient during the last three solar maximum periods. We cover a broad range of heliospheric distances, from 1 AU to 80 AU. In the previous work [6] was described the radial profiles in term of a simple model that took into account diffusion and convection. Those authors found a transition region between 10 and 20 AU where a sharp change in the gradient takes place. They argue that the changes in the interplanetary medium producing the modulation from solar minimum to solar maximum occurs in the outer region, related to the formation of the global merged

interaction regions. In this work we use a more realistic model that includes the adiabatic energy loss, and compare our results with those obtained in [6].

Data analysis

In this study we have used the data from the IMP 8 Goddard Medium Energy Detector (R.E. McGuire, P.I.), Pioneer 10 Cosmic Ray Telescope (F.B. McDonald, P.I.), the Voyager Cosmic Ray Subsystem (E.C. Stone, P.I.) and the high-altitude balloon experiment BESS (data from [7]). The time interval for our analysis is the last three solar maximum periods in 1981 (cycle 21), 1990 (cycle 22) and 2001 (cycle 23). The observations covered the radial distances from 1 AU to 80 AU. For the radial gradients we analyze the galactic cosmic ray (GCR) H in the energy range of 130–220 MeV and GCR He of 150–380 MeV/n.

In [5] and [6] the authors found that latitudinal intensity gradients are small between the ecliptic plane and the position of the two Voyager spacecraft. Therefore, we will not take into account the latitudinal gradients, and will concentrate in the radial intensity gradients during solar maximum epochs.

The aim of the present study is to compute the gra-

dients using a more realistic model that includes the adiabatic energy loss. This model is a one dimensional model, described in [1].

Model

The gradients are studied with the numerical solution of the cosmic ray transport equation ([8]). For the omnidirectional part of the cosmic ray distribution function, $f(\vec{r}, p, t)$, in one dimensional approximation, $f(r, p)$, this equation takes the form,

$$V \frac{\partial f}{\partial r} - \frac{1}{r^2} \frac{\partial}{\partial r} (r^2 \kappa \frac{\partial f}{\partial r}) - \frac{1}{3r^2} \frac{\partial}{\partial r} (r^2 V) \frac{\partial f}{\partial \ln p} = 0, \quad (1)$$

where V is the solar wind velocity, p is the particle momentum, and κ is the diffusion coefficient. The radial solar wind velocity is 400 km/s at solar maximum ([4]). At $r = r_b$ we impose the local instellar spectra (LIS) for H and He given in [9]. This boundary is a parameter that we change in order to fit the observations. Diffusion coefficient κ in one dimensional case is a function of r as follow:

$$\kappa(r) = \begin{cases} \kappa_0 r^a, & r \leq r_t \\ \kappa_0 \left(\frac{r}{r_t}\right)^b r_t^a, & r > r_t \end{cases} \quad (2)$$

where r_t is the position of the transition region, κ_0 , a and b are the other parameters that we change to obtained a good fit to the observations. The mesured cosmic ray intensity, j_T , with respect to kinetic energy per nucleon, T , is related to the omnidirectional distribution function f through $j_T = p^2 f$. Take into account this relationship, we calculate the radial intensity gradient between two points (at r_1 and r_2) inside the modulation region, by:

$$g_r = \frac{\ln(j_{T1}/j_{T2})}{r_2 - r_1}. \quad (3)$$

Results and discussion

Using our numerical model we made a parameter variation. Following [2], first we used a constant value of the diffusion coefficient κ . In figure 1 we present the energy spectra for H (panel

A) and for He (panel B) for the last solar maximum. Although, it is difficult to fit all the observations with a one dimensional model, we obtained a reasonable fitting with a boundary at 120 AU and $\kappa = 2.4 \times 10^{22} \beta P(\text{GV}) \text{ cm}^2/\text{s}$. This means that in principle one can reproduce the observed intensities without any transition region where the modulation conditions change. In this figure we observe that the spectra reach the adiabatic limit at low energies ($j_T \propto T$), and that is only possible with a model that considers the energy loss.

The next step is to introduce a transition region. In order to compare our results with the results in [6], we used $b = 1$ in (2). We took $a = 0$ because we want to see if we can explain the observations with the simplest diffusion coefficient. In this case the best fit is obtained with $r_t = 40$ AU, the boundary at $r_b = 150$ AU and $\kappa_0 = 1.8 \times 10^{22} \beta P(\text{GV}) \text{ cm}^2/\text{s}$. This value of r_b for the heliopause is more realistic (see [3]). In figure 2A and 2B we show the radial intensity profiles observed by those missions over the last three solar maxima. Radial gradients are shown in panels 2C and 2D. In the inner heliosphere (< 20 AU) the gradient for H is smaller than in the outer region. Those authors assumed that $j \propto r^\beta$, and from this calculated the radial gradient ($g_r = (1/j) \partial j / \partial r$). In that analysis ([6]), they considered a transition region from 10 to 20 AU, that separates the inner from the outer intensity gradients. In the inner heliosphere they obtained a similar gradient from H and He, ($10/r$ %/AU), while in the outer heliosphere the intensity gradient for H is $139/r$ %/AU and $73/r$ %/AU for He.

In our study, from the numerical solution of (1), we found different radial gradients, in particularly in the inner heliosphere. In this region we obtained an average radial gradient of ≈ 3 %/AU for H and ≈ 2.2 %/AU for He (shown in figure 2C and 2D). Our model naturally reproduces the small gradient due to the fact that it includes the adiabatic energy loss, that are more significant in the inner part of the heliosphere ([1]).

In the outer region the modulation is essentially driven by diffusion and convection, and in this case our results for the radial gradients are of the same order of those in [6]. However, some differences are clear. Beyond 40 AU the gradients calculated from our model do not follow the $1/r$ dependence,




Figure 1: Intensity spectra for H (panel A) and He (panel B) for 2000. The lines are the results from a one dimensional model described in the text.

even when $\kappa \propto r$. The reason for that yields also in the adiabatic term that modifies the form of the gradient derived from the force field model ($g_r = CV/\kappa$), used in [6].

Conclusions

We analyzed the galactic cosmic ray modulation at solar maximum using a one dimensional model that includes diffusion, convection and adiabatic energy loss. With this model we can conclude:

1. The observed radial profiles can be explained with a constant value of the diffusion coefficient, however, we get the best fit with a transition region at about 40 AU and the heliopause at 150 AU.
2. In the inner heliosphere, < 40 AU, the adiabatic energy loss is more important, and the radial intensity gradients are determined by this physical process.

3. In the outer heliosphere the radial intensity gradient can be explained mainly by diffusion and convection terms.

Acknowledgements

This work was supported by UNAM-DGAPA grant IN106105.

References

- [1] R. A. Caballero-Lopez and H. Moraal. Limitations of the Force Field Equation to Describe Cosmic Ray Modulation. *J. Geophys. Res.*, 109:A01101, doi:10.1029/2003JA010098, 2004.
- [2] R. A. Caballero-Lopez, H. Moraal, and F. B. McDonald. Galactic Cosmic Ray Modulation: Effects of the Solar Wind Termination Shock and the Heliosheath. *J. Geophys. Res.*, 109:A05105, doi:10.1029/2003JA010358, 2004.




Figure 2: Radial intensity profiles (panel A for H and panel B for He) and radial gradient (panel C for H and panel D for He). Full lines are from a one dimensional model with a transition region at 40 AU and heliopause at 150 AU. Dashed, dotted and dash-dotted lines are from [6].

- [3] D. A. Gurnett, W. S. Kurth, and E. C. Stone. The Return of the Heliospheric 2-3 KHz Radio Emission During Solar Cycle 23. *Geophys. Res. Lett.*, 30(23):8-1, doi:10.1029/2003GL018514, 2003.
- [4] D. J. McComas, H. A. Elliott, N. A. Schwadron, and et al. The three-dimensional solar wind around solar maximum. *Geophys. Res. Lett.*, 30(10):1517, doi:10.1029/2003GL017136, 2003.
- [5] F. B. McDonald. Cosmic Ray Modulation in the Heliosphere. A Phenomenological Study. *Space Sci. Rev.*, 83:33-50, 1998.
- [6] F. B. McDonald, Z. Fujii, B. Heikkila, and N. Lal. The Radial Distribution of Cosmic Rays in the Heliosphere at Solar Maximum. *Adv. Space Res.*, 32(4):633-638, 2003.
- [7] Z. D. Myers et al. Cosmic ray ^3He and ^4He spectra from BESS 98. In *28th Inter. Cosmic Ray Conf.*, volume 4, pages 1805-1808, 2003.
- [8] E. N. Parker. The passage of energetic particles through interplanetary space. *Planet. Space Sci.*, 13:9-13, 1965.
- [9] W. R. Webber and L. A. Lockwood. Voyager and Pioneer Spacecraft Measurement of Cosmic Ray Intensities in the Outer Heliosphere: Toward a New Paradigm for Understanding the Global Modulation Process: 1. Minimum Solar Modulation (1987 and 1997). *J. Geophys. Res.*, 106(12):29323-29332, 2001.