



Contribution ID : 807

Type : **Poster**

## The effect of a modified Parker field on the modulation of the galactic cosmic rays

*Monday, 9 July 2007 14:45 (0:00)*

### Abstract content

The effects on the galactic cosmic ray modulation of a Parker field modified by a latitudinal-dependent solar wind speed are numerically investigated. The calculation of the propagation of the galactic cosmic ray is made by solving a coupled set of the stochastic differential equations which is equivalent to the so-called diffusion convection partial differential equation. The stochastic numerical code adapted for the wavy heliospheric current sheet in a Parker field has been developed by Miyake and Yanagita [1]. The 3D code has developed into a Parker field modified by a latitudinal-dependent solar wind speed. The differences between the effect of a standard Parker field and a modified Parker field on the solar modulation of the galactic cosmic ray will be presented.

[1] Miyake, S. and Yanagita, S., Proc. 29th ICRC, 2, 203-206, 2005.

**If this papers is presented for a collaboration, please specify the collaboration**

### Summary

### Reference

Proceedings of the 30th International Cosmic Ray Conference; Rogelio Caballero, Juan Carlos D'Olivo, Gustavo Medina-Tanco, Lukas Nellen, Federico A. Sánchez, José F. Valdés-Galicia (eds.); Universidad Nacional Autónoma de México, Mexico City, Mexico, 2008; Vol. 1 (SH), pages 445-448

**Primary author(s) :** Ms. MIYAKE, Shoko (Faculty of Science, Ibaraki University)

**Co-author(s) :** Prof. YANAGITA, Shohei (Faculty of Science, Ibaraki University)

**Presenter(s) :** Ms. MIYAKE, Shoko (Faculty of Science, Ibaraki University)

**Session Classification :** Posters 3 + Coffee

**Track Classification :** SH.3.1