



Contribution ID : 851

Type : **Poster**

Search for Signatures of Extra-Terrestrial Neutrinos with a Multipole Analysis of the AMANDA-II Sky Map

Monday, 9 July 2007 14:45 (0:00)

Abstract content

In this analysis 3329 neutrino events detected by AMANDA-II during the years 2000-2003 are analysed for anisotropies or unexpected structures in their arrival direction. The structures could arise due to the presence of a signal from many weak and therefore unresolved cosmic neutrino sources, extended sources or a few brighter sources.

For the first time a technique, well known from the analysis of the cosmic micro-wave background, is applied to AMANDA-II data. The sky-distribution of arrival directions (sky-map) is expanded in a series of spherical harmonics and the power in each multipole moment is calculated. The application of this technique is challenging because of small event statistics and a non-uniform acceptance of the detector. However, the analysis is a model-independent probe of unexpected deviations of the sky-map from the expectation for atmospheric neutrinos. Compared to previous AMANDA-II analyses, it provides a new complementary approach, in particular in the search for very weak individual astro-physical sources.

The experimentally measured angular power spectrum agrees well with the expectation for a sky-map of purely atmospheric neutrinos and no excess from extra-terrestrial sources is found. Statistical errors as well as systematic errors related to the uncertainty of the angular distribution of the atmospheric neutrinos are quantified using the Feldman-Cousins unified approach.

Limits for contributions from extra-terrestrial sources to the sky-map are derived as function of the average source strength and the spectral index of the energy spectrum for different sky-distributions: weak sources isotropically distributed in the northern sky, sources located in the galactic and super-galactic plane.

The tested average flux per source varies between $\phi = 5 \cdot 10^{-13} \text{ cm}^{-2} \text{ s}^{-1}$ and $\phi = 5 \cdot 10^{-11} \text{ cm}^{-2} \text{ s}^{-1}$ at the earth, assuming an E^{-2} power spectrum in the sensitive energy range between 1.6 TeV and 1.6 PeV. The number of such sources in the sky can be limited at 90% C.L. to be less than 3000 for the weakest and less than 30 for the strongest case.

If this papers is presented for a collaboration, please specify the collaboration

IceCube

Summary

Reference

Proceedings of the 30th International Cosmic Ray Conference; Rogelio Caballero, Juan Carlos D'Olivo, Gustavo Medina-Tanco, Lukas Nellen, Federico A. Sánchez, José F. Valdés-Galicia (eds.);

Primary author(s) : HÜLB, Jan-Patrick (RWTH Aachen University)

Co-author(s) : WIEBUSCH, Christopher (RWTH Aachen University)

Presenter(s) : HÜLB, Jan-Patrick (RWTH Aachen University)

Session Classification : Posters 3 + Coffee

Track Classification : HE.2.3