



Evidences for the Influence of the Earth's Magnetic Field on EAS Muon Component

A.G. BOGDANOV¹, R.P. KOKOULIN¹, A.A. PETRUKHIN¹, O. SAAVEDRA², V.V. SHUTENKO¹, I.I. YASHIN¹

¹Moscow Engineering Physics Institute, 115409 Moscow, Russia

²Dipartimento di Fisica Generale dell'Università di Torino, 10125, Torino, Italy

IYashin@mephi.ru

Abstract: The dependence of the intensity of muon bundles detected at the Earth's surface on the angle between muon arrival direction and geomagnetic field vector (pitch angle) has been analysed. It is found that muon bundle intensity decreases with the increase of the transverse component of the magnetic field in comparison with calculations performed under assumption of azimuthal uniformity of the flux, the effect being enhanced with the increase of zenith angle. Comparison with CORSIKA-based simulations has shown that this effect is explained by a distortion of EAS muon lateral distribution functions caused by propagation of the shower in the Earth's magnetic field. Another effect - appearance of a coplanar component in directions of EAS muons in a plane defined by shower axis and Lorentz force vector - has been also observed.

Introduction

The effects of the influence of the Earth's magnetic field (EMF) on ground level muon flux characteristics are well known. For example, the azimuth asymmetry of low energy muon intensity related with primary particles geomagnetic cut-off is observed [1]. The influence of the magnetic field on the trajectories of secondary particles (muons) modifies the observed value of their charge ratio; the effect is the most significant for East-West direction at low geomagnetic latitudes near horizon [2]. Considerably less studied is the influence of EMF on multi-muon events. Experimental indications for a destruction of the axial symmetry of muon lateral distribution function (LDF) were found in giant air showers at large zenith angles [3-4]. Though theoretically the EMF influence on EAS muon component was considered by several authors [5-6], the quantitative experimental data on these effects are practically absent. In this paper, the first results of the analysis of EMF influence on muon bundle characteristics measured by means of coordinate-tracking detector DECOR [7] are presented. The setup is

located in Moscow. The absolute value of the EMF induction vector $B_0 = 52 \mu\text{T}$ [8]; the declination is about 9° East; the inclination is equal to 71° (that is, the zenith angle of EMF vector $\theta = 19^\circ$).

Angular Dependence of Muon Bundle Intensity

As it was noticed earlier [6], in a small-angle approximation the magnetic displacement of muons in a plane orthogonal to the shower axis is proportional to the transverse component of the magnetic field, inversely proportional to the particle momentum, and directly proportional to the squared geometrical path (that is, nearly proportional to the squared zenith angle secant). As a result, at large zenith angles low energy muons are swept out to shower periphery, the particles are separated in charge sign and momentum. Axial symmetry of the shower is destroyed, and muon LDF becomes "8-shaped" with the main axis parallel to the Lorentz force vector. Muon density in the central part of the shower considerably decreases.




Figure 1: The ratio of the observed number of muon bundles to the expected one as a function of azimuth angle (left) and of the pitch angle sinus (right)

When muon bundles are detected with a small-size setup (smaller than typical sizes of the shower), the event rate is determined primarily by muon density near the shower axis [9]. In this case, the EMF influence leads to the decrease of the bundle intensity, and, if the magnetic field vector is not vertical, for a fixed zenith angle the dependence of the event rate on the azimuth angle must appear. In order to explore this effect experimentally, we have used the data on muon bundles obtained by means of the coordinate-tracking detector DECOR in series of 2004-2005 measurements (about 4200 hours "live" registration time). The side part of DECOR [7] has a total area of 70 m² and consists of 8 supermodules; each of them includes 8 vertical planes of plastic streamer tube chambers. Charged particles are registered by means of external strip readout system in two orthogonal views, that allows reconstruct particle tracks in space. To ensure high statistics of the data, events with 3 muons detected in three different supermodules (the minimal number determined by trigger conditions) were selected. Total number of events in four zenith angle intervals (35°-45°, 45°-55°, 55°-65°, and 65°-75°) amounted to about 150 thousands.

The expected number of events for different zenith and azimuth angle bins was calculated taking into account the effective setup area, registration conditions and selection criteria. In calculations, a power dependence of the flux of incident events with local muon density D was assumed [9]:

$$dF/dD \sim D^{-(\beta+1)} \cos^\alpha \theta; \quad (1)$$

the parameters ($\alpha \sim 4.5$; $\beta \sim 1.95 - 2.10$, slightly increasing with zenith angle) were estimated from the experimental distributions of muon bundle characteristics. The expected event number was computed under assumption of azimuthal uniformity of the flux; then the total number of events was normalized to the observed one for every zenith angle interval. In Fig.1 (left panel), the ratio of the measured and expected intensity of muon bundles is plotted as a function of arrival azimuth angle in the laboratory frame (direction to geomagnetic South corresponds to $\varphi = 26^\circ$). As it is seen from the figure, this ratio is far not uniform. In the right panel, the same ratio is shown as a function of sinus of the angle between muon bundle direction and EMF vector (pitch angle γ). In this case, the dependence becomes regular and




Figure 2: Scatter diagrams of relative track directions in muon bundles; $\theta = 55^\circ - 65^\circ$; different azimuth angle intervals. Ellipses show 2σ -contours of 2D-Gaussian fits; solid lines: main axes of the ellipses; dotted lines: directions of Lorentz force vector

monotonous; the slope is increasing with the increase of zenith angle. Solid curves in the right part of Fig.1 represent the results of calculations of the expected muon bundle intensity (also normalized to the observed number of events for every zenith angle bin) taking into account the Earth's magnetic field, obtained on a basis of muon LDFs simulated by means of the CORSIKA code [10] (version 6.502) for fixed zenith angles and different values of the pitch angle. Comparison of the curves and the data exhibits a good qualitative (and a reasonable quantitative) agreement and confirms that the observed angular dependence of the event intensity is explained by the changes of EAS muon LDF under the influence of the geomagnetic field.

Coplanarity of Tracks in the Bundles

Apart from stochastic processes of the angular deflection of muon trajectory from the direction of the primary particle (transverse momenta in

hadronic interactions, parent meson decay kinematics, multiple Coulomb scattering in air), there exists a regular component of the deflection caused by the Earth's magnetic field. An equivalent transverse momentum may be estimated as

$$p_{\text{mag}} = 0.3 B_0 \sin\gamma H_0 \sec\theta. \quad (2)$$

Here H_0 is the effective muon production altitude; usual system of units (GeV/c, T, m) is used. A typical value of this momentum in conditions of the present experiment for 60° zenith angle equals to about 0.3 - 0.5 GeV/c, which is comparable to the average transverse momentum for meson production; contribution of multiple scattering is several times less.

Muon bundles detected by a small-size setup contain muons initially emitted in different directions and slightly turned (differently, depending on the momentum and sign) by the magnetic

field. Hence, in the directions of particles that hit the detector a kind of alignment (coplanarity) in a plane defined by EAS axis and Lorentz force vector must appear. Remarkably, for EAS arrival directions close to the geomagnetic meridian (from North or South) the Lorentz force vector is horizontal; for EAS arriving from East or West the regular component of the transverse momenta must be inclined relative to horizon.

To search for this effect, we have selected muon bundle events with four (and only four) quasi-parallel particles detected in four different super-modules of DECOR. The last condition was necessary for unambiguous reconstruction of individual particle tracks in space. The events with zenith angles 55° - 65° in four narrow (20° -wide) sectors of azimuth angle (nearly equivalent from the view-point of track registration and geometry reconstruction, but differently oriented relative to geomagnetic meridian) were analyzed.

For every muon bundle, the average direction was determined, and then deviations of individual tracks from this average bundle vector in a vertical plane ($\Delta\theta$) and across it ($\Delta\chi = \sin\theta \cdot \Delta\phi$) were calculated. In Fig.2, scatter diagrams of the track directions in ($\Delta\chi$, $\Delta\theta$) coordinates for muon bundles arriving from four different azimuth angle intervals are presented. Ellipses in the figures correspond to 2σ -contours of 2D-Gaussian fit of the points distribution; solid lines indicate the main axes of the ellipses, dotted ones show the directions of the Lorentz force (average for every data sample). As it is seen from the figure, at the background of measurement errors and random scattering factors a regular component (with rms-value about 0.4° - 0.5°) aligned with the Lorentz force vector is really observed in muon track arrival directions.

Conclusion

Analysis of muon bundles registered in coordinate-tracking detector DECOR has shown that the event intensity significantly decreases with the sinus of the angle between the muon bundle arrival direction and geomagnetic field vector; this effect is enhanced with the increase of zenith angle. Comparison with CORSIKA-based simula-

tions shows that this phenomenon is explained by the distortion of muon LDF resulting from propagation of EAS muons in the geomagnetic field. A reasonable agreement between calculations and data may be considered as a validation of CORSIKA treatment of EMF influence on EAS muon component. A new phenomenon - coplanarity of muon bundle tracks in a plane determined by the Lorentz force vector and EAS axis - has been found.

Acknowledgments

The work was supported by Federal Agency for Education, Federal Agency for Science and Innovations and RFBR (grant 06-02-17000).

References

- [1] Pham Ngoc Diep et al. Nucl. Phys. B, 678 (2004) 3.
- [2] O.C.Allkofer et al. Nucl.Phys.B, 259 (1985) 1.
- [3] A.M. Hillas et al. Proc. 11th ICRC, Budapest, 3 (1969) 533.
- [4] L. Nellen et al. Proc. 29th ICRC, Pune, 7 (2005) 183.
- [5] L.G. Dedenko et al. Proc. 24th ICRC, Roma, 1 (1995) 309.
- [6] M. Ave et al. Astropart. Phys., 14 (2000) 19.
- [7] M.B. Amelchakov et al. Proc. 27th ICRC, Hamburg, 3 (2001) 1267.
- [8] <http://www.ngdc.noaa.gov/ngdc.html>
- [9] N.S. Barbashina et al. Nucl. Phys. B (Proc. Suppl.), 165 (2007) 317.
- [10] D. Heck et al. <http://www-ik.fzk.de/corsika>.