



Contribution ID : 324

Type : Oral

## The shower size spectrum reconstructed with KASCADE-Grande

Friday, 6 July 2007 12:30 (0:12)

### Abstract content

The Grande array as main part of the KASCADE-Grande experiment consists of 37x10 sqm scintillation detectors spread over an area of 700x700 sqm. Grande enables triggers and reconstruction of primary cosmic rays in the energy range of  $\sim 30$  PeV to 1 EeV. This contribution discusses the reconstruction of the shower size spectrum, i.e. the total number of charged particles. The KASCADE-Grande set-up allows, for a subsample of the registered showers, detailed comparisons of the data with measurements of the original KASCADE array (252 detectors, 490 sqm sensitive area spread over 200x200 sqm) on an event-by-event basis. The obtained reconstruction accuracy of  $N_{\text{charged}}$  will be presented as well as the lateral charged particle distributions and the absorption length of the charged particle component. In the overlapping size region the resulting shower size spectra for different zenith angular ranges will be compared with the KASCADE size spectra.

### If this papers is presented for a collaboration, please specify the collaboration

KASCADE-Grande

### Summary

### Reference

Proceedings of the 30th International Cosmic Ray Conference; Rogelio Caballero, Juan Carlos D'Olivo, Gustavo Medina-Tanco, Lukas Nellen, Federico A. Sánchez, José F. Valdés-Galicia (eds.); Universidad Nacional Autónoma de México, Mexico City, Mexico, 2008; Vol. 4 (HE part 1), pages 195-198

**Primary author(s) :** Dr. DI PIERRO, Federico (University of Torino)

**Presenter(s) :** Dr. DI PIERRO, Federico (University of Torino)

**Session Classification :** HE 1.3.A

**Track Classification :** HE.1.3.A