



Contribution ID : 1097

Type : Oral

## Atmospheric neutrino oscillation in Super-Kamiokande

*Saturday, 7 July 2007 12:41 (0:12)*

### Abstract content

The Super-Kamiokande started observation in April of 1996 and continued the data taking for five years of initial running period (SK-I) till the maintenance in July of 2001. The Super-Kamiokande continued the data taking with the half PMT density in the second period (SK-II) from resuming in December of 2002 to shutdown for the reconstruction to return the PMT density in October 2005. Using the atmospheric neutrino data of SK-I and SK-II, various types of oscillation analyses have been performed. The results from the  $\nu_{\mu} \leftrightarrow \nu_{\tau}$  2 flavor oscillation analysis and the 3 flavor oscillation analysis with the aim of searching for a non-zero 1-3 mixing will be presented. The results from an oscillation analysis including the sub-dominant oscillation effects driven by the solar oscillation terms ( $\theta_{12}$  and  $\Delta m_{21}^2$ ) will also be reported.

### If this papers is presented for a collaboration, please specify the collaboration

The Super-Kamiokande Collaboration

### Summary

### Reference

Proceedings of the 30th International Cosmic Ray Conference; Rogelio Caballero, Juan Carlos D'Olivo, Gustavo Medina-Tanco, Lukas Nellen, Federico A. Sánchez, José F. Valdés-Galicia (eds.); Universidad Nacional Autónoma de México, Mexico City, Mexico, 2008; Vol. 5 (HE part 2), pages 1319-1322

**Primary author(s) :** Ms. TAKENAGA, Yumiko (ICRR, University of Tokyo)

**Presenter(s) :** Ms. TAKENAGA, Yumiko (ICRR, University of Tokyo)

**Session Classification :** HE 2.2

**Track Classification :** HE.2.2