



Contribution ID : 185

Type : Oral

## A conservative dark matter annihilation bound from the diffuse gamma ray flux

*Tuesday, 10 July 2007 08:54 (0:12)*

### Abstract content

We discuss the general observational upper limits on the total annihilation rate of dark matter derived from the diffuse gamma ray background measured by the EGRET satellite. We assume that the dark matter annihilates at tree level in the least detectable final states in the Standard Model, namely neutrinos. Any other decay channel would lead to stronger constraints. Electroweak jet cascading imply that, for masses above  $\sim 100$  GeV, interesting bounds apply which are much more stringent than well-known general bounds as the unitarity one and comparable or only a few times weaker than the bounds recently derived from atmospheric neutrino data. We briefly discuss the perspectives of improvement of these constraints on the light of forthcoming GLAST satellite data.

**If this paper is presented for a collaboration, please specify the collaboration**

### Summary

### Reference

Proceedings of the 30th International Cosmic Ray Conference; Rogelio Caballero, Juan Carlos D'Olivo, Gustavo Medina-Tanco, Lukas Nellen, Federico A. Sánchez, José F. Valdés-Galicia (eds.); Universidad Nacional Autónoma de México, Mexico City, Mexico, 2008; Vol. 4 (HE part 1), pages 701-704

**Primary author(s) :** Dr. SERPICO, Pasquale Dario (Fermilab)

**Co-author(s) :** Prof. KACHELRIESS, Michael (Department of Physics, NTNU - Trondheim, Norway)

**Presenter(s) :** Dr. SERPICO, Pasquale Dario (Fermilab)

**Session Classification :** HE 3.3

**Track Classification :** HE.3.3