



Contribution ID : **1249**

Type : **Poster**

## The end of the Galactic spectrum

*Wednesday, 4 July 2007 14:45 (0:00)*

### Abstract content

We use two complementary models, diffusion and explicit particle propagation, to analyze the end of the Galactic cosmic ray spectrum and its mixing with the extragalactic cosmic ray flux. The extragalactic cosmic ray flux is estimated by comparing Fly's Eye, HiRes and AGASA spectra with Galactic spectrum expectations for different injection models, populations of Galactic accelerators and magnetized Halo structures.

**If this papers is presented for a collaboration, please specify the collaboration**

### Summary

### Reference

Proceedings of the 30th International Cosmic Ray Conference; Rogelio Caballero, Juan Carlos D'Olivo, Gustavo Medina-Tanco, Lukas Nellen, Federico A. Sánchez, José F. Valdés-Galicia (eds.); Universidad Nacional Autónoma de México, Mexico City, Mexico, 2008; Vol. 2 (OG part 1), pages 219-222

**Primary author(s) :** DE DONATO, Cinzia (Physics Department of the University and INFN, Milano, Italy)

**Co-author(s) :** Dr. MEDINA TANCO, Gustavo (Dep. Altas Energías, Inst. de Ciencias Nucleares, Universidad Nacional Autónoma de México, México DF, CP 04510)

**Presenter(s) :** DE DONATO, Cinzia (Physics Department of the University and INFN, Milano, Italy)

**Session Classification :** Posters 1 + Coffee

**Track Classification :** OG.1.3