30th International Cosmic Ray Conference



Contribution ID : 473

Type : Poster

Analytical solutions of the Moliere series terms of higher orders for multiple Coulomb scattering

Wednesday, 4 July 2007 14:45 (0:00)

Abstract content

General higher terms of Moliere series are solved analytically, in Moliere- Heisenberg definite integral and/or Goldstein series. The terms of higher orders up to n=6 are practically obtained. Applicable region of Moliere series is extended to shorter depths of penetration down to B=5 by the results.

If this papers is presented for a collaboration, please specify the collaboration

Summary

Reference

Proceedings of the 30th International Cosmic Ray Conference; Rogelio Caballero, Juan Carlos D'Olivo, Gustavo Medina-Tanco, Lukas Nellen, Federico A. Sánchez, José F. Valdés-Galicia (eds.); Universidad Nacional Autónoma de México, Mexico City, Mexico, 2008; Vol. 4 (HE part 1), pages 149-152

Primary author(s) : Prof. NAKATSUKA, Takao (Okayama Shoka University)

Co-author(s) : Dr. OKEI, Kazuhide (Graduate School of Natural Science and Technology, Okayama University); Dr. TAKAHASHI, Naoya (Graduate School of Natural Science and Technology, Okayama University)

Presenter(s): Prof. NAKATSUKA, Takao (Okayama Shoka University)

Session Classification : Posters 1 + Coffee

Track Classification : HE.1.2.B