



Contribution ID : 474

Type : **Poster**

## **Accuracy of numerical functional transforms applied to derive Moliere series terms and comparison with analytical results**

*Wednesday, 4 July 2007 14:45 (0:00)*

### **Abstract content**

Accuracies of numerical Fourier and Hankel transforms are examined with the Takahasi-Mori theory of error evaluation. The higher Moliere terms both for spatial and projected distributions derived by these methods agree very well with those derived analytically. The methods will be valuable to solve other transport problems concerning fast charged particles.

**If this papers is presented for a collaboration, please specify the collaboration**

### **Summary**

### **Reference**

Proceedings of the 30th International Cosmic Ray Conference; Rogelio Caballero, Juan Carlos D'Olive, Gustavo Medina-Tanco, Lukas Nellen, Federico A. Sánchez, José F. Valdés-Galicia (eds.); Universidad Nacional Autónoma de México, Mexico City, Mexico, 2008; Vol. 4 (HE part 1), pages 153-156

**Primary author(s) :** Dr. TAKAHASHI, Naoya (Graduate School of Natural Science and Technology, Okayama University)

**Co-author(s) :** Dr. OKEI, Kazuhide (Graduate School of Natural Science and Technology, Okayama University); Prof. NAKATSUKA, Takao (Okayama Shoka University)

**Presenter(s) :** Prof. NAKATSUKA, Takao (Okayama Shoka University)

**Session Classification :** Posters 1 + Coffee

**Track Classification :** HE.1.2.B