



Contribution ID : 759

Type : Oral

## Particle Acceleration and Turbulence in Cosmic Ray Shocks: Possible Pathways beyond the Bohm limit

### Abstract content

We suggest a physical mechanism whereby the acceleration time of cosmic rays by shock waves can be significantly reduced. This creates the possibility of particle acceleration beyond the knee energy at  $10^{15}$  eV. The acceleration results from a nonlinear modification of the flow ahead of the shock supported by particles already accelerated to the knee momentum. The particles gain energy by bouncing off converging magnetic irregularities frozen into the flow in the shock precursor and not so much by re-crossing the shock itself. The acceleration rate is thus determined by the gradient of the flow velocity and turns out to be formally independent of the particle mean free path. The velocity gradient is, in turn, set by the knee-particles. The acceleration rate of particles above the knee does not decrease with energy, unlike in the linear acceleration regime. The knee forms because particles above it are effectively confined to the shock while they are within limited domains in the momentum space, while other particles fall into “loss-islands”, similar to the “loss-cone” of magnetic traps. This also maintains the steep velocity gradient and high acceleration rate.

**If this papers is presented for a collaboration, please specify the collaboration**

### Summary

### Reference

**Primary author(s) :** Dr. MALKOV, Mikhail (University of California San Diego)

**Co-author(s) :** Prof. DIAMOND, Patrick (University of California San Diego)

**Presenter(s) :** Dr. MALKOV, Mikhail (University of California San Diego)

**Session Classification :** OG 1.4

**Track Classification :** OG.1.4