



Contribution ID : 391

Type : Oral

## On the GeV-TeV connection of Galactic Gamma-Ray Sources, II. VHE gamma-ray sources without EGRET-detected counterparts and vice versa

*Saturday, 7 July 2007 12:29 (0:12)*

### Abstract content

Recent observations by atmospheric Cherenkov telescopes such as H.E.S.S., and MAGIC have revealed a large number of new sources of very-high-energy (VHE) gamma-rays above 100 GeV, mostly concentrated along the Galactic plane. At lower energies (100 MeV - 10 GeV) the satellite based instrument EGRET revealed a population of gamma-ray sources clustering along the Galactic Plane. Here we investigate those gamma-ray sources, which currently lack a connection in their adjacent energy band. We conclude on similarities and differences among the EGRET resp. VHE gamma-ray sources, and complement the discussion with consistently determined upper limits in their proximate energy band. An interpretation will invoke characteristics of known object classes at high-energy gamma-rays, and both observational and theoretical arguments to explain situations where no counterpart can be expected in one or the other energy regimes.

**If this papers is presented for a collaboration, please specify the collaboration**

### Summary

### Reference

Proceedings of the 30th International Cosmic Ray Conference; Rogelio Caballero, Juan Carlos D'Olivo, Gustavo Medina-Tanco, Lukas Nellen, Federico A. Sánchez, José F. Valdés-Galicia (eds.); Universidad Nacional Autónoma de México, Mexico City, Mexico, 2008; Vol. 2 (OG part 1), pages 613-616

**Primary author(s) :** Dr. REIMER, Olaf (HEPL/KIPAC, Stanford University)

**Co-author(s) :** Dr. FUNK, Stefan (Kavli Institute for Particle Astrophysics and Cosmology, SLAC); Prof. TORRES, Diego F. (ICREA/IEEC-CSIC, Barcelona, Spain)

**Presenter(s) :** Prof. TORRES, Diego F. (ICREA/IEEC-CSIC, Barcelona, Spain)

**Session Classification :** OG 2.2, OG 2.1

**Track Classification :** OG.2.2