



Contribution ID : 505

Type : Oral

Photon spectra from final stages of a primordial black hole evaporation in different theoretical models

Monday, 9 July 2007 13:17 (0:12)

Abstract content

Possibilities of an experimental search for gamma-ray bursts from primordial black hole (PBH) evaporations in space are reconsidered. It is argued that the corresponding constraints which can be obtained in experiments with cosmic ray detectors strongly depend on theoretical approach used for a description of the PBH evaporation process. Predictions of several theoretical models for gamma-ray spectra from final stages of PBH life (integrated over time) are given.

If this papers is presented for a collaboration, please specify the collaboration

Summary

Reference

Proceedings of the 30th International Cosmic Ray Conference; Rogelio Caballero, Juan Carlos D'Olivo, Gustavo Medina-Tanco, Lukas Nellen, Federico A. Sánchez, José F. Valdés-Galicia (eds.); Universidad Nacional Autónoma de México, Mexico City, Mexico, 2008; Vol. 3 (OG part 2), pages 1123-1126

Primary author(s) : Prof. BUGAEV, Edgar (Institute for Nuclear Research, Moscow, Russia)

Co-author(s) : Dr. KLIMAI, Peter (Institute for Nuclear Research, Moscow, Russia); Dr. PETKOV, Valery (Institute for Nuclear Research, Moscow, Russia)

Presenter(s) : Prof. BUGAEV, Edgar (Institute for Nuclear Research, Moscow, Russia)

Session Classification : OG 2.4

Track Classification : OG.2.4